Zeitschrift:	Entomologica Basiliensia
Herausgeber:	Naturhistorisches Museum Basel, Entomologische Sammlungen
Band:	25 (2003)
Artikel:	Towards knowledge of the Indian Clytrinae (Coleoptera, Chrysomelidae)
Autor:	Medvedev, Lev N.
DOI:	https://doi.org/10.5169/seals-980880

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

# Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Entomologica Basiliensia	25	275-291	2003	ISSN 0253-24834

# Towards knowledge of the Indian Clytrinae (Coleoptera, Chrysomelidae)

by Lev N. Medvedev

Abstract. A list of Clytrinae (Chrysomelidae) from southern India in the collection of the Natural History Museum Basel (Switzerland) is given, as well as a key to the genera of southern India and keys for genera *Pantocometis* Lacordaire, 1848, *Epimela* Weise, 1903 and *Aetheodactyla* Lacordaire, 1848. Seven new species are described: *Epimela minuta* (southern India), *E. tristis* (Nepal), *Aetheodactyla orientalis* (eastern India), *A. regalini, Aspidolopha dembickyi, Aetheomorpha sculpturata*, and *Smaragdina pacholatkoi* (southern India) spp.nov. *Epimela downesii* (Baly, 1865) comb.nov. is removed from the genus *Lachnaea* Chevrolat, 1837. *Aetheodactyla flavobasalis* (Jacoby, 1908) comb.nov. is removed from the genus *Smaragdina* Chevrolat, 1837. *Aetheodactyla minor* Duvivier, 1891 is confirmed as a valid species (a new status for *Pseudoclytra plagiata* var. *minor* Duvivier, 1891). Two new synonyms are established: *Epimela downesii* (Baly, 1865) (= *Epimela indica* Duvivier, 1891 syn.nov.) and *Aetheodactyla flavobasalis* (Jacoby, 1908) (= *Aetheodactyla incisipes* L. Medvedev, 1984 syn.nov.).

Key words. Coleoptera - Chrysomelidae - Clytrinae - India - taxonomy - keys - new species

### Introduction

During my last visit to the Basel Museum of Natural History I had an opportunity to study a large and interesting body of Clytrinae material collected in southern India in the year 2002. This included 16 species, mostly represented with large series, among them 4 species new to science.

The Clytrinae of southern India are still very poorly studied and no keys exist. Because of this, I include a key to the genera of southern India in the article, as well as revisions of 3 genera: *Pantocometis* Lacordaire, 1848, *Epimela* Weise, 1903 and *Aetheodactyla* Lacordaire, 1848, also with descriptions of a few new species.

# Material of the southern Indian Clytrinae in Natural History Museum Basel (The species collected in 2002.)

### Pantocometis subfasciata Jacoby, 1908

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 7 exx.

**Remark.** The black spots on the elytra are occasionally connected along the side margin.

#### Aetheodactyla regalini sp.nov.

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 1 male.

**Remark.** The species described below.

#### L. N. MEDVEDEV

# Aspidolopha distincta Duvivier, 1891

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 8 exx.

# Aspidolopha decora Fabricius, 1801

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 6 females.

# Aspidolopha dembickyi sp.nov.

**Material examined.** Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 10 females; Tamil Nadu, Nilgiri Hills, 15 km SE Kotagiri, Kunchappanai, 17–22.V.1997, leg. Dembický & Pacholátko, 1 male.

Remark. The species described below.

# Aetheomorpha cribellata Jacoby, 1908

**Material examined.** Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 19 exx.; Karnataka, W Ghats, 20 km W Talguppa, Jog Falls, 500 m, 22–28.V.2002, leg. P. Pacholátko, 1 ex. Aedeagus – Fig. 17.

**Remarks.** This species is very variable, and is represented by the following colour forms:

#### 1. Prothorax fulvous:

- Elytra entirely fulvous 2 exx.
- Elytra with black band beyond mid-point 4 exx.
- Elytra with 2 spots in anterior half and black band beyond mid-point (typical form) 1 exx.

# 2. Prothorax black:

- Elytra entirely fulvous 2 exx.
- Elytra with black band beyond mid-point 5 exx.
- Elytra with 2 spots in anterior half (sometimes connected) and black band beyond mid-point 6 exx.

# Aetheomorpha nigropicta (Lefevre, 1891)

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 1 female.

# Aetheomorpha sculpturata sp.nov.

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 3 females.

## Remark. The species described below.

#### Aetheomorpha suturata Jacoby, 1898

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 2 exx; Karnataka, W Ghats, 20 km W Talguppa, Jog Falls, 500 m, 22–28.V.2002, leg. P. Pacholátko, 9 exx.

#### Aetheomorpha tibialis L. Medvedev, 2000

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 9 exx.

**Remark.** One male with entirely fulvous prothorax; in other specimens there are well developed black spots.

#### Aetheomorpha bifurcata L. Medvedev, 2001

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 1 male, 41 females; Karnataka, W Ghats, 20 km W Talguppa, Jog Falls, 500 m, 22–28.V.2002, leg. P. Pacholátko, 2 females.

#### Miochira lefevrei Jacoby, 1895

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 15 exx; Karnataka, W Ghats, 20 km W Talguppa, Jog Falls, 500 m, 22–28.V.2002, leg. P. Pacholátko, 5 exx.

#### Ceroclytra cornuta Jacoby, 1895

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 1 exx.; Karnataka, W Ghats, 20 km W Talguppa, Jog Falls, 500 m, 22–28.V.2002, leg. P. Pacholátko, 3 exx.

#### Smaragdina frontalis Jacoby, 1908

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 9 exx.

#### Smaragdina divisa Jacoby, 1889

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 63 exx.

## Smaragdina pacholatkoi sp.nov.

Material examined. Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 57 exx.

Remark. The species described below.

### Smaragdina nilgiriensis Jacoby, 1903

Material examined. Tamil Nadu, Nilgiri Hills, 15 km SE Kotagiri, Kunchappanai, 17–22.V.1997, leg. Dembický & Pacholátko, 1 male.

#### Abbreviations

NHMB	 Naturhistorisches Museum, Bas	sel
LM	 Author's collection (L. N. Medvedev), Mosco	w
SMNS	 Staatliches Museum für Naturkunde, Stuttga	art

# Taxonomy

## Key to genera of Clytrinae of southern India

- 1(4) Upperside or at least prothorax pubescent. First segment of tarsi more or less elongate. Pygidium not covered by elytra.
- 2(3) Propleurae not pubescent. Upperside, head, pygidium and legs with short, light pubescence and long, erect black hairs. Elytra confusedly punctate ... *Pantocometis* Lacordaire, 1848
- 4(1) Upperside not pubescent.
- 6(5) Epipleurae not pubescent.
- 7(10) Frons and vertex with dense, erect hairs. Anterior coxae divided with thin but very high lamellate prosternum. Antennal segments 6–9 distinctly transverse. Pygidium covered by elytra. Tarsi broad, 2nd segment broader than long. Body large.

- 10(7) Frons and vertex glabrous or with short, mostly adpressed pubescence. Anterior coxae contiguous, prosternum low and not seen between coxae. Tarsi narrow with segments 1 and 2 mostly longer than broad.
- 11(16) Pygidium not covered by elytra.
- 12(15) Prosternal triangle between coxa, pleural suture and anterior margin with dense adpressed publication. Elytra with well developed epipleural lobe, epipleurae almost vertical.
- 14(13) Propleurae pubescent. Apex of pygidium mostly emarginate in female. Head with pubescent stripes along eyes, sometimes connected on frons. Body narrows towards the front and the rear. Prothorax with broadly rounded hind angles, usually punctate. ..... Aspidolopha Lacordaire, 1848
- 16(11) Pygidium covered by elytra.
- 17(18) Anterior legs of male strongly elongate. Head of male not enlarged. Body black, elytra fulvous with 2 black bands. [Single species, *M. lunulata* (Fabricius, 1781).]
   Merilia Lacordaire, 1848
- 18(17) Anterior legs of male feebly elongate or not at all, in latter case body not black.

#### L. N. MEDVEDEV

- 20(19) In male, anterior legs not elongate and head not enlarged. First antennal segment feebly widened or not at all.
- 22(21) Tarsi longer, at least tarsal segment 1 narrow and elongate.
- 23(24) Hind angles of prothorax distinct, obtuse. Tibiae often comparatively thick. ..... *Physosmaragdina* L. Medvedev, 1971
- 24(23) Hind angles of prothorax broadly rounded. Tibiae thin.
- 25(26) Head of male with horn on frons. Upperside fulvous with variable dark spots. ..... *Ceroclytra* L. Medvedev, 1962

# Key to southern Indian species of the genus Pantocometis Lacordaire, 1848

- 1(4) Elytra with large subtriangular spot at humeri and with black transverse band in front of apex.
- 2(3) Elytral spots not connected along side margin. Pygidium entirely fulvous. Aedeagus, Fig. 1. Length 4.2–5.3 mm. ....... *P. subfasciata* Jacoby, 1908
- 4(1) Elytra with broad black band beyond centre, widened externally. Length 5 mm. Poorly known species. ..... *P. hirsuta* Lacordaire, 1848

### Key to southern Indian species of the genus Epimela Weise, 1903

- 3(2) Elytra without apical spot.

- Prothorax dark metallic, very narrowly fulvous margined, especially on 4(13)sides.
- Pubescence of head and prothorax adpressed. 5(12)
- 6(7)Legs entirely fulvous. Aedeagus with elongate apical triangle, more or less separated from central part (Fig. 2). Length 5.3-6 mm. Northern India: Jammu, Assam. ..... E. viridicollis (Jacoby, 1899)
- Legs dark metallic with fulvous tibiae and sometimes tarsi. 7(6)
- 8(9) Elytra unspotted or with small spot near suture beyond centre. Length 5.1-6 mm. Aedeagus as in E. trimaculata (Fabricius, 1798), apical triangle somewhat more elongate. Nepal. [Perhaps a subspecies of *E. trimaculata*. Type series studied.] ..... ..... E. nepalica Takizawa, 1987
- 9(8) Spots on elytra more developed.
- 10(11) Apex of aedeagus in form of short triangle (Fig. 3). Postmedian band on elytra reduced to small spot or absent. Labrum fulvous. Length 5.5 mm. Distribution unclear, type from "India", other specimens from "India orientalis". [syn. Diapromorpha cicticollis var. obliterata Pic, 1941] ..... E. obliterata (Pic, 1941)
- 11(10) Apex of aedeagus in form of elongate triangle (Fig. 4). Full complement of spots on elytra or humeral spot absent. Labrum dark. Length 5–6.7 mm. Southern India, Sri Lanka; Nepal records need confirmation and perhaps belong to E. nepalica. [syn. E. insularis Weise, 1903, E. dilutipes Jacoby, 1908, E. viridescens Pic, 1941, E. viridescens ab. interrupta Pic, 1941, Diapromorpha cicticollis Pic, 1941] ..... ..... E. trimaculata Fabricius, 1798
- Pubescence of head and prothorax longer and partly erect (especially in 12(5)lateral view). Antennae and labrum fulvous. Aedeagus with very elongate apical part (Fig. 5). Body small, 4.3 mm. Southern India. ..... E. minuta sp.nov.
- 13(4)Prothorax entirely dark metallic blue, without fulvous emargination. Legs entirely blackish-blue. Length 6.4 mm. Nepal. [Male unknown.] ..... E. tristis sp.nov.
- Elytra with sparse erect pubescence, especially in humeral area, more or 14(1)less regularly punctate, fulvous, usually with dark spot or band beyond centre. Prothorax with erect hairs, dark metallic, sometimes with very narrow fulvous emargination. ..... (subgenus Paraepimela L. Medvedev, 1984)
- 15(16) First abdominal sternite without smooth spot on the sides. Elytra parallelsided, about 1.7 times as long as wide. Labrum fulvous. Aedeagus, Fig. 6. Length 6.5–9 mm.



Figs 1–8. Aedeagus (d – dorsal, v – ventral, l – lateral, a – apical): 1, Pantocometis subfasciata Jacoby (d); 2, Epimela viridicollis Jacoby (d); 3, Epimela obliterata Pic (d); 4, Epimela trimaculata Fabricius (d); 5, Epimela minuta sp.nov. (d); 6, Epimela downesii Baly (d); 7, Aetheodactyla andrewesi Jacoby (v, l, a) (after TAKIZAWA 1990); 8, Aetheodactyla wallardiensis Jacoby (d, l, a) (after TAKIZAWA 1990).

16(15) First abdominal segment with smooth bare spot on the sides. Elytra elongate ovate, about 1.5 times as long as wide. Labrum dark. Length 6.3–6.8 m.
Sri Lanka. [Male unknown.] ...... E. zaitzevi L. Medvedev, 1984

# Epimela (s.str.) minuta sp.nov.

Material examined. Holotype (male): southern India, Nilgiri Hills, Moyar-Camp, 3000 ft., V. 1954, leg. Nathan (LM).



Figs 9–14. Aedeagus (d – dorsal, v – ventral, l – lateral, a – apical): 9, Aetheodactyla orientalis sp.nov. (v); 10, Aetheodactyla lateralis Lacordaire (v, l, a) (after TAKIZAWA 1990); 11, Aetheodactyla plagiata Duvivier (v, l, d); 12, Aetheodactyla minor Duvivier (v, l, d); 13, Aetheodactyla regalini sp.nov. (v, l); 14, Aetheodactyla flavobasalis Jacoby (v).

**Description.** Dark metallic green, labrum, antennae, narrow emargination of prothorax, elytra, tibiae and tarsi fulvous; on elytra humeral spot, another spot before midpoint near suture and transverse band beyond centre violaceous blue; extreme apex of suture with small, dark-brown mark (Fig. 19).

Morphologically identical with *E. trimaculata*, only head and pronotum with longer and more or less erect hairs. Aedeagus (Fig. 5) with very elongate apical triangle.

Length: 4.3 mm.

# Distribution. Southern India.

**Diagnosis.** Being very near to *E. trimaculata*, differs sharply in fulvous antennae and labrum, erect hairs of head and prothorax, trace of additional spot on apex of elytra, smaller size and different form of aedeagus.

## Epimela (s.str.) tristis sp.nov.

**Material examined.** Holotype (female). Nepal 471, Myagdi Distr., lower Marang Khola, Bim, 1150 m, 29. V. 1995, leg. Martens & Schawaller (SMNS).



Figs 15–18. Aedeagus (d – dorsal, v – ventral, l – lateral, a – apical): 15, Aetheodactyla dimidiatipennis Baly (v); 16, Aspidolopha dembickyi sp.nov. (d); 17, Aetheomorpha cribellata Jacoby (v); 18, Smaragdina pacholatkoi sp.nov. (v).

**Description.** Dark metallic green, 3 basal antennal segments and elytra fulvous, on elytra humeral and subhumeral spots and large patch in middle (connected sutural spot and postmedian band) violaceous.

Clypeus glossy, finely punctate, frons and vertex densely punctate, with adpressed pubescence. Antennae distinctly serrate from the 4th segment. Prothorax broadest before base, with adpressed pubescence, very densely punctate on sides, finely and sparsely in the middle. Elytra strongly and irregularly punctate, partly rugose, not pubescent.

Length: 6.4 mm.

Distribution. Nepal.

**Diagnosis.** Differs clearly from all species of the subgenus in its naked and smooth clypeus, the sculpture of the prothorax and entirely metallic legs.

# Epimela (Paraepimela) downesii (Baly, 1865) comb.nov.

Pantocometis downesii Baly, 1865 Lachnaea downesii: BRYANT (1923). Epimela indica (Duvivier, 1891) **syn.nov.** 

**Remarks.** This species is identical with *E. indica* (Duvivier, 1891) syn.nov. It is really surprising that BRYANT (1923) includes it in the Mediterranean genus *Lachnaea* Chevrolat, 1837.

# A key to Indian species of the genus Aetheodactyla Lacordaire, 1848

- 1(6) Upperside entirely fulvous. Hind tibiae of male notched before apex.
- 2(5) Body entirely fulvous, including antennae and legs. Pubescence of abdomen simple.
- 3(4) Epipleurae with short and sparse pubescence. Male: tarsi elongate, segment 1 of fore-tarsi about 2.5 times as long as wide; hind tibiae almost straight, narrowly notched just before apex, with triangular plate on extreme apex (Fig. 24). Aedeagus (Fig. 7) not bifurcate at apex. Length 6–8 mm.

Southern India. ..... A. andrewesi (Jacoby, 1895)

4(3) Epipleurae with long and dense pubescence. Male: tarsi feebly elongate, segment 1 of fore-tarsi a little longer than wide; hind tibiae angulate in apical third, distinctly notched before apex. Aedeagus (Fig.9) bifurcate at apex. Length 7.2 mm.

Eastern India. ..... A. orientalis sp.nov.

- 6(1) Elytra bicolorous.
- 7(8) Underside, legs, antennae apart from basal segments black, sometimes partly fulvous. Pubescence of underside dark. Upperside fulvous with rear half of elytra black, or with black postmedian band, sometimes broadly interrupted on suture. Male: segment 1 of mid-tarsi strongly widened (Fig. 26), segment 1 of fore-tarsi about 3 times as long as wide. Hind tibiae straight, widened to apex, without preapical notch. Aedeagus (Fig. 10) with rounded apex, underside with deep and broad preapical groove (broader than lateral margins). Length 6.3–9 mm.

Southern India. [syn. A. externesignata Pic, 1943] ... A. lateralis Lacordaire, 1848

- 8(7) Underside, legs and antennae fulvous or partly black. Pubescence of underside light. Male: segment 1 of mid-tarsi slightly widened or not at all, apex of aedeagus not rounded.
- 9(12) Elytra with broad black band beyond centre, sometimes widened to the rear, but apex of elytra fulvous. Hind tibiae of male feebly notched before apex. Antennae, legs and underside fulvous. Abdomen of male without hairy brushes.

- 12(9) Apical half of elytra entirely black. Antennae, legs and underside fulvous or partly black.
- 13(16) Hind tibiae of male with very deep triangular incisure on underside before apex; base of abdomen with very dense and long pubescence. (Species from southern India, Sri Lanka.)
- 15(14) Epipleurae with short, somewhat indistinct pubescence. Male: abdominal sternites 1–3 with a pair of hairy brushes directed obliquely backwards. Segment 1 of fore-tarsi about 1.2–1.3 times as long as wide, hind tibiae strongly curved and widened to apex. Aedeagus (Fig. 14) with truncate apex and triangular apical tip, its latero-apical angles distinct, more or less produced, underside with narrow median groove before apex. Length of male 4.3–5.7 mm; of female 5–7 mm.

Southern India, Sri Lanka. [*A. incisipes* L. Medvedev, 1984 **syn.nov.**] ...... *A. flavobasalis* (Jacoby, 1908) **comb.nov.** 

16(13) Male: hind tibiae not emarginate, but with transverse ridge on underside before apex, moderately widened apically and slightly curved. Anterior tarsi elongate, segment 1 about 2–2.2 times as long as wide, 2nd broadly triangular, as long as wide, 3rd broad, with basal part as long as apical lobes. Abdomen with simple pubescence. Aedeagus with elongate triangular apex and narrow median groove on underside (Fig. 15). Length of male 6.3–7 mm; of female 7.3–8.2 mm.

North India, Nepal, Assam. ..... A. dimidiatipennis Baly, 1878

#### Aetheodactyla orientalis sp.nov.

Material examined. Holotype (male): East India, Ramanborog (LM).

**Description.** Male: Anterior tarsus not elongate, 1st segment widened, 1.1 times as long as wide, 2nd triangular, as long as wide, 3rd 1.1 times as long as wide, with its basal part about 4 times shorter than apical lobes. Mid-tarsi practically the same. Hind-tibia curved, angulate in apical third, with shallow emargination on underside before apex. Epipleurae with long and dense pubescence. Aedeagus (Fig. 9) with unusual bidentate apex, with feeble and narrow median groove on underside, deeper before apex; lamellae of orifice triangularly widened laterally; viewed laterally, practically in line with the underside.

Length: 7.2 mm.

**Distribution.** East India.

**Differential diagnosis.** Very near to *A. andrewesi* Jacoby, 1895 morphologically and fully identical in coloration, differs only in secondary sexual characters of male.

### Aetheodactyla minor (Duvivier, 1891) stat.nov.

Pseudoclytra plagiata var. minor Duvivier, 1891

**Remarks.** Types of both taxa were studied. *A. minor* differs clearly from *A. plagiata* in the quite different form of the aedeagus and is therefore a distinct species.

#### Aetheodactyla flavobasalis (Jacoby, 1908) comb.nov.

Gynandrophthalma flavobasalis Jacoby, 1908 Smaragdina flavobasalis: TAKIZAWA (1990) Aetheomorpha incisipes L. Medvedev, 1984 syn.nov.

**Remarks.** I have studied a male specimen of *A. flavobasalis* determined and illustrated by TAKIZAWA (1990) and found it identical with *A. incisipes* I described from Sri Lanka. Moreover, the species has to be removed from *Smaragdina* Chevrolat, 1837 to *Aetheomorpha*.

#### Aetheodactyla regalini sp.nov.

**Material examined.** Holotype (male): southern India, Nilgiri Hills, Moyar Camp, 900 m, V. 1954, leg. Nathan (LM). Paratypes: same locality, 3 males, 4 females (LM); Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 1 male (NHMB).

**Description.** Red fulvous, rear half of elytra black. Sometimes antennae, legs and abdomen more or less darkened. Epipleurae with long, dense, light yellow pubescence.

Male: anterior abdominal sternites with longer and denser pubescence than on other parts of underside but not arranged in two brushes as in *A. flavobasalis*, and all hairs directed backwards, not occurring obliquely. Fore-tarsi not elongate, 1st segment 1.5 times, 2nd as long as wide, 3rd with basal part about 4 times shorter than apical lobes.

Mid-tarsi with 1st segment feebly enlarged. Hind tibiae feebly curved, moderately widened apically; preapical incisure of the same form and as deep as in *A. incisipes*. Aedeagus (Fig. 13) with rounded triangular apex, ending in a short tip, without distinct lateroapical angles; underside with deep rounded ovate fovea before apex; lamellae of orifice longer, almost straight, forming longitudinal split between them (in *A. incisipes* these are distinctly arcuate and forming a more or less round hole between them).

Length of male: 5.2–6.7 mm. Female more widened towards the rear, length 6.8–7.6 mm.

Distribution. Southern India.

**Differential diagnosis.** Very near to *A. flavobasalis*, differs mostly in secondary sexual characters.

**Derivatio nominis**. I dedicate this species to Mr. Renato Regalin, a specialist in the Clytrinae.

### Aspidolopha dembickyi sp.nov.

**Material examined.** Holotype (male): Tamil Nadu, Nilgiri Hills, 15 km SE Kotagiri, Kunchappanai, 17–22.V.1997, leg. Dembický & Pacholátko (NHMB). Paratypes: Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko, 10 females (NHMB, 2 exx. – LM).

**Description.** Metallic blue, including abdominal tergites, labrum, basal segments of antennae, elytra and basal half of all tibiae fulvous; elytra often with humeral spot, sometimes also with median spot and traces of preapical spot dark blue (Figs 20–22).

Head glossy, densely punctate. Antennae distinctly serrate from the 5th segment, 4th segment triangular, but much smaller than 5th. Prothorax narrowed towards the front, strongly punctate throughout, but more sparsely in the middle, glossy. Elytra 1.4 times as long as wide, narrowed towards the rear, strongly punctate; punctures in posterior half arranged in irregular rows. 1st abdominal sternite with impunctate glabrous spot on each side. Male pygidium without longitudinal ridge, with apex narrowly margined. Aedeagus with elongate apex (Fig. 16). Female pygidium with longitudinal ridge and triangularly emarginate apex (Fig. 27).

Length: male 5 mm, female 6–6.3 mm.

Distribution. Southern India.

**Differential diagnosis.** The new species differs clearly from all known species in the form of the aedeagus, but in any case it is near *A. decora* Fabricius, 1801, from which it differs in fulvous labrum and basal parts of tibiae, females also in ridged pygidium. JACOBY (1908) indicated that the pygidium of *A. decora* is "carinate in the middle", but this is definitely a mistake, because the typical form of *A. decora* (with fulvous scutellum) has a pygidium that is not carinate. Specimens of this new species with spotted elytra are very similar to *A. distincta* Duvivier, 1891, which however has bicolourous prothorax and fulvous abdominal tergites.

Derivatio nominis. The species is dedicated to its collector, Mr. M. Dembický.



Figs 19–28. 19–23, elytron pattern: 19, *Epimela minuta* sp.nov.; 20–22, *Aspidolopha dembickyi* sp.nov.; 23, *Aetheomorpha sculpturata* sp.nov. 24–25, hind-tibia of male: 24, *Aetheodactyla andrewesi* Jacoby; 25, *A. wallardiensis* Jacoby. 26, *A. lateralis* Lacordaire: mid tarsus of male. 27, *Aspidolopha dembickyi* sp.nov.: pygidium of female. 28, *Smaragdina pacholatkoi* sp.nov.: antenna.

### Aetheomorpha sculpturata sp.nov.

**Material examined.** Holotype (female): Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko (NHMB). Paratypes. Same locality, 2 females (NHMB, LM).

**Description.** Head black with fulvous clypeus, antennae black with fulvous basal segments, prothorax reddish-fulvous with 4 black spots in a transverse row, scutellum black, elytra flavous with black pattern: humeral spot, lateral stripe narrowing towards the rear, ovate spot before centre and near side margin, common sutural spot in anterior third, prolonged towards the rear in sutural stripe black, extreme apex very narrowly margined in black (Fig. 23). Propleurae fulvous with black spot, meso- and metasternum black, abdomen fulvous with more or less black centre, pygidium black with fulvous base or fulvous with black apex, legs fulvous with tarsi slightly darkened.

L. N. MEDVEDEV

Body cylindrical, rather robust. Head without distinct impressions, clypeus bare and almost impunctate, frons and vertex pubescent, densely punctate, frons 2.5 times as wide as eye. Antennae feebly serrate from the 5th segment, segment 3 very small, 4 elongate triangular, next segments approximately as long as wide, subtriangular or subquadrate. Prothorax 1.8 times as wide as long, broadest beyond centre, very densely and coarsely punctate. Scutellum triangular with rounded apex, distinctly punctate and pubescent in basal half. Elytra 1.4 times as long as wide, strongly and densely punctate, more feebly on apical slope, punctures in lateral half form irregular rows. Pygidium pubescent, densely punctate, with triangular emargination on apex, not covered with elytra. Propleurae not pubescent.

Length: 3.5–3.9 mm.

Distribution. Southern India.

**Differential diagnosis.** Propleurae not pubescent. Differs from all species of *Aetheomorpha* in the very strong sculpture of upperside, especially prothorax, and specific pattern of upperside.

## Smaragdina pacholatkoi sp.nov.

**Material examined.** Holotype (male): Tamil Nadu, Nilgiri Hills, 11 km SE Kotagiri, Kunchappanai, 1100 m, 3–15.V.2002, leg. Pacholátko (NHMB). Paratypes. Same locality, 56 exx. of both sexes (NHMB, 4 exx. – LM).

**Description.** Entirely fulvous, only antennal segments 4–11 darkened to black.

Body cylindrical (male) or slightly widened towards the rear (female). Head glossy, impunctate, with 3 grooves on frons, which is 1.6–1.8 times as wide as eye in male, 2.0–2.2 times in female. Antennae serrate from the 4th segment, segment 5–10 feebly transverse, with more or less rounded innerside and apical angle not acute (Fig. 28). Prothorax twice as wide as long, broadest beyond centre, glossy and impunctate. Scutellum triangular with rounded apex, impunctate. Elytra 1.4–1.5 times as long as wide, very distinctly and confusedly punctate except for apical slope which is almost smooth. Segment 1 of fore-tarsi slightly widened in male, subcylindrical. Aedeagus (Fig. 18) with long finger-like apical process.

Length: male 3.7–4.0 mm, female 4.3–4.7 mm.

Distribution. Southern India.

**Differential diagnosis.** *S. divisa* Jacoby, 1889, is very similar, but is distinctly larger, with black tarsi, pale apical part of elytra, antennal segments 5–10 sharply triangular, with straight innerside and acute apical angle; aedeagus is also quite different, without finger-like apical process.

Derivatio nominis. The species is dedicated to its collector, Mr. Petr Pacholátko.

# Acknowledgements

I am grateful to Dr. M. Brancucci for opportunity to study the interesting material in the Naturhistorisches Museum Basel.

## References

BRYANT G. E. (1923): Synonymy in the Clytrinae. Annals and Magazine of Natural History 12(9): 134.
JACOBY M. (1908): Fauna of British India. London, 534 pp.
TAKIZAWA H. (1990): Notes on Chrysomelid beetles (Coleoptera, Chrysomelidae) of India and its neighbouring areas. Part 9. Japan Journal of Entomology 58(4): 746–760.

#### Address of author:

Prof. Lev. N. Medvedev Institute for Problems in Ecology and Evolution Russian Academy of Sciences Leninsky prospect 33 Moscow 119071 RUSSIA