

**Zeitschrift:** Entomologica Basiliensia et Collectionis Frey  
**Herausgeber:** Naturhistorisches Museum Basel, Entomologische Sammlungen  
**Band:** 30 (2008)

**Artikel:** The head and alimentary canal of a larva of *Byturus tomentosus* DeGeer, 1774 (Coleoptera, Byturidae)  
**Autor:** Kolibá, Jií  
**DOI:** <https://doi.org/10.5169/seals-981053>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 20.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## The head and alimentary canal of a larva of *Byturus tomentosus* DeGeer, 1774 (Coleoptera, Byturidae)

by Jiří Kolibáč

**Abstract.** The ventral side of the cranium, the mouth parts (maxilla, mandible, labium, labrum) and alimentary canal of a larva of *Byturus tomentosus* DeGeer, 1774 were studied. Ink drawings and photographs of the body parts are provided. The relations of the observed structures to those in Cleroidea larvae are briefly discussed.

**Keywords.** Coleoptera – *Byturus tomentosus* – larva – morphology

### Introduction

The family Byturidae used to be classified within the traditional superfamily Cucujoidea and considered a relative of Biphyllidae, often placed near Erotylidae (CROWSON 1955; LAWRENCE & NEWTON 1982, 1995). In a recent phylogenetic study based on molecular data (HUNT *et al.* 2007), the authors classify Byturidae and Biphyllidae as basal taxa of Cleroidea, together with Phloiophilidae. This means that a morphology of Byturidae is important to the future study of relations among families of Cleroidea and Cucujoidea (see also BEUTEL & POLLOCK 2000, BEUTEL & SLIPINSKI 2001, LESCHEN *et al.* 2005).

The larva of the raspberry beetle *Byturus tomentosus* DeGeer, 1774 is well known and has been studied several times (e.g., BÖVING & CRAIGHEAD 1931, KLAUSNITZER 2001, LAWRENCE 1991, VAN EMDEN 1942). In this communication, some body parts are examined in detail.

### Larva of *Byturus tomentosus* DeGeer, 1774

(Figs 1–9)

**Description.** Head prognathous and protracted, distinctly pigmented, transverse, sides rounded, with about 20 long setae on the dorsal surface (Fig. 6) and about 12 shorter setae on the ventral surface (Fig. 7).

Cranium (Figs 6, 7, 8): frontal arms curved (lyriform), contiguous at base (Fig. 6); endocarina and epicranial stem absent; five ocelli in pattern 3–2; frontoclypeal suture absent, clypeus protracted (Fig. 6); gular sutures distinct; gular area with two apodemes; hypostomal rods as long as half of cranium; paragular sclerites absent; hypostomal bracon absent (Fig. 7); socket and short, stout seta (Fig. 8) situated in anterior corner of hypostomal margin near ventral mandibular articulation (term after LAWRENCE 1991).<sup>1)</sup>

<sup>1)</sup>The “ventral mandibular articulation” is the site in the hypostomal area of the ventral surface of the cranium where the mandible is joined to the cranium.

Antenna (Figs 4, 8): first antennal segment transverse, second and third elongate; sensory appendix nearly as long as third segment.

Maxilla (Figs 2, 9): cardo transverse, triangular, much smaller than stipes; stipes elongate, coalescent with mala; maxillary palpi 3-segmented, first and second segments weakly transverse, third elongate; apex of lacinia with about 9 stout, pale setae and 3 blunt, dark-pigmented thorns (Fig. 2: several pale setae covering the thorns removed); pedunculate seta absent.

Labium (Fig. 2): ligula membranous, semitransparent in upper portion and pigmented at base, with fine setae along margin; prementum well-developed; mentum fused with prementum or reduced in size; setose, pigmented and sclerotized band situated along each side margin of anterior part of labium (marked as “band” in Fig. 2); two long setae occur in membrane between postmentum and prementum; postmentum elongate, weakly sclerotized, with four long setae.

Labrum (Fig. 1): strongly transverse, anterior margin rounded; setae variable in length, irregularly arranged; basal 2/3 of labrum weakly pigmented. Torma composed of single heavily sclerotized complex formation; torma is figured together with labrum in Fig. 1 but is attached between mandibles and hypopharynx, not to epipharynx (hypopharyngeal sclerome?).

Mandible (Fig. 5): symmetrical; bidentate, ventral apical tooth situated next to the dorsal tooth but at a slightly lower level; medial tooth absent; mola large, tuberculate, with hyaline setose lobe at base; prostheca absent; ventral condyle situated approximately halfway along mandibular base; basal notch shallow, sinuate.

Alimentary canal (Fig. 3): surface of ileum and ventriculus smooth, without papillae or net-like structure; six Malpighian tubules present, all of same length; pyloric valve not visible, without outer lobes; recurvature of ileum weak; ileum relatively long, approximately as long as ventriculus.

**Remarks.** When compared with known larvae of Cleroidea, the following similarities in selected characters appear (references marked by a superscript number are listed at the end of the section):

- (1) Gula with two anteriorly-situated apodemes. This state also occurs in *Thymalus limbatus* Fabricius, *Peltis ferruginea* Linnaeus, *Phloiophilus edwardsi* Stephens, *Protopeltis viridescens* Broun, *Larinotus umblicatus* Carter et Zeck (all Phloiophilidae, Trogossitidae<sup>1</sup>). This feature is considered a plesiomorphy. It also occurs, for example, in *Elateroides dermestoides* Linnaeus and *Lymexylon navale* Linnaeus (Lymexylidae<sup>4</sup>; referred to as “tentorium” by KOLIBÁČ 2002). It has never been found in families of Cleroidea other than Trogossitidae.
- (2) Hypostomal bracon or hypostomal margin (in anterior part of cranium) absent, cranium ventrally “open”. The same state can be found in the Melyrid branch of Cleroidea, especially in Malachiidae<sup>7</sup>.
- (3) Clypeus protracted. This state is unknown in Cleroidea, in which the clypeus is absent and the anterior dorsal (epistomal) margin of the cranium is mostly straight or concave (emarginate)<sup>2,6</sup>.

- (4) Socket and short, stout seta situated in protracted anterior corner of hypostomal margin near ventral mandibular articulation. This character is unknown in Cleroidea, in which the ventral condyle of the mandibles is jointed in the ventral mandibular articulation in the cranium<sup>2,6</sup>. In *Byturus tomentosus*, the ventral condyle is shifted towards the middle of the mandibular base and touches the seta in the socket when the mandible moves (observed in an undissected specimen).
- (5) Ventral condyle situated near the middle of mandibular base: a state unknown in Cleroidea, in which a differently-shaped condyle is always situated at the outer edge of the mandibular base and is joined to the cranium by the ventral mandibular articulation<sup>2,3,6</sup>.
- (6) Setose lobe beneath mandibular mola. This character is unknown in mandibles of Cleroidea, whether the taxa are with or without mola<sup>2,3,5,6</sup>.
- (7) The formation shown together with the labrum (Fig. 1) is very similar to that found in *Idgia* sp. (Prionoceridae)<sup>3</sup>; however, the sclerite in *Byturus tomentosus* is attached between the mandibles and hypopharynx and it is probably homologous with a “hypopharyngeal sclerome”<sup>8</sup>. A torma or tormal sclerite with various apodemes should be attached to the epipharynx<sup>2,3,6</sup>. This state can be found in all known cleroid larvae; it should, however, be checked.
- (8) Maxillary mala with two large thorns and one small, pigmented thorn. Such thorns are unknown in Cleroidea. However, the bidentate projection of the mala (*Thymalus limbatus* Fabricius, *Phlophilus edwardsi* Stephens, *Protopeltis viridescens* Broun) may be derived from these thorns<sup>1,6</sup>.
- (9) Labium without mentum, only “prementum” and postmentum developed. This state is common in all Trogossitidae<sup>6</sup> and some Cleridae<sup>2</sup>. Separation of prementum, mentum and postmentum is developed in Thanerocleridae<sup>5</sup>, Metaxinidae<sup>5</sup> and some Cleridae<sup>2</sup>.
- (10) Sensory appendix large, nearly as long as the third antennal segment. This state is known in *Thymalus limbatus* Fabricius (Trogossitidae)<sup>6</sup> and *Phlophilus edwardsi* Stephens (Phlophilidae)<sup>6</sup>; the sensory appendix is also large in *Melyris scutellaris* Mulsant (Melyridae)<sup>7</sup> and *Lophocateres pusillus* Klug (Trogossitidae)<sup>6</sup>.

**Literature.** CROWSON 1964<sup>1</sup>; KOLIBÁČ 1997<sup>2</sup>, 1999<sup>3</sup>, 2002<sup>4</sup>, 2004<sup>5</sup>, 2006<sup>6</sup>; KOLIBÁČ *et al.* 2005<sup>7</sup>; LAWRENCE 1991<sup>8</sup>.

### Acknowledgement


The Ministry of Culture of the Czech Republic supported this project with grant ref. MK00009486201.

## References


- BEUTEL R. G. & POLLOCK D. A. (2000): *Larval head morphology of Phycosecis litoralis (Pascoe) (Coleoptera: Phycosecidae) with phylogenetic implications*. Invertebrate Taxonomy **14**: 825–835.
- BEUTEL R. G. & SLIPINSKI S. A. (2001): *Comparative study of head structures of larvae of Sphindidae and Protocucujidae (Coleoptera: Cucujoidea)*. European Journal of Entomology **98**: 219–232.
- BÖVING A. G. & CRAIGHEAD F. C. (1931): *An illustrated synopsis of the principal larval forms of the order Coleoptera*. Entomologica Americana, 1–351.
- CROWSON R. A. (1955): *The natural classification of the families of Coleoptera*. Nathaniel Lloyd & Co., Ltd., London, 187 pp.
- CROWSON R. A. (1964a): *A review of the classification of Cleroidea (Coleoptera), with descriptions of two genera of Peltidae and of several new larval types*. Transactions of the Royal entomological Society of London **116**: 275–327.
- EMDEN F. I., van (1942): *Larvae of British beetles. III. Keys to the families*. Entomological monthly Magazine **78**: 206–226, 253–272.
- HUNT T., BERGSTEN J., LEVKANICOVA Z., PAPADOPOLOU A., ST.JOHN O., WILD R., HAMMOND P. M., AHRENS D., BALKE M., CATERINO M. S., GOMEZ-ZURITA J., RIBERA I., BARRACLOUGH T. G., BOCAKOVA M., BOCAK L. & VOGLER A. P. (2007): *A Comprehensive Phylogeny of Beetles Reveals the Evolutionary Origins of a Superradiation*. Science **318(5858)**: 1913–1916.
- KLAUSNITZER B. (2001): *83. Familie: Byturidae*. Pp. 205–208. In: KLAUSNITZER B. (ed.): *Die Larven der Käfer Mitteleuropas. Band 6*. Spektrum Akademischer Verlag, Heidelberg – Berlin, 310 pp.
- KOLIBÁČ J. (1997): *Classification of the subfamilies of Cleridae (Coleoptera: Cleroidea)*. Acta Musei Moraviae, Scientiae naturales **81**: 307–361.
- KOLIBÁČ J. (1999): *Comparative morphology of mandible, epipharynx and alimentary canal in larval and adult Cleroidea (Coleoptera)*. Acta Musei Moraviae, Scientiae biologicae **84**: 11–69.
- KOLIBÁČ J. (2002): *Description of larvae of Lymexylon navale (Linnaeus, 1758) and Elateroides dermestoides (Linnaeus, 1761) (Coleoptera, Lymexylidae)*. Acta Musei Moraviae, Scientiae biologicae **87**: 47–54.
- KOLIBÁČ J. (2004): *Metaxinidae fam.nov., a new family of Cleroidea (Coleoptera)*. Entomologica Basiliensis **6**: 239–268.
- KOLIBÁČ J. (2006): *A review of the Trogositidae. Part 2: Larval morphology, phylogeny and taxonomy (Coleoptera, Cleroidea)*. Entomologica Basiliensis et Collectionis Frey **28**: 105–153.
- KOLIBÁČ J., MAJER K. & ŠVIHLA V. (2005): *Cleroidea. Brouci nadčeledi Cleroidea Česka, Slovenska a sousedních oblastí. Beetles of the superfamily Cleroidea in the Czech and Slovak republics and neighbouring areas*. Clarion, Praha, 186 pp.
- LAWRENCE J. F. (1991): *Order Coleoptera*. Pp. 144–237. In: STEHR F. W. (ed.): *Immature Insects. Vol. 2*. Kendall/Hunt Publishing Company, Dubuque, Iowa, 658 pp.
- LAWRENCE J. F. & NEWTON A. F., Jr. (1982): *Evolution and classification of beetles*. Annual Revue of Ecology and Systematics **13**: 261–290.
- LAWRENCE J. F. & NEWTON A. F., Jr. (1995): *Families and subfamilies of Coleoptera (with selected genera, notes, references and data on family-group names)*. Pp. 779–1006. In: PAKALUK J. & SLIPINSKI S. A. (eds): *Biology, Phylogeny, and Classification of Coleoptera. Papers Celebrating the 80th Birthday of Roy A. Crowson*. Muzeum i Instytut Zoologii PAN, Warszawa, 1006 pp.
- LESCHEN R. A. B., LAWRENCE J. F. & SLIPINSKI S. A. (2005): *Classification of basal Cucujoidea (Coleoptera: Polyphaga): cladistic analysis, keys and review of new families*. Invertebrate Systematics **19**: 17–73.

### **Author's address:**



Dr. Jiří Kolibáč  
 Moravian Museum  
 Department of Entomology  
 Hvězdoslavova 29a  
 627 00 Brno  
 CZECH REPUBLIC  
 E-mails: ento.kol@volny.cz, jkolibac@mzm.cz




**Figs 1–2.** 1 – labrum and hypopharyngeal sclerite or torma, 2 – maxillolabial complex (left maxilla removed, setae covering pigmented thorns in mala removed).




Figs 3–5. 3 – alimentary canal, 4 – antenna, 5 – right mandible dorsally.



**Figs 6–7.** 6 – Cranium dorsally (labrum, mandibles, maxillolabial complex removed); 7 – Cranium ventrally (labrum, mandibles, maxillolabial complex removed). Photo by I. Malenovský and J. Kolibáč (Olympus BX41 + C 5060, Helicon Focus 4.21©).



**Fig. 8.** Cranium and antenna, ventral view. A detail of the ventral mandibular articulation area. Photo by I. Malenovský and J. Kolibáč (Olympus BX41 + C 5060, Helicon Focus 4.21©).



9

**Fig. 9.** Right maxilla (including all setae in mala). Photo by I. Malenovský and J. Kolibáč (Olympus BX41 + C 5060).

