

**Zeitschrift:** Entomologica Basiliensia et Collectionis Frey  
**Herausgeber:** Naturhistorisches Museum Basel, Entomologische Sammlungen  
**Band:** 31 (2009)  
  
**Artikel:** A revision of the fulvous species of the genus *Mimastra* Baly, 1865 from Vietnam (Chrysomelidae, Galerucinae)  
**Autor:** Medvedev, Lev N.  
**DOI:** <https://doi.org/10.5169/seals-981043>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 31.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## A revision of the fulvous species of the genus *Mimastra* Baly, 1865 from Vietnam (Chrysomelidae, Galerucinae)

by Lev Medvedev

**Abstract.** A key to fulvous species of the genus *Mimastra* Baly, 1865 from Vietnam is proposed. Six new species are described: *Mimastra fulvipes*, *M. fulviventris*, *M. korotyaevi*, *M. tamdaoana*, *M. similis*, *M. tarsalis* spp.nov. (all from Vietnam). The following new synonyms are proposed: *M. uncitarsis* Laboissière, 1940 (= *M. chennelli* Baly, 1879); *M. soreli* Baly, 1878, *M. limbata* Baly, 1879, *M. guerrei* Laboissière, 1929 (= *M. cyanura* (Hope, 1831)).

**Key words.** *Galerucinae* – *Mimastra* – Vietnam – key – new species – new synonyms

### Introduction

The many Oriental genera of *Galerucinae* include a large set of “fulvous” species (at least on the upperside of the body), which often appear homogeneous and are usually poorly studied in comparison with the “spotted” or “banded” species. Such a situation once held in such genera as *Aulacophora* Chevrolat, 1836, *Hoplasoma* Jacoby, 1884, *Haplosomoides* Duvivier, 1890, *Paridea* Baly, 1886 and still exists in *Mimastra* Baly, 1865, *Monolepta* Chevrolat, 1836, *Dercetina* Gressitt et Kimoto, 1963, *Arthrotus* Motschulsky, 1857 and others. The genus *Mimastra* is keyed for India (MAULIK 1936), China (OGLOBLIN 1936, GRESSITT & KIMOTO 1963, ZHANG *et al.* 2006) and Indochina (KIMOTO 1989) but identification of fulvous species is difficult in all of them and figures of diagnostic characters (especially aedeagi) are lacking.

In this work I revise the fulvous species of *Mimastra* from only Vietnam, since a very large body of material exists for this region. As a result of this, a key for this group is given, 6 species are described as new for science and 4 new synonyms are stated. Males differ mainly in the form of the aedeagus, females in the proportions of the basal antennal segments and the colour of the underside of the body.

### Material

The following abbreviations are used for the places in which the new species are deposited:

NHMB ..... Naturhistorisches Museum, Basel, Switzerland  
LM ..... The Lev Medvedev collection, Moscow, Russia

### Taxonomy

#### A key to the *Mimastra* of Vietnam with entirely fulvous elytra

- 1(10) Underside entirely fulvous. Antennae longer than body, at least apical segments darkened.
- 2(9) Proportions of segments 2–4 approx. 1–2–3, segment 3 not more than twice as long as 2.
- 3(4) Legs entirely fulvous. Apical antennal segments slightly darkened, segments 2–4 proportions 7–16–22. Elytra lacking ridges. Aedeagus (Fig. 1) feebly narrowing to apex, longitudinally concave in apical third of underside. Body length 6.9–7.7 mm. South Vietnam. .... *M. fulvipes* sp.nov.
- 4(3) Tibiae and tarsi black or distinctly darker than femora.
- 5(6) Elytra with sharp ridge along lateral margin beyond centre. Apical antennal segments slightly darkened, segments 2–4 proportions 7–16–22. Aedeagus almost parallel-sided, with deep ovate impression on apical fifth of underside (Fig. 2). Body length 6.5–7.6 mm. South Vietnam, Laos. .... *M. apicalis* Kimoto, 1989
- 6(5) Elytra without sharp ridge along lateral margin beyond centre.
- 7(8) Antennae piceous with 3–4 basal segments fulvous, segment 4 about 1.5 times as long as 3 (segments 2–4 proportions 7–15–21). Upperside light fulvous. Tibiae and tarsi piceous, but not black. Transverse impression of prothorax usually interrupted halfway. Elytra lacking more or less distinct longitudinal costae. Aedeagus parallel-sided with ovate transparent impression in apical fifth of underside (Fig. 3). Body length 6.3–7.6 mm. Vietnam, Laos, Thailand, Burma, China. .... *M. gracilicornis* Jacoby, 1889
- 8(7) Antennae black or with 1–2 basal segments dark fulvous, segment 4 about 1.2–1.25 times as long as 3 (segments 2–4 proportions 8–18–22). Upperside red-fulvous, scutellum sometimes darkened. Legs black, but femora often fulvous with black apices. Elytra with 3–4 feeble longitudinal costae. Aedeagus (Fig. 4) parallel-sided, with broad, longitudinal impression on apical third of underside, partly membranaceous. Body length 6.8–8.5 mm. Laos, Thailand, South India, Burma, possible in Vietnam. .... *M. scutellata* Jacoby, 1904
- 9(2) Antennal segment 3 about 2.5–3 times as long as 2. Aedeagus – Fig. 5. Body length 6.7–7.1 mm. North Vietnam. Specimens with completely fulvous underside. See also item 15. .... *M. fulviventris* sp.nov.
- 10(1) Underside not entirely fulvous.
- 11(18) Metasternum and abdomen differ in colour.



- 12(13) Metasternum fulvous, abdomen black (except apex). Femora fulvous with black stripe on upperside, tibiae and tarsi more or less piceous. Antennal segments 2–4 proportions 8–13–23. Aedeagus narrowed before rounded apical part, its underside with longitudinal impression in apical third, widened to apex (Fig. 6). Body length 7.2–7.8 mm. Extreme North Vietnam (Fanxipan), China, North India, Nepal. .... ***M. gracilis* Baly, 1878**
- 13(12) Metasternum black, abdomen fulvous, sometimes darkened but not black. Femora fulvous without black stripe on upperside. Antennae longer than body, brown or piceous with fulvous basal segment.
- 14(17) Antennal segment 3 distinctly longer than 2, segments 2+3 about as long as 4. Legs fulvous, tibiae sometimes slightly darkened on dorsal side.
- 15(16) Antennal segment 3 about 2.5–3 times as long as 2. Aedeagus – Fig. 5. Body length 6.7–7.1 mm. North Vietnam. Specimens with metasternum black. See also item 9. .... ***M. fulviventris* sp.nov.**
- 16(15) Antennal segment 3 about 1.5–1.6 times as long as 2. Abdomen sometimes more or less piceous, but not black. Aedeagus – Fig. 7. Body length 5.3–7.3 mm. North Vietnam. See also item 20. .... ***M. korotyaevi* sp.nov.**
- 17(14) Antennal segments 2 and 3 subequal, segments 2+3 twice as short as segment 4 (segments 2–4 proportions 5–5–19). Femora fulvous, tibiae and tarsi piceous. Aedeagus slightly widened to apical quarter and then narrowed again, underside without distinct impressions, extreme apex curved downward in lateral view (Figs 8, 8a). Body length 6.8–7.5 mm. .... ***M. pectoralis* Kimoto, 1989**
- 18(11) Metasternum and abdomen black or piceous, sometimes with metallic lustre.
- 19(32) Segment 1 of anterior tarsus of male simple, narrow and elongate. Head and prothorax without black spots.
- 20(21) Abdomen more or less piceous, but not black. Antennal segment 3 about 1.5–1.6 times as long as 2. Aedeagus – Fig. 7. Body length 5.3–7.3 mm. North Vietnam. See also item 16. .... ***M. korotyaevi* sp.nov.**
- 21(20) Abdomen black, sometimes with metallic lustre.
- 22(23) Underside with distinct metallic lustre, apical abdominal sternite and pygidium not fulvous. Proportions of antennal segments 2–4 approx. 1–2–4. Antennae longer than body. Aedeagus narrowed to apex, extreme apex triangular, but curved upward, lending it a narrowly truncate appearance in ventral view (Fig. 9). Body length 7.5–8.2 mm. North Vietnam. .... ***M. tamdaoana* sp.nov.**

- 23(22) Underside black with apical abdominal sternite and pygidium fulvous.
- 24(25) Antennae shorter than body, about  $\frac{3}{4}$  or  $\frac{5}{6}$  of body length, segment 3 about 1.3–1.65 times as long as 2, segment 4 more than twice as long as 3. Antennae, tibiae and tarsi infusate. Aedeagus cuneiform with narrowly rounded apex (Fig. 10). Body length 5.3–7.7 mm. South Vietnam, Laos, Thailand, Burma (*M. badia* Kimoto, 1989). ..... ***M. polita* Jacoby, 1989**
- 25(24) Antennae longer than, or as long as, body. Aedeagus distinctive or, if cuneiform, then not with acute apex.
- 26(29) Antennal segment 3 not more than twice as long as segment 2. Proportions of antennal segments 2–4 approx. 1–2–3.
- 27(28) Aedeagus thin and very long, about 7–9 times as long as width before base, parallel-sided, longitudinally impressed on underside (Fig. 11). Body length 6.7–7.1 mm. Vietnam. .... ***M. kabakovi* sp.nov.**
- 28(27) Aedeagus cuneiform with acute apex, about 5 times as long as width before base, longitudinally concave on underside (Fig. 12). Body length 5.7–7.4 mm. North Vietnam. .... ***M. similis* sp.nov.**
- 29(26) Antennal segment 3 about 2.5 times as long as segment 2. Segments 2–4 proportions 1–2.5–3.7.
- 30(31) Aedeagus unknown. Body length 6.0–7.5 mm (according to original description). South Vietnam (Dalat), Laos (Houei Kong). .... ***M. persimilis* Kimoto, 1989**
- 31(30) Aedeagus lanceolate, longitudinally concave on underside (Fig. 13). Elytra sometimes with poorly delimited darkened area. Body length 6.6–7.4 mm. North Vietnam. .... ***M. persimilis* Kimoto, 1989**
- 32(19) Segment 1 of anterior tarsus strongly widened in male. Underside with feeble metallic lustre. Antennae a little shorter than body length, piceous or brown-piceous with basal segments fulvous. Legs fulvous with most of tibiae and tarsi blackish.
- 33(36) In male segment 1 of anterior tarsus very broad, not curved laterally; its upperside lustrous, concave and impunctate.
- 34(35) Segment 1 of anterior tarsus of male irregularly round or feebly elongate ovate, not more than 1.2 times as long as wide, its upperside feebly concave and not ridged at the sides (Fig. 18). Sculpture of underside asymmetrical, with deep oblique impression in basilateral area (Fig. 17). Antennal segments 2–4 proportions 6–9–19. Vertex with black spots: two on hind margin (mostly covered by prothorax) and one elongate in middle; prothorax with a few dark spots. Apex of elytra often bordered with dark metallic stripe. Aedeagus (Fig. 14) narrowed to apex,



underside with deep longitudinal impression, sharply delimited at sides. Body length 7.8–8.8 mm. Vietnam, Laos, Thailand, South China. (*Mimastra lunata* (Kollar et Redtenbacher, 1848); *M. davidis* (Fairmaire, 1878); *M. limbata* Baly, 1879; *M. apicalis* Baly, 1886; *M. soreli* Baly, 1878; *M. latimanus* Allard, 1889; *M. guerrei* Laboissière, 1929). .....

..... ***M. cyanura* (Hope) (fulvous form)**

- 35(34) Segment 1 of anterior tarsus of male transversely ovate, 1.4 times as wide as long, its upperside deeply concave and ridged along base and at the sides (Fig. 22 ); sculpture of underside symmetrical, with deep basal impression and apical area divided into three parts (Fig. 21). Antennal segments 2–4 proportions 6–12–24. Two spots and longitudinal stripe on vertex and 4 poorly delimited spots on prothorax piceous. Aedeagus (Fig. 15) very much as in preceding species. Body length 9.9 mm. ....

..... ***M. tarsalis* sp.nov.**

- 36(27) Male: segment 1 of fore-tarsus strongly curved, its upperside lustrous and convex (Fig. 20), underside with longitudinal ridge that divides ventral surface into three concave areas; one of these is round or ovate, lustrous and impunctate (Fig. 19). Vertex and prothorax with a few dark spots, as in preceding species. Apex of elytra not bordered with dark stripe. Aedeagus (Fig. 16) almost parallel-sided with deep longitudinal impression in apical two-fifths, sharply bordered at the sides. Body length 7.4–9.0 mm. Vietnam, China, Assam. (*Mimastra uncitarsis* Laboissière, 1940). ..... ***M. chennelli* Baly, 1879 (fulvous form)**

### ***Mimastra fulvipes* sp.nov.**


**Material examined.** Holotype (male): South Vietnam, Gialai-Contum prov., 40 km N Ankhe, Buon Loi, 10.VI.1980, leg. L. Medvedev (LM). Paratypes: same locality, VI–VII.1983, 9 ex. (LM, 3 ex. NHMB); – South Vietnam, Dongnai Prov, Mada, 24.VI.1991, 1 male (LM).

**Description.** Body entirely fulvous, only apical antennal segments slightly darkened.


Body elongate, slightly widening towards the rear. Head impunctate, frontal tubercles triangular, convex, sharply delimited posteriorly. Antennae longer than body, proportions of segments: 18–7–16–22–22–22–23–23–23–21–22, preapical segments approx. 10 times as long as wide. Prothorax 1.35 times as wide as long, subrectangular, broadest at anterior margin and narrowed to base, with sides almost straight, surface lustrous and impunctate. Scutellum triangular. Elytra 1.8 times as long as wide, surface with dense, feeble punctures and fine microsculpture. Segment 1 of fore-tarsi simple in both sexes. Aedeagus feebly narrowed to apex, longitudinally concave in apical third of underside (Fig. 1).

Body length: 6.9–7.7 mm.

**Differential diagnosis.** Near *M. gracilicornis* Jacoby, 1899, differs in colour of legs, broader body and form of aedeagus.



**Figs 1–10.** Aedeagi, ventral view: 1 – *Mimastrea fulvipes* sp.nov., 2 – *M. apicalis* Kimoto, 3 – *M. gracilicornis* Jacoby, 4 – *M. scutellata* Jacoby, 5 – *M. fulviventris* sp.nov., 6 – *M. gracilis* Baly, 7 – *M. korotyaevi* sp.nov., 8 – *M. pectoralis* Kimoto (8a – extreme apex in lateral view), 9 – *M. tamdaoana* sp.nov., 10 – *M. polita* Jacoby.



**Figs 11–16.** Aedeagi, ventral view: 11 – *M. kabakovi* sp.nov., 12 – *M. similis* sp.nov., 13 – *M. persimilis* Kimoto, 14 – *M. cyanura* (Hope), 15 – *M. tarsalis* sp.nov., 16 – *M. chennelli* Baly.

***Mimastra fulviventris* sp.nov.**


**Material examined.** Holotype (male): Vietnam, Vinh Phu prov., Tam Dao, 24–31.V.1985, leg. L. Medvedev (LM). Paratypes: same locality and date, 7 ex. (LM, 2 ex. NHMB); – same locality, 11–13.V.1975, 6 ex. (LM, 1 ex. NHMB); 12–22.IV.1986, 1 ex. (LM).

**Description.** Fulvous, 3 or 4 apical antennal segments slightly darkened, underside with metasternum black or completely fulvous.

Body elongate, widened towards the rear. Head impunctate, frontal tubercles convex, short triangular, sharply delimited posteriorly. Antennae longer than body, proportions of segments: 14–7–20–20–20–20–20–20–20–18, preapical segments about 15 times as long as wide. Prothorax 1.7 times as wide as long, rectangular, feebly narrowed to base, with almost straight hind margins, surface lustrous and impunctate. Scutellum triangular with rounded apex. Elytra 1.6 times as long as wide, surface lustrous, densely and strongly punctate. Segment 1 of fore-tarsi simple in both sexes. Aedeagus long and thin, almost parallel-sided with finger-like apical process, underside longitudinally concave (Fig. 5).

Body length: 6.7–7.1 mm.





**Figs 17–22.** Anterior male tarsi from below and above (Figs 17–20 after LABOISSIÈRE 1940): 17, 18 – *Mimastra cyanura* (Hope); 19, 20 – *M. chennelli* Baly; 21, 22 – Segment 1 of male anterior tarsus of *M. tarsalis* sp.nov. from below and above.

**Differential diagnosis.** Near *M. fulvipes* sp.nov., differs clearly in form of aedeagus and proportions of basal antennal segments, the main character for assessing females, especially specimens with entirely fulvous underside.

***Mimastra korotyaevi* sp.nov.**

**Material examined.** Holotype (male): Vietnam, 100 km W Thanhhoa, Langthanh, 400 m, 23.I.1989, leg. B. Korotyaev (LM). Paratypes: same locality, 1 female (NHMB); – North Vietnam, Kim Shon, 7–8.III.1962, leg. O. Kabakov, 1 female (LM); North Vietnam, mountains NW Qui Chau, 200–400 m, 7.III.1962, leg. O. Kabakov, 1 female (LM).

**Description.** Fulvous, antennae with darkened apical segments, metasternum black, abdomen mostly piceous with fulvous apex, sometimes almost fulvous.

Body moderately elongate, widened towards the rear. Frontal tubercles triangular, convex, sharply delimited posteriorly. Antennae longer than body, proportions of segments approx. 15–5–10–19–19–17–20–19–19–17–17, preapical segments about 7–8 times as long as wide. Prothorax 1.5 times as wide as long, very feebly narrowed to base, with side margins straight or feebly concave, surface lustrous and impunctate. Scutellum triangular. Elytra 1.7 times as long as wide, surface finely and densely punctate. Segment 1 of fore-tarsi simple in both sexes. Aedeagus almost parallel-sided, slightly narrowing to triangular apex, underside with shallow longitudinal impression in middle (Fig. 7).

Body length: 5.3–7.3 mm.

**Etymology.** The species is dedicated to its collector, Dr. B. Korotyaev.

**Differential diagnosis.** Near *M. flaviventris* sp.nov., differs in proportions of basal antennal segments and form of aedeagus.

***Mimastra tamdaoana* sp.nov.**

**Material examined.** Holotype (male): Vietnam, Prov. Vinh-Phu, Tam Dao, 800–1200 m, forest, 12–22.IV.1986, leg. L. Medvedev et al. (LM). Paratypes: same locality and date, 1 male, 1 female (LM, NHMB); same locality, VI.1983, leg. L. Medvedev, 1 female (LM).

**Description.** Fulvous, head and prothorax lacking dark spots, antennae piceous with fulvous basal segments, elytra with apical margin very narrowly black, metasternum, abdomen and pygidium black with distinct metallic lustre, tibiae and apical margin of femora black.

Body narrow and elongate, almost parallel-sided. Head impunctate, frontal tubercles triangular, convex, sharply delimited posteriorly. Antennae longer than body (male) or as long as body (female); proportions of segments: 15–6–11–20–20–20–19–15–12–19, preapical segments about 6–8 times as long as wide. Prothorax 1.7 times as wide as long, almost rectangular, broadest in anterior third, surface lustrous and impunctate. Scutellum triangular with rounded apex, very finely strigose. Elytra 1.7–1.8 times as long as wide, surface with dense and moderately strong punctures. Segment 1 of fore-tarsi simple in both sexes. Aedeagus long and thin, narrowed to apex, extreme apex triangular, but curved upward, lending it a narrowly truncate appearance in ventral view, underside with unsclerotized central part (Fig. 9).



Body length: 7.5–8.2 mm.

**Differential diagnosis.** Differs from all species in simple anterior tarsi of male and in the metallic lustre of the underside.

***Mimastra kabakovi* sp.nov.**

**Material examined.** Holotype (male): South Vietnam, Prov. Gialai-Contum, 50 km N Ankhe, Hanung, 14.VI.1980, tropical forest, leg. L. Medvedev (LM). Paratypes: same locality, 14–15.VI.1980, 1 male, 1 female (LM); – same locality, VI.1983, 3 males (LM); North Vietnam, mountains NW Dong Hoi, 300–800 m, 20–24.III.1963, leg. O. Kabakov, 4 males (LM, 1 ex. NHMB); – Vietnam, mountains NW Cui Chau, 200–400 m, 15.III.1961, leg. O. Kabakov, 1 male (LM), 1 female (NHMB); – Vietnam, mountains SW Bai Thuong, 300 m, 18–20.III.1961, leg. O. Kabakov, 2 males, 2 females (LM).

**Description.** Fulvous, apical antennal segments darkened, metasternum and abdomen black (apical sternite and pygidium may be fulvous).

Body elongate, slightly widened towards the rear. Head impunctate, frontal tubercles triangular, convex, well delimited posteriorly. Antennae longer than body, proportions of segments: 15–6–12–18–18–17–17–16–16–15–17, preapical segments about 7–8 times as long as wide. Prothorax 1.5 times as wide as long, subrectangular, very feebly narrowed to base, surface lustrous and impunctate. Scutellum triangular. Elytra 1.6–1.7 times as long as wide, surface with deep, dense punctures. Segment 1 of fore-tarsi simple in both sexes. Aedeagus thin and very long, about 7–9 times as long as wide before base, parallel-sided with triangular apex, underside longitudinally grooved along mid-line (Fig. 11).

Body length: 6.7–7.1 mm.

**Etymology.** The species is dedicated to its collector, Mr. O. Kabakov.

**Differential diagnosis.** Very near to *M. persimilis* Kimoto, 1989, differs in proportions of antennal segments and form of aedeagus.

***Mimastra similis* sp.nov.**

**Material examined.** Holotype (male): Vietnam, Prov. Vinh-Phu, Tamdao, 800–1200 m, forest, 12–22.IV.1986, leg. L. Medvedev et al. (LM). Paratypes: same locality and date, 2 males, 1 female (LM, 1 ex. NHMB); – North Vietnam, mountains SW Bai Thuong, 20.III.1964, leg. O. Kabakov, 2 males (LM); – North Vietnam, Kim Shon, 7–8.III.1962, leg. O. Kabakov, 1 male (LM); – North Vietnam, Thanh Hoa, 9.I.1962, leg. O. Kabakov, 1 male (LM).

**Description.** Fulvous, antennae piceous with fulvous basal segments, metasternum and abdomen black, but apical sternite and pygidium fulvous.

Body narrow, elongate, slightly widened towards the rear. Head impunctate, frontal tubercles triangular, convex, well delimited posteriorly. Antennae longer than body, proportions of segments: 13–5–10–16–16–15–15–15–14–13–15, preapical segments about 7–8 times as long as wide. Prothorax 1.5–1.6 times as wide as long, broadest in anterior third and narrowed to base, surface lustrous and impunctate. Scutellum triangular. Elytra 1.7–1.8 times as long as wide, surface with dense, deep punctures. Segment 1 of fore-tarsi simple in both sexes. Aedeagus cuneiform with acute apex, about 5 times as long as wide before base, longitudinally concave on underside (Fig. 12).



Body length: 5.7–7.4 mm.

**Differential diagnosis.** Very near to *M. kabakovi* sp.nov., differs mostly in form of aedeagus.

***Mimastra persimilis* Kimoto, 1989**

**Remark.** I was unable to examine a type of this species described from Dalat, but I have a good series from North Vietnam (Bai Thuong, Kon Kuong, Kim Kuong, River Kon) which are identical with the original description of *M. persimilis*. On the other hand, this species does not appear in my very extensive material from South Vietnam.

***Mimastra cyanura* (Hope, 1831)**

*Mimastra lunata* (Kollar et Redtenbacher, 1848)

*Mimastra davidis* (Fairmaire, 1878)

*Mimastra limbata* Baly, 1879 **syn.nov.**

*Mimastra apicalis* Baly, 1886

*Mimastra soreli* Baly, 1878 **syn.nov.**

*Mimastra latimanus* Allard, 1889

*Mimastra guerrei* Laboissière, 1929 **syn.nov.**

**Remark.** All three new synonyms have a widened segment in the male (Figs 17, 18) and aedeagus (Fig. 14) exactly the same as in *M. cyanura* Hope. They differ only in the pattern on the elytra; *M. soreli* Baly, the most common in Vietnam, has entirely fulvous elytra. In general, the elytral pattern of *M. cyanura* is very variable, but this appears to be merely a colour aberration or apparent only in local populations.

***Mimastra tarsalis* sp.nov.**

**Material examined.** Holotype (male): Vietnam, NW Tam Dao ridge, Shon-Duong, 200 m, 4.IV.1962, leg. O. Kabakov (LM).

**Description.** Fulvous, antennae piceous except 4 basal segments, 2 spots and longitudinal stripe on vertex and 4 poorly delimited spots on prothorax piceous, metasternum and abdomen black with feeble metallic gloss. Colour very possibly variable.

Morphologically identical with *M. cyanura* Hope, 1831, differs mainly in form and sculpture of the first segment of anterior tarsi in male. It is 1.4 times as wide as long, more or less trapeziform, upperside deeply concave, with lateral and basal margins strongly elevated and ridged (Fig. 22); underside in basal half deeply concave, impunctate and lustrous, apical half with round shallow impression on each side, with rough sculpture; these impressions are divided by a flat central part covered with very fine, dense punctures (Fig. 21). Proportion of antennal segments 2–4 proportions 6–12–24. Aedeagus – Fig. 15.

Body length: 9.9 mm.

**Differential diagnosis.** See key.

***Mimastra chennelli* Baly, 1879**

*Mimastra uncitarsis* Laboissière, 1940 (syn.nov.)

*Mimastra unicitarsis*: GRESSITT & KIMOTO (1963) (misspelling); KIMOTO (1989) (misspelling).

**Remark.** Both species are entirely identical in form and sculpture of segment 1 of foretarsi of male (Figs 19, 20) as well as in form of aedeagus (Fig. 16). They differ only in colour of elytra: entirely fulvous in *M. chennelli* and having a metallic blue apical spot or elongate stripe in *M. uncitarsis*. These species have indubitably to be united. In Vietnam specimens with unspotted elytra are dominant; I have only one spotted specimen, collected together with the unspotted form.

**References**

- GRESSITT J. L. & KIMOTO S. (1963): *The Chrysomelidae (Coleopt.) of China and Korea, part 2*. Pacific Insect Monograph **1B**: 301–1026.
- KIMOTO S. (1989): *Chrysomelidae (Coleoptera) of Thailand, Cambodia, Laos and Vietnam. VI. Galerucinae*. Esakia **27**: 1–241.
- LABOISSIÈRE V. (1940): *Observations sur les Galerucinae des collections du Musée Royal d'Histoire Naturelle de Belgique et descriptions de nouveaux genres et espèces*. Bull. Mus. Roy. Hist. Nat. Belgique **16(37)**: 1–41.
- MAULIK S. (1935): *The Fauna of British India including Ceylon and Burma. Coleoptera. Chrysomelidae, Galerucinae*. London, Francis & Taylor, 648 pp.
- MEDVEDEV L. N. (2002): *Jacoby's types of Chrysomelidae (Coleoptera) from Burma in the Museo Civico di Storia Naturale "Giacomo Doria", Genoa. Part 3*. Annali del Museo Civico di Storia Naturale "Giacomo Doria" (Genoa) **94**: 249–264..
- OGLOBLIN D. A. (1936): *Listoedy, Galerucinae*. In: *Fauna SSSR. Nasekomye Zhestkokrylye [Chrysomelidae, Galerucinae. Faune de l'URSS, Insectes Coléoptères]*. n.s. 8, Volume 26, part 1. Moskva-Leningrad, Izdatelstvo Akademii Nauk SSSR, xiv + 455 pp.
- ZHANG LI-JIE, YANG XING-KE, CUI JUN-ZHI & LI WEN-ZHU (2006): *A key to the genus Mimastra Baly (Coleoptera: Chrysomelidae: Galerucinae) from China, with the description of a new species*. Entomological News **117(2)**: 203–210.

**Author's address:**

Prof. Lev N. Medvedev  
 Institute for Problems of Ecology and Evolution  
 Russian Academy of Sciences  
 Leninsky prospect 33  
 Moscow 119071  
 RUSSIA  
 E-mail: lev.n.medvedev@mail.ru