Zeitschrift: Entomologica Basiliensia et Collectionis Frey

Herausgeber: Naturhistorisches Museum Basel, Entomologische Sammlungen

Band: 33 (2011)

Artikel: A contribution to knowledge of Oriental species of Cneorane Baly, 1865

(Chrysomelidae, Galerucinae)

Autor: Medvedev, Lev N.

DOI: https://doi.org/10.5169/seals-980974

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 25.10.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

A contribution to knowledge of Oriental species of *Cneorane* Baly, 1865 (Chrysomelidae, Galerucinae)

by Lev N. Medvedev

Abstract. A preliminary key to Oriental *Cneorane* Baly, 1865 is given. Three species (*C. bhutana* – Bhutan, *C. sprecherae*, *C. costipennis* – both Thailand) are described as new for science. *C. nigripennis* Laboissière, 1922 is the new synonym of *C. cariosipennis* Fairmaire, 1888. Four species (*C. ephippiata* Laboissière, 1930, *C. abdominalis* Jacoby, 1890, *C. semipurpurea* Jacoby, 1884, *C. borneensis* Jacoby, 1894) are excluded from the genus.

Keywords. Galerucinae - Cneorane - Oriental Region - key - new species - new combination

Introduction

The Paleotropical genus *Cneorane* Baly, 1865, largely Oriental, is quite well studied. Most of the Oriental species were described in the 19th century and only eight species have been added in the last 50 years: three from China (Gressitt & Kimoto 1963, Chen 1964, Chen & Jiang 1976), four from Nepal (Chûjô 1966, Kimoto & Takizawa 1972, Medvedev 1992, 2004), and one from Bhutan (Kimoto 1977). Keys exist for almost all regions: "British India" (Maulik 1936), Russia and neighbouring regions (Ogloblin 1936), China (Gressitt & Kimoto 1963), Indochina (Kimoto 1989), and Taiwan (Kimoto & Takizawa 1997), but all these keys are essentially based on the colour of the antennae, legs and underside and partly on the sculpture of the elytra. Many species are very briefly described and only poorly known.

I had the opportunity to study almost all the types described by M. Jacoby and the types of species described in recent years. The main problem was with Fairmaire's species, because the types were not found (GRESSITT & KIMOTO 1963).

In the key that follows I have used some additional characters, including aedeagus and spermatheca. In any event, this is a preliminary contribution, based mainly on substantial material held by the Naturhistorishes Museum in Basel and in the author's collection, including a large series from Vietnam.

Abbreviations for the collections housing the material examined

NHMB	. Naturhistorisches Museum Basel, Switzerland
LM	. Lev N. Medvedev collection, Moscow, Russia
JB	. Jan Bezděk collection, Brno, Czech Republic

Taxonomy

A key to the species

- 2(1) Body not entirely fulvous.
- Upperside, including prothorax, black with distinct metallic lustre. Head, breast and legs black with metallic lustre, abdomen pitchy-brown with 2–4 apical segments fulvous. Prothorax and elytra minutely and quite closely punctured. Length 6.5–8.1 mm. China (Sikang/Sichuan).

 C. sikanga Gressitt et Kimoto, 1963
- 4(3) Prothorax fulvous or red, sometimes darkened to entirely black, but always without metallic lustre.
- Apex of elytra broadly rounded. Anterior coxal cavities open. Anterior tibiae not widened in angulate fashion at apex and most of abdomen without brushes in male. Prothorax red or fulvous, rarely darkened. Aedeagus never strongly constricted before apex.
- 7(10) Elytra with erect hairs.
- 9(8) Fulvous including antennae and epipleurae, elytra bluish-black with two basal thirds of lateral margin, a basal quarter of sutural margin and entire basal margin sometimes reddish-fulvous; metasternum and abdomen black. Elytra with dense, distinct punctures and granulate interspaces. Aedeagus slightly constricted in apical quarter, its apex rounded with a short apical tip (Fig. 4). Length 4.3–7.0 mm. Nepal.

 C. hirsuta Kimoto et Takizawa, 1972
- 10(7) Elytra bare or with a few hairs on side and apical margins.
- 11(12) Elytra bluish-black (without metallic lustre) with lateral, apical and (very narrow) anterior margin fulvous (Fig. 43). Head fulvous with black vertex, remainder of body fulvous, underside sometimes partly

- 12(11) Elytra unicolorous, metallic or black (sometimes epipleurae fulvous).
- 13(16) Antennae fulvous, usually with apex of apical segment darkened and segments 1 and 2 slightly darkened above.

- 16(13) Antennae usually black with a few basal segments, sometimes also with one or two apical segments, red or fulvous; rarely entirely black.
- 17(22) Head fulvous with black or bluish-black vertex.
- 18(21) Species from the Himalayas.

- 22(17) Head entirely fulvous.
- 23(28) Elytra with sharp ridges.
- 25(24) Dorsal part of elytra with ridges. Body much larger.

- 28(23) Elytra without distinct ridges.
- 29(46) Legs entirely fulvous, sometimes tarsi and apices of tibiae more or less darkened.
- 30(35) Antennal segments 8–10 thickened in males.
- 31(32) Thickened antennal segments distinctly elongate (about twice as long as each of segments 8 and 9, about 1.5 times at segment 10 (Fig. 42). Metasternum deep blue, tarsi mostly blackish, apical antennal segments black. Elytra closely punctate. Length 6.0–7.0 mm. Northern India, Nepal. Indication for Burma (MAULIK 1936) needs confirmation.

 C. braeti Duvivier, 1892
- 32(31) Thickened antennal segment only slightly elongate, almost subquadrate (Fig. 37). Two or three apical antennal segments fulvous.

- 35(30) Apical antennal segments of male not thickened. Elytra dull, with microsculptured interspaces of punctures.
- 36(37) Species from Sumatra. Scutellum and elytra metallic green, sides of mesosternum, metasternum and abdomen bluish-green, head and prothorax with green tint. Antennae short, reach anterior third of elytra. Prothorax with dense, weak punctures, elytra densely and very finely

punctate. Length 6.0 mm. Sumatra. Unclear species. 37(36) Continental species. Scutellum fulvous to black. 38(41) Underside and pygidium entirely fulvous. Scutellum fulvous. Elytra very densely and comparatively strongly punctate, interspaces very half-puncture diameter, narrow. about convex, microsculptured. Antennae with 1–2 apical segments reddish. 39(40) Antennae with two apical segments reddish, segment 4 about twice as long as 2. Length 7.6–8.1 mm. SW China (Fukien, Kweichow). 40(39) Antennae with one apical segment reddish, segment 4 about 2.2-2.7 times as long as 2. Length 7.5-9.5 mm. Myanmar, northern India, 41(38) Metasternum and abdomen black or metallic. 42(43) Species from northern India (Manipur) and Nepal. Antennae reach apical area of elytra. Prothorax sparsely punctate, punctures somewhat larger and more crowded on sides and near base. Elytra lustrous metallic blue, very sparsely and finely punctate, tarsi piceous. Length 8.0 mm. Poorly known species. 43(42) Species from China and Korea. 44(45) Aedeagus comparatively thick in lateral view, with short apical triangle (Fig.11). Tarsi sometimes more or less darkened. Spermatheca – Fig. 34. Length 5.5–8.0 mm. South of Russian Far East, eastern Mongolia, all eastern China from Russia to Vietnam, Korea. Indication for Taiwan (Кімото & Такіzawa 1997) is in fact C. cyanipennis Chûjô, item 72. [syn. C. elegans Fairmaire, 1887, C. rufipes Weise, 1889]. Feeding on 45(44) Aedeagus very thin in lateral view, with longer apical triangle (Fig. 12). Tibiae and tarsi from entirely fulvous to entirely black, mostly partly blackish. Spermatheca – Fig. 35. Length 6.5–8.5 mm. Southern China, to the north reaching Sichuan and Shansi. Feeding was recorded, with some degree of doubt, on Cunninghamia sp. (GRESSITT & KIMOTO 1963). See also item 52. Very possible that this species is identical with C. episcopalis Fairmaire, 1889. C. fokiensis Weise, 1922 (pars) 46(29) Legs distinctly bicolorous or entirely black. 47(62) All femora fulvous, tibiae and tarsi black. 48(49) Species from Japan. Elytra greenish-blue, finely punctate, with interstices broader than diameter of punctures, which are quite deep. Aedeagus – Fig. 13. Length 6.3–7.0 mm. Honshu, Shikoku, Kyushu.

49(48) Continental species.

50(53) Species from China. Elytra violaceous, dark violaceous or violaceous blue.

- 53(50) Species from northern India and Nepal.
- 55(54) Apical antennal segments black, in male not widened or only very slightly so.
- 57(56) Body larger than 6 mm. The following three species are very poorly known.
- 59(58) Elytra dull, finely or moderately punctate, with interspaces wider than diameter of punctures.
- 61(60) Elytral punctures fine and very dense with interspaces distinctly microsculptured. Scutellum black. Length 6 mm. "India orientalis".

 C. alutacea Allard, 1889
- 62(47) At least hind-femora black.
- 63(84) Fore- and mid-legs black with red or fulvous femora, hind legs completely black.
- 64(81) Antennae thin, nitidiform, preapical segments not thickened in male. Prothorax usually with very weakly impressed lines near side margins, sometimes quite indistinct.

- 65(76) Species from Indochina and mostly from China and Taiwan.
- 67(66) Apical antennal segments black.
- 69(68) Body larger than 6.5 mm.

71(70) Interspaces of elytra punctures without irregular convexities, lustrous or microsculptured.

- 73(72) Elytra with strong, very dense punctures, which are wider than interspaces. Prothorax usually with impressed line along lateral margin.
- 75(74) Prothorax without groove near hind angles, but often with very weak impressed line parallel to side margin (Fig. 33). Elytra mostly violaceous or violaceous blue, usually lustrous in both sexes. Prothorax 1.3–1.5 times as wide as long. Segment 1 of mid-tarsi parallel-sided and thinner in male. Aedeagus with acute apex, on underside in apical quarter flat or slightly concave, with very slight central ridge (Fig. 14). Length 7.0–9.5 mm. China (Hupeh, Sichuan, Kwantung, Kweichow, Tibet, Yunnan). See also item 51.

- 76(65) Species from the Himalayas and Myanmar.
- 77(80) Elytra strongly and densely punctate, interspaces not wider than diameter of punctures. Aedeagus with triangular apex.

- 81(64) Preapical antennal segments (usually 8–10) more or less thickened in
- 83(82) Elytra with distinct but shallow postbasal impression, punctures dense and moderately large, interspaces mostly not wider than punctures themselves, colour blue or violaceous. Male: antennal segments 8–10 widened, mostly broader than in preceding species, quite variable in form, segment 11 twice as narrow as 10 (Figs 39–41), segment 1 of fore- and mid-tarsi more elongate and parallel-sided. Aedeagus not constricted preapically, with longitudinal groove on underside (Fig.24). Length 6.8–8.0 mm. China (Yunnan, Xizang).
- 84(63) Legs black or metallic, sometimes articulations may be more or less reddish.
- 85(86) Apex of elytra emarginated. Head and prothorax red-fulvous or more or less darkened or almost black. Legs black. Antennal segment 4 equal to

- 86(85) Apex of elytra broadly rounded. Antennal segment 4 distinctly longer than 3.

Descriptions of new species and taxonomical notes

Cneorane sprecherae sp.nov.

Material examined. Holotype (male): Thailand, Umfang river, 16° 07′N, 90° 00′E, 1000 m, 28.IV-6.V.1991, leg. V. Kuban (NHMB). Paratypes: same locality and date, 5 ex. (NHMB, 2 ex. LM).


Description. Head fulvous with black vertex, antennae, prothorax and scutellum fulvous, elytra bluish-black, without lustre; fulvous margin along side and apex, very narrowly on anterior margin (Fig. 43); underside and legs fulvous, metasternum mostly more or less darkened to black, abdomen sometimes darkened on sides.

Body elongate, parallel-sided. Head impunctate, clypeus triangular with straight anterior margin, interantennal space carinate, frontal tubercles feebly convex, delimited at the rear by a sharp, straight, impressed line. Antennae reach apical slope of elytra, nitidiform, proportions of segments: 8-3-8-8-8-8-8-8-8-6-8, preapical segments about five times as long as wide. Prothorax 1.3 times as wide as long, broadest in anterior quarter, side margins beyond broadest point almost straight, surface evenly convex, lustrous, with very sparse microscopic punctures. Scutellum as wide as long, broadly rounded at apex, lustrous, impunctate. Elytra 1.5 times as long as wide, parallel-sided, with feeble basal convexity and postbasal impression, surface dull, densely microsculptured and very finely punctate, but punctures not very distinct among dense microsculpture. Anterior coxal cavities almost closed. Aedeagus (Fig. 5) narrow, elongate with acute apex and deep longitudinal groove in middle of underside. Segment 1 of fore- and mid-tarsi practically not widened.


Length of body: 4.2-4.5 mm.

Etymology. This species is dedicated to Dr. E. Sprecher, curator of the entomological collection in the Naturhistorisches Museum Basel.


Differential diagnosis. Differs well from almost all species in its very distinct fulvous margin of elytra. The only exception is *C. hirsuta* Kimoto et Takizawa, 1972, which sometimes has two-thirds of lateral margin, a quarter of the sutural and the entire basal margin of the elytra fulvous, but also differs well in pubescent elytra and aedeagus.



Figs 1–9. Aedeagus (d – dorsal, v – ventral, l – lateral): 1 – *C. modesta* Jacoby, d, l, type; 2 – *C. feae* Jacoby, d, l, type; 3 – *C. medvedevi* Beenen, v, l, holotype; 4 – *C. hirsuta* Kimoto, l, v; 5 – *C. sprecherae* sp.nov., v, l, holotype; 6 – *C. fulvicornis* Jacoby, d, l, type; 7 – *C. siamensis* Laboissière, d, l; 8 – *C. minuta* L. Medvedev, d, l, holotype; 9 – *C.variabilis* Kimoto, d, l.



Figs 10–18. Aedeagus (d – dorsal, v – ventral, l – lateral): 10 – C. subcoerulescens Fairmaire, d, l; 11 – C. violaceipennis Allard, v; 12 – C. fokiensis Weise, d; 13 – C. elegans Baly, v, l; 14 – C. intermedia Fairmaire, d, l; 15 – C. rubricollis (Hope), d, l; 16 – C. tibialis Chûjô, d, l, and extreme apex; 17 – C. apicicornis Jacoby, d, l, type; 18 – C. cyanipennis Chûjô, v, d, l.



Figs 19–27. Aedeagus (d – dorsal, v – ventral, l – lateral): 19 – *C. femoralis* Jacoby, d, l; 20 – *C. varipes* Jacoby, d, l; 21 – *C. rugulipennis* Baly, d; 22 – *C. bhutana* sp.nov., v, holotype; 23 – *C. subaenea* Jacoby, d; 24 – *C. crassicornis* Fairmaire, d, l; 25 – *C. abdominalis* Jacoby, d, l, type; 26 – *C. borneensis* Jacoby, d, l, type; 27 – *C. semipurpurea* Jacoby, d, l, type.

Cneorane costipennis sp.nov.

Material examined. Holotype (female): northern Thailand, Chiang Mai, Bella Villa Resort, 18°48′N, 98° 50′E, 500 m, 24–27.VIII.2010, leg. O. Gorbunov (LM).

Paratypes: same locality and date, 3 females (LM), – Thailand, N, Mae Hong Son prov., Soppong env., 600 m, 28.V–2.VI.1999, M. Riha leg., 2 females (JB), – Thailand, N, Mae Hong Son prov., SE of Soppong, 1500 m, 19°27′N, 98°20′E, 23–7.V.1999, M. Riha leg., 1 female (JB), – Thailand, Mae Hong Son prov., Soppong, 1500 m, 19°27′N, 98°20′E, 7–12.V.1996, S. Becvar leg., 4 females (JB).

Description. Head, prothorax, scutellum and legs fulvous, antennae black with three basal and two or three apical segments fulvous, elytra metallic blue to greenish-blue, underside black with last abdominal sternites and pygidium fulvous.

Body elongate ovate. Clypeus with deep, sparse punctures and straight anterior margin, interantennal space quite broad, convex but not carinate, with a few punctures, frontal tubercles triangular, convex, sharply divided posteriorly; with straight, impressed line, as well as from each other; vertex convex, with a few punctures. Antennae reach beyond midpoint of elytra, proportions of segments: 7-3-6-6-6-6-6-6-5-5-8, preapical segments about three times as long as wide. Prothorax 1.5 times as wide as long, broadest in anterior quarter, sides slightly rounded, anterior and posterior angles distinct and produced. Scutellum semicircular, impunctate. Elytra 1.4 times as long as wide, broadest beyond midway, surface with a trace of basal convexity, lustrous, densely punctate, punctures not large but deep, interspaces mostly as wide as diameter of punctures, sharp ridge running from humerus to midpoint of elytra, parallel to side margins; also two additional slight, short ridges, one inside, the other outside, the main ridge (Fig. 47). Prosternum distinct between coxae, anterior coxal cavities half closed. Last abdominal sternite with broadly rounded hind margin.

Length of body: 3.7-4.3 mm.


Differential diagnosis. Differs from all species of the genus in the sharp ridge on the sides of the elytra. This species, having almost closed coxal cavities, appears somewhat atypical for *Cneorane*; conversely, its spermatheca is practically same as in most species of this genus.

Cneorane bhutana sp.nov.

Material examined. Holotype (male): Bhutan, Chasilakha, 6425'[?], 1978, Doriee, Khandu [?collector], with Kimoto's label – *Cneorane varipes* Jacoby (NHMB).

Description. Black, head, three basal antennal segments, prothorax, femora and bases of tibiae of fore- and mid-legs reddish-fulvous, elytra violaceous black.

Body elongate, widening towards the rear. Head impunctate, clypeus short with straight anterior margin, interantennal space with narrow, low ridge, frontal tubercles convex, sharply delimited at the rear by a straight impression. Antennae reach beyond midpoint of elytra, proportions of segments: 11-4-9-10-10-9-10-9-9-9-11, preapical segments about three times as long as wide. Prothorax 1.4 times as wide as long, broadest in anterior third, with distinct anterior angles and produced posterior angles, side margin feebly arcuate, slightly emarginated before hind angles, surface lustrous, convex, impunctate, without impressed lines near lateral margin. Scutellum semicircular, impunctate. Elytra 1.5 times as long as wide, broadest in apical third, surface lustrous,



Figs 28–36. 28 –30. *C. feae* Jacoby: 28, 29 – abdomen ventral and lateral; 30 – anterior tibia and tarsus of male. Figs 31–33. Lateral margin of prothorax with impressions: 31 – *C. cariosipennis* Fairmaire; 32 – *C. femoralis* Jacoby; 33 – *C. intermedia* Fairmaire. 34–36. Spermatheca: 34 – *C. violaceipennis* Allard; 35 – *C. fokiensis* Weise; 36 – *C. sudha* Maulik.

with very slight basal convexity and postbasal impression, finely and not densely punctate, punctures on apical slope very faint, far from distinct, interspaces flat and mostly wider than diameter of punctures. Segment 1 of fore- and mid-legs stick-like, about 2.5 times as long as wide. Aedeagus (Fig. 22) very feebly curved in lateral view, its apex broadly rounded with a very short, small apical tip, underside broadly and deeply concave in apical part.

Length of body: 6.0 mm.

I also have a female from Nepal that might belong to this species, being of practically the same colour and structure of elytra apart from a distinctly punctate apical slope, but at the moment I prefer not to include it in paratypes.

Differential diagnosis. This species alike at *C. rugulipennis* Baly, 1866 and *Cneorane* varipes Jacoby, 1896 but differs in the sculpture of elytra and especially in the form of aedeagus, which has a broadly rounded apex with very short apical tip.

Cneorane modesta Jacoby, 1886

Remark. This species has all the formal characters of *Cneorane*, but differs from typical representatives of the genus in colour and an unusual structure of aedeagus, lacking the typical covering plate on the upperside. Possibly in future, after studying a good series of specimens, *C. modesta* might be removed to another genus.

Cneorane feae Jacoby, 1892 (comb. orig.)

Cneorane rubyana Maulik, 1936 Cassena rubyana: Kimoto, 1989

Remark. This species has closed anterior coxal and basal grooves on the prothorax, and because of these characters it was removed from genus *Cneorane* Baly, 1865 to *Cassena* (KIMOTO 1989). In strictly formal terms, this is correct, but this species differs from all species of *Cassena* and its generic position is far from clear. The basal grooves of the prothorax in this species are very slight and sometimes absent. It also differs from typical *Cneorane* Baly, 1865 in having closed anterior coxal cavities and brushes on the apex of the abdomen. *Cneorane rubyana* Maulik, 1936 is a synonym of *Cneorane feae* Jacoby, 1892 (MEDVEDEV 2002). I exclude this species from the genus *Cassena* and return it to *Cneorane*.


The specimens were collected in Vietnam on *Prunus walicii* and *Fabaceae*, but I am not sure that these are its actual food plants.

Cneorane cariosipennis Fairmaire, 1888

Remark on synonymy. *C. nigripennis* Laboissière, 1922 is without doubt a new synonym of *C. cariosipennis* Fairmaire, 1888. Nearly the only difference is the black colour of the elytra, but this is only a colour aberration, something that is also known in *C. cariosipennis*.

Cneorane cyanipennis Chûjô, 1938 valid species

Remark. This species from Taiwan was united with *C. violaceipennis* Allard (KIMOTO 1969), but differs from it in having black legs with red fore- and mid-femora (*C. femoralis* species group), smaller size, and another form of aedeagus. I studied two males kindly sent to me by Dr. H.Takizawa.




Figs 37–48. 37–42. Five apical antennal segments of males: 37 – C. dilaticornis Chen; 38 – C. subaenea Jacoby; 39–41 – C. crassicornis Fairmaire; 42 – C. braeti Duvivier. 43–46. Elytral pattern: 43 – C. sprecherae sp.nov.; 44–46 – Liroetis ephippiata (Laboissière). 47–48. Elytral sculpture: 47 – C. costipennis sp.nov.; 48 – C. birmanica Jacoby.

Species excluded from the Cneorane

Liroetis ephippiata (Laboissière, 1930) (comb.nov.)

Remark. Cneorane ephippiata already differs from other species of this genus in its fulvous elytra with black spots (Figs 44–46); it also has very narrow epipleurae, typical of the genus Liroetis Weise, 1889. Females of this species have a very complicated postpygidium with four acute teeth (Fig. 51). Spermatheca – Fig. 52. I have only females in my collection, so the aedeagus was studied and figured by my colleagues Dr. Eva Sprecher (Fig. 49) and Mr. Ron Beenen (Fig. 50). Its structure differs sharply from all Cneorane, but is typical of Liroetis. Because of all the characters mentioned above I place this species in the genus Liroetis.



Figs 49–52. *Liroetis ephippiata* Laboissière: 49 – aedeagus dorsal and lateral, by E. Sprecher; 50 – aedeagus lateraly by R. Beenen; 51 – postpygidium; 52 – spermatheca.

Charaea abdominalis (Jacoby, 1890) (comb.nov.)

Remark. Cneorane abdominalis Jacoby, 1890, the type of which was studied, must be placed in another genus, possibly Charaea Baly, 1878. The prothorax of this species is scarcely broader than long, with sides widened midway, anterior coxal cavities closed and a type of coloration unusual for Cneorane. Aedeagus – Fig. 25.

Hyphaenia borneensis (Jacoby, 1894) (comb.nov.)

Remark. Cneorane borneensis Jacoby, 1894 has several characters quite atypical of the genus Cneorane. The antennae of this species have basal segments practically glabrous and the following ones densely pubescent, with a few erect hairs; further, segments 2 and 3 are short and equal, while segment 4 is much longer than the two preceding combined; hind angles of prothorax broadly rounded; aedeagus very thin and curved twice in lateral view (Fig. 26), differing sharply from all known Cneorane. The type was studied. As a preliminary measure, I place it in the genus Hyphaenia.

Hyphaenia semipurpurea (Jacoby, 1884) (comb.nov.)

Remark. Jacoby (1884) placed this species in the genus *Cneorane*, with some doubt, and noted that he could not indicate any important characters to "justify the establishment of another genus for its reception", but also mentioned that it resembles the genus *Theopea* Baly, 1864. The male of this species has antennal segment 4 as long as the three preceding combined and the punctures of the elytra are partly arranged in semi-regular rows. I think that these characters are enough to exclude the species in question from *Cneorane*; it appears to be quite near *C. borneensis* Jacoby, 1894, mentioned above. Its aedeagus is also thin but not curved in lateral view (Fig. 27). The type was studied.

Acknowledgements

I am very grateful to Dr. Eva Sprecher, Dr. Haruo Takizawa, Dr. Jan Bezděk and Mr. Ron Beenen for their invaluable help in the preparation of this publication.

References

- CHEN S.H. (1964): New genera and species of Galerucinae from China. Acta Ent. Sinica 13: 201-211
- CHEN S.H. & JIANG S. (1976): New leaf beetles from West China. In: CHEN YU & WANG JIANG, Acta Ent. Sinica 19: 205–219.
- CHÛJÔ M. (1966): Chrysomelid beetles from Northeast Nepal. Mem. Fac. Educ. Kagawa Univ. 2(45): 1-35.
- GRESSITT J.L. & KIMOTO S. (1963): The Chrysomelidae of China and Korea, Part 2. Pac. Ins. Monogr. 1B: 301–1026.
- Кімото S.H. (1969): Notes on Chrysomelidae of Taiwan II. Esakia 7: 1-65.
- KIMOTO S.H. (1977): Ergebnisse der Bhutan-Expedition 1972 des Naturhistorischen Museums in Basel, Coleoptera: Fam. Chrysomelidae Subfam. Galerucinae. Entomologica Basiliensia 2: 351–392.
- KIMOTO S. (1989): Chrysomelidae of Thailand, Cambodia, Laos and Vietnam. VI. Galerucinae. Esakia 27: 1–241.
- KIMOTO S. & TAKIZAWA H. (1972): The Chrysomelid-beetles of Nepal, collected by the Hokkaido University scientific expedition to Nepal Himalaya, 1968. Part 1. Kontyû 40: 215–223.
- KIMOTO S. & TAKIZAWA H. (1994): Leaf beetles (Chrysomelidae) of Japan. Tokai University Press, Tokyo, 539 pp.
- KIMOTO S. & TAKIZAWA H. (1997): Leaf beetles (Chrysomelidae) of Taiwan. Tokai University Press, Tokyo, 581 pp.
- MAULIK S. (1936): The Fauna of British India including Ceylon and Burma. Coleoptera. Chrysomelidae, Galerucinae. Taylor and Francis, London, 648 pp.
- Medvedev L.N. (1992): Chrysomelidae from the Nepal Himalayas, III (Insecta, Coleoptera). Stuttg. Beitr. Naturk. (Ser. A.) 485: 1–36.
- Meducedev L.N. (2002): Jacoby's types of Chrysomelidae (Coleoptera) from Burma in the Museo Civico di Storia Naturale "Giacomo Doria", Part 3. Ann.Mus. Stor. Nat., 94(2): 249–264
- MEDVEDEV L.N. (2004): New species of Chrysomelidae (Coleoptera) from Nepal. Veroeffentlichung Naturkundemuseum Erfurt 23: 203–206.
- MEDVEDEV L.N. (2009): Revision of the genus Cassena Weise, 1892 (Coleoptera, Chrysomelidae). Entomologica Basiliensia et Collectionis Frey 31: 219–238
- MEDVEDEV L.N. & DANG THI DAP (1982): *Trophical connections of Chrysomelidae in Vietnam*. In: Animal world of Vietnam. Moscow, Nauka, pp 84–97 (in Russian).
- Medvedev L.N. & Sprecher-Uebersax E. (1999): Katalog der Chrysomelidae von Nepal. Entomologica Basiliensia 21: 261–354

Author's address:

Prof. Lev N. Medvedev Institute for Problems of Ecology and Evolution Russian Academy of Sciences Leninsky prospect 33 Moscow 119071 RUSSIA

E-mail:lev.n.medvedev@mail.ru