Zeitschrift: Mitteilungen der Entomologischen Gesellschaft Basel

Herausgeber: Entomologische Gesellschaft Basel

**Band:** 31 (1981)

Heft: 1

**Artikel:** Annotated list of sawflies (Hym., Symphyta) from Kanton Unterwalden.

Part II.

**Autor:** Liston, A.D.

**DOI:** https://doi.org/10.5169/seals-1042558

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 24.11.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# ANNOTATED LIST OF SAWFLIES (HYM., SYMPHYTA) FROM KANTON UNTERWALDEN. Part II.

#### A.D. Liston

For an explanation of collecting dates and localities, the introduction to the first part of this record should be consulted (LISTON, 1980). The main sources from which I have drawn distributional and biological data are not repeated if they are already given in the previous paper, unless they are of particular importance or obscurity. The months given for the flight of adult insects are based on dates of capture throughout the whole of Europe but they should apply fairly well to Switzerland where species often appear over an unusually long period of time as suitable conditions gradually move up to higher altitudes.

## Tenthredinidae

Nematinae (concl.)

## Nematus PANZER

These species are often multi-brooded and they were therefore the best represented Nematinae at lower levels in July. Keys to the European forms are given by MUCHE (1974: 5-10) and BENSON (1958: 211-231).

## N. fagi ZADDACH (= mimus KONOW)

1♀, Dallenwil/NW. 12.vii.78. In copse containing Fagus, the larval foodplant. Larve is a solitary, leaf-edge feeder. Adult flies in two broods: V-VI and VII-VIII. Northern, Central and Southern Europe (found in Albania), but commonest in the south-east. Not apparently previously recorded in Switzerland.

## N. ferrugineus FOERSTER

29, 10, Ende der Welt/OW. 15. vii. 79. On tips of Salix twigs (cinere and purpurea L.). Larva on rough-leaved species of willow, and possibly Betula. Northern and Central Europe, and Siberia. In two broods: V-VI and VII-IX. New record for Switzerland.

#### N. melanaspis HARTIG

Larvae abundant on Populus nigra L. at Hergiswil/NW, 4.vii.79. Presumably these larvae had resulted from eggs laid by the spring (V-VI) generation of sawflies. The second generation flies VII-VIII. The larvae of this species are distinctive and are easily detected because they feed gregariously. No adults were bred from these Swiss larvae, but I already know them well from experience of melanaspis in Britain. Distributed over North and Central Europe and Siberia. Widely recorded in Switzerland from Wallis, Solothurn and Zürich.

#### N. myosotidis (FABRICIUS)

2 ♀, Wiesenberg/NW. 12. vii. 78. 2♂, Dallenwil/NW. 6. vii. 79. ? Larvae

on Vicia cracca L. near Dallenwil, 12. vii.78. According to LORENZ and KRAUS (1957) these larvae seem to belong to another species, namely papillosus (THOMSON, 1871), but this species was placed as a synonym of myosotidis by KONTUNIEMI (1951). Larvae of the true myosotidis are otherwise recorded from Trifolium ssp. and Onobrychis viciifolia SCOPOLI. A very common species throughout Europe to W. Siberia in two or more broods: V-IX, Throughout Switzerland.

## S. oligospilus FOERSTER (=glutinosae CAMERON)

1 ♀, Wolfenschiessen/NW. 14. vii. 78. From a grassy verge beside the main road, but probably flew from willows growing beside the River Aa. Larva on Salix spp., Betula and ? Ulmus. Double brooded: V-VI and VII-VIII. Hitherto recorded from Graubünden and Luzern.

## N. ribesii (SCOPOLI)

4 ♀, and larvae, Hergiswil/NW. 14.vii.78. In gardens. All Europe, east to the Caucasus and introduced to North America. In two or three broods: IV-IX. Previously noted from Zürich, Bern, Neuenburg and Graubünden.

# N. spiraeae ZADDACH

5 \( \), Wiesenberg/NW. 15. vii. 78. Larvae were found to be very abundant on Aruncus silvester (KOSL.) growing in woodland glades in Fagus forest at Wiesenberg and also below Niederrickenbach/NW. The adults at Wiesenberg were flying in a deeply shaded area in the late afternoon. Common in most of Central Europe and reaching quite far north (Finland). In many broods: IV-IX. Parthenogenetic. Not previously recorded in Switzerland.

#### N. tibialis NEWMAN (= hortensis HARTIG)

Two Nematine larvae of what is almost certainly <u>tibialis</u> were found feeding singly on the leaf edges of Robinia pseudacacia L. which were growing near the Winkelried monument in the main square of Stans/NW. 11.vii.79. This species is an introduction from North America which is now found in most areas where this tree is planted in Europe. Widely recorded in Switzerland. I have seen larvae beside a motorway in Baselland (4.vii.79). Also found in Geneva, Bern, Zürich and Wallis.

## N. viridis STEPHENS

1 \, D\, D\, D\, orfli/NW. 14. vii. 78. This is a very pale specimen, with a completely pale scutellum (a point which often serves to distinguish it easily from bergmanni DAHLBOM, which is superficially similar) and completely green abdomen. Forms from Southern Europe and warmer areas generally, as well as late summer specimens, tend to have less extensive black markings than spring and northern forms. Larva on Betula and perhaps Salix. Not, surprisingly, recorded from Switzerland until now. Very common in most of North and Central Europe and also found in the South (Turkey). At least two broods: IV-V and VII-VIII.

#### Pachynematus KONOW

Unlike <u>Nematus</u>, fewer of these species are multibrooded, so it is not surprising that most of the specimens which I obtained came from subalpine

and alpine levels. MUCHE (1974:70-72) gives a key to Central European forms, and BENSON (1958: 231-41) a key to those that occur in Britain.

## P. apicalis (HARTIG) (= lichtwardti KONOW)

1 \,\text{\$\pi\$, Tr\"u\text{ebsee}/NW. 10.vii.79. Probably over most of Europe but? subalpine in Switzerland. Recorded from Britain, including Ireland, Germany (East and West), Sweden, Finland, North-West European USSR and Bulgaria. Larva on Gramineae. Flies IV-VII. The only Swiss record that gives a specific locality is for Graub\"u\text{onden}.

# P. declinatus (FOERSTER) (= inopinatus BENSON nec LINDQVIST)

1 of, Trüebsee/NW. 10. vii. 79. From Carex, but the foodplant of this species is not known for certain. Apparently distributed only in Central Europe: East and West Germany, and subalpine and alpine Switzerland. Flies V-VII. Recorded from Wallis and Graubünden in Switzerland. 8th tergite of male figured by BENSON (1955).

## P. extensicornis (NORTON) (= truncatus BENSON)

2 ♀, Ende der Welt/OB. 15.vii.79. Holarctic species occurring commonly over Central and Northern Europe. Larva on various Gramineae. Flies V-VII. Previously recorded from Graubünden.

# P. laevigatus (ZADDACH & BRISCHKE) (= chambersi BENSON)

1 \( \), Rinderbühl-Stockhütte/NW. 15. vii. 79. Central and Northern European species (Sweden, England, Finland, Germany and Switzerland). Rather scarce. The synonymy with chambersi, its previously unknown male, was made by BENSON (1967). I cannot agree with MUCHE (l.c.) who synonymises chambersi with clitellatus. It seems to me that though the females can only be distinguished with difficulty, the differences shown by the males are too profound to represent mere variation in clitellatus. Larva probably on Gramineae. Flies V-VII. Previously recorded from the St. Gotthard by HELLÉN (1974).

#### P. obductus (HARTIG)

1  $^{\circ}$ , Fräkmuntegg/NW. 5.vii.79. All Europe, North America. Flies IV-IX in two or more broods. Larva on many Gramineae. Probably widespread in Switzerland and recorded from Wallis, Graubünden, Bern, Zürich and Tessin.

## P. smithae ROSS (= smithiae: error)

1 of, Trüebsee/NW. 11. vii. 78. An interesting species which was described from males found on Mount Washington, New Hampshire, USA and later found by BENSON in Lapland, Northern Scotland and the high Swiss Alps. A specimen was also found by BRIDGMANN on the Norfolk coast in the south of Britain in the late 19th century. The female was described by BENSON, 1967, but once again it is difficult to distinguish from clitellatus. Larva probably on Carex, but not yet described. Flies V-VII.

## Acknowledgement

Once again I thank Miss G.A. LISTON for her care in preparing the abstract for this work and also for translating the German language texts from which I drew some information.

# Zusammenfassung

Dieser Bericht schliesst die vorbereitende Untersuchung der Hymenoptera Symphyta aus dem Kanton Unterwalden ab, die der Autor im Sommer der Jahre 1977-79 durchführte.

Der Hauptteil der Angaben über die gefundenen Arten ist schon in dieser Zeitschrift erschienen (LISTON, 1980). Im vorliegenden Bericht werden 15 Arten der Unterfamilie Nematinae behandelt. Anscheinend sind vier dieser Arten, Nematus ferrugineus, N. fagi, N. spiraeae und N. viridis, erst jetzt in der Schweiz festgestellt worden. Künftige Forschungen in diesem Gebiet dürften zur Entdeckung weiterer interessanter Arten führen.

## References


BENSON, R.B.	1955	Some high alpine Nematine Sawflies (Hym., Tenthredinidae). Entomologist's mon. Mag., 91: 103-5.
-	1958	Hymenoptera Symphyta. Handbks. Ident. Br. Insects, 6(2c): 139-252.
-	1967	Pachynematus laevigatus ZADD. as a British Sawfly and the females of P. chambersi BEN-SON, P. smithiae ROSS and P. sulcatus BENSON. Entomologist's mon. Mag., 103: 141-3.
HELLÉN, W.,	1974	Die Nematinen Finnlands III (Hymenoptera, Tenthredinidae). Gattung <u>Pachynematus</u> KONOW. Notul. Ent., 54: 65-80.
KONTUNIEMI, T.	1951	Zur Kenntnis des Lebenszyklus der Sägewespen in Finnland. Acta ent. Fenn., 9: 1-92.
LISTON, A.D.	1980	An Annotated List of Sawflies (Hym., Symphyta) from Kanton Unterwalden, Central Switzerland. Part 1. Mitt. ent. Ges. Basel, (N.F.) 30: 4-28.
LORENZ, H. and KRAUS, M.	1957	Die Larvalsystematik der Blattwespen. Abh. Larvalsyst. Insekt., 11, viii+ 340 pp.
MUCHE, W.H.	1974	Die Nematinengattungen Pristiphora LATREILLE, Pachynematus KONOW und Nematus PANZER (Hym., Tenthredinidae). Dt. ent. Z., (N.F.) 21: 1-137.

## Author's address:

A.D. Liston 99 Clermiston Road Edinburgh. EH12 6UU

Scotland

Addendum: The easiest and most reliable way to separate Nematus oligospilus from viridis is by the comparative lengths of the sawsheath and cerci (Figs. 1 & 2).



Figs. 1 & 2: Sawsheath of  $\varphi$  viewed dorsally

in 1: Nematus oligospilus

2: N. viridis