Mitteilungen der Entomologischen Gesellschaft Basel			
erausgeber: Entomologische Gesellschaft Basel			
32 (1982)			
2			
Insects from high-altitude summer snow in Austria, 1981			
Liston, A.D. / Leslie, A.D.			
https://doi.org/10.5169/seals-1042475			

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

# Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

STECK, T. 1893 Beiträge zur Kenntnis der Hymenopterenfauna der Schweiz. I. Tenthredinidae. Mitt. Schweiz. Ent. Ges., 9: 1-45.

Anschriften der Verfasser:

Andrew D. Liston 99 Clermiston Road

Edinburgh EH12 6UU Schottland

Bruno Peter Rigistr. 169 CH-<u>6340 Baar</u> Schweiz

INSECTS FROM HIGH-ALTITUDE SUMMER SNOW IN AUSTRIA, 1981

A.D. Liston & A.D. Leslie

From 20.8. to 9.9.1981, the authors took a mountaineering holiday in Austria. We had no plans for serious collecting because of the late date of our visit, but once in the mountains we were fascinated by the diversity and occasional abundance of the insects seen dead or crawling over snow patches and Firn.

Little data has been published on the phenomenon of insects in such seemingly inhospitable areas. MASUTTI (1978) has characterised the insect fauna of late winter and spring snow in the Carnic and Julian Alps (N. Italy). His studies revealed that certain insects, e.g. Isotoma hiemalis (Collembola), <u>Boreus</u> ssp. (Mecoptera), <u>Trichoptera</u> ssp., <u>Trichocera hiemalis</u> (Diptera), are typically active during winter on alpine snow. Other groups of insects, particulary Coleoptera Carabidae, have been indicated as characteristic of "new ground" at receding snow patch edges (JANETSCHEK, 1949).

In summer, however, by far the greatest number of insect species and individuals found on snow are not resident to areas of permanent snow at all, but are either members of summer subalpine-alpine communities or are of more distant lowland origin. It is probably true, as has often been stated in the past, that most organisms found inactive on high-altitude snow have merely been blown or carried up into the mountains by wind or convection currents. This is almost always the right explanation for biological debris on snow in non-alpine areas, e.g. the Scottish Mountains(BRUCE in THORN-LEY, 1896). But for certain types of summer, alpine insects assemblages we think that large scale migrations of lowland species, rather than passive/ accidental transport by wind, play an important role in introducing insects to the snowfields in some areas. Our data from the Alps of Kärnten (Glockner-Gruppe) seem to support this interpretation. Below are listed, in two sections, the insects which we collected in 1) Hochstubai, Nordtirol.

2) Glockner-Gruppe, Kärnten.

For each species a few voucher specimens were preserved and written notes taken on numbers of individuals and their state of preservation, i.e. living, torpid, dead (and undamaged: therefore probably recent arrivals on the snow), dead and badly preserved (probably earlier arrivals). Authorities are named where identifications were made by entomologists other than the authors.

# I. Hochstubai, Nordtirol, 23.8.-26.8.1981

- A: Firn snow on Wilder Freiger. Some insects found on very summit of this mountain. 23.8.
- 2 <u>Plutella xylostella</u> (L.)(Lepidoptera, Plutellidae). 3500m. Alive, 1 other seen. Det. S.WHITEBREAD.
- 3 <u>Zeiraphera diniana</u> (Gn.)(Lepidoptera, Tortricidae). 1 at 3000m, 2 at 3500m. All alive, no others seen. Det. S.WHITEBREAD.
- 2 <u>Episyrphus balteatus</u> (Deg.)(Diptera, Syrphidae). 3300m and 3500 m. Dead (undamaged), no others. Det. K.WATT.
- B: Firn snow and lower snow patches on walk from Nürnberger Htt. Ost. Feuerstein - Bremer Htt. 25.8.
- 5 <u>Zeiraphera diniana</u> (Gn.). 3100m. All alive, no others. Det. S.WHITE-BREAD.
- 1 Oscinella frit (L.)(Diptera, Chloropidae). Below Ost. Feuerstein (Simmingferner).2800-3000m. Alive and dead, a few flying over snow. Countless numbers.
- C: Single large snow patch above Bremer Htt. at 2500m, 25.8.
- 1 Leuctra sp. (Plecoptera). Alive, no others.
- 1 Aphid sp. (Aphidoidea). Alive, no others.
- 2 Episyrphus balteatus (Deg.). Dead, no others.
- 1 <u>Bibio pomonae</u> (F.)(Diptera, Bibionidae). Dead and badly preserved, no others.
- 2 Muscidae sp. (Diptera). Dead (undamaged), no others.
- 2 <u>Myrmica ruginodis</u> Nyland (Hymenoptera, Formicoidea). Winged sexuals. Dead (undamaged-badly damaged), 4 or 5 others seen. Det. B.BOLTON.
- 1 Apis mellifera L. (Hymenoptera, Apoidea). Worker. Dead (very badly preserved), no others. Being fed on by Aphodius sp. A &below).
- 1 <u>Aphodius</u> sp. A (Coleoptera, Scarabaeidae). Alive, feeding on body of <u>Apis mellifera</u> (above), no others.
- 1 Aphodius sp. B. Alive, crawling on snow, no others.
- 1 Chrysomelidae sp. (Coleoptera). Dead (undamaged), no others.
- D: Snow patches on Habicht, above Innsbrucker Htt. 2500m. 26.8.

- 1 Trichoceridae sp. (Diptera). Alive, no others.
- II. Glockner-Gruppe, Kärnten, 2.9.81.

Fresh snow lying on Firn.

- 1 <u>Autographa gamma</u> (L.)(Lepidoptera, Noctuidae). Johannisberg (3100m). Torpid and dead, 3 others seen, all with very rubbed wings.
- 2 <u>Scotia ipsilon</u> HUFN. (Lepidoptera, Noctuidae). Above Oberwalder Htt. (3100m). Torpid, 11 others seen, good condition.
- 1 <u>Sympetrum sanguineum</u> MUELLER (Odonata). Johannisberg (3100m). Dead, badly broken, no others. Det. I. BALDWIN.
- 1 Chrysopa carnea L. (Neuroptera). Johannisberg (3000m). Dead, 3 others.
- 1 Aphid sp. Very large black species. Johannisberg (3000m). Dead, no others.
- 1 <u>Eristalis tenax</u> (L.)(Diptera, Syrphidae). Johannisberg (3100m). Torpid, one other seen. Det. K.WATT.
- 4 <u>Episyrphus balteatus</u> (Deg.). Johannisberg (2900-3200m). All dead, some apparently fresh. Very many: probably hundreds.
- 1 <u>Scaeva pyrastri</u> (L.)(Diptera, Syrphidae). Oberwalder Htt. to 3200m on Johannisberg. All dead, badly damaged. About 50 others seen.
- 2 <u>Coccinella septempunctata</u> L. (Coleoptera, Coccinellidae). Johannisberg (3100m). Dead, 2 others seen.

# Discussion

Nearly all of the species from the peaks of the Stubai were obviously lowland or subalpine in origin and had probably been blown up to the snowfields during the day that we observed them. The weater was poor, with low cloud, high winds, occasional falls of snow and freezing temperatures. It is unthinkable that any of the species found near the summits of the Wilder Freiger or Ost. -Feuerstein had arrived more than half an hour before they were observed. Any earlier arrivals would have been buried in the fresh powder snow after they succumbed to the cold. In terms of numbers (apart from Oscinella frit), we saw far fewer insects in the Stubai than we did in the Glockner area. This was probably partly due to the differences in weather conditions (Figs. 1-4). The only abundant insect in the Stubai was O.frit, which gave the snow a dirty appearance on parts of the Simmingferner.

The <u>Plutella xylostella</u> found on the Wilder Freiger may have been on migration, but could equally well have been transported by wind from a local population. The moth is cosmopolitan and a well-known migrant. Its larva has been noted at 2800m in the Alps.

S. WHITEBREAD and E. de BROS have both told me of finding hundreds of <u>Zeiraphera diniana</u> on permanent snow at 3400m on the Mettelhorn, near Zermatt, 31.7.1981. E. de BROS also reports having 200 dead <u>diniana</u> in his Heath trap during a single night at Zermatt. According to S. WHITE-BREAD, the specimens on the Mettelhorn were concentrated in one place only: on a col immediately above an almost sheer drop to the larch forest below. Our Austrian specimens were found  $2^{1/2}$  km the nearest area of <u>Larix</u>,

in the Langental. Loss of adults to high altitudes and subsequent death may be an important control acting on this defoliator in alpine areas.

The species from the relatively low snow patch above the Bremer Hütte are likewise predominantly from lower altitudes. <u>Aphodius</u> sp. A, however, appeared to be making use of the insect debris as a food source. Only the Trichoceridae sp. may be a real representative of the permanent snow community.

Of the insects from the Glockner-Gruppe, at least the two Noctuidae are well documented migrants (REZBANYAI, 1979, 1980), particularly in the Alps, where they are amongst the commonest moths taking part in mass migrations. The vast numbers of <u>Episyrphus balteatus</u> may also indicate that migrations of this hoverfly passed through the area. It is interesting to note that on the late afternoon of 2.9.81 we observed several <u>Vanessa atalanta</u> flying steadily northwards over the snowfield below the Johannisberg.

Other species such as <u>Chrysopa carnea</u> and <u>Sympetrum sanguineum</u> may have been individual "wanderers" which had unsuccessful post-emergence dispersal flights. Dispersal flights of this type are well documented in <u>C.carnea</u> (DUELLI, 1980).

The differences in composition of insect assemblages on snow between the Hochstubai and the Glockner-Gruppe may partly be due to the fact that the Johannisberg (Glockner) lies at the head of a major alpine valley and its surrounding snowfields are the highest point on what is otherwise probably a useful migration route between south-eastern and more central Europe. Conversely, there are several easier routes round the Hochstubai rather than over these mountains.

Finally, the main difference between the late summer insect assemblages described here and the spring ones found by MASUTTI is the absence of Hymenoptera Symphyta from the summer ones. This was only to be expected: the only adult sawflies that we encountered during the whole visit were <u>Athalia rosae</u> (L.), <u>Tenthredo aegra</u> ENSLIN and <u>T.velox</u> (F.)(all on the Franz-Josef-Höhe, Kärnten, 2100m, 1.9.81). Even these were very scarce.

## Acknowledgements

The Merchant Company of Edinburgh made our visit possible with a grant. We thank S. Whitebread (Herznach) in particular for his help in determination. K. WATT (Aberdeen), A. BOLTON (London) and I.BALDWIN (Edinburgh) also assisted with identification.

## Summary

Insects were recorded from high altitude snow in the Hochstubai and in Kärnten, Austria, during August and September 1981. Some of these summer assemblages contained remarkable numbers of migrant species (e.g. <u>Scotia</u> <u>ipsilon</u> and <u>Autographa gamma</u>). An <u>Aphodius</u> species was found scavenging on dead insect remains in the Stubai. Sawflies, typically common in spring snow assemblages, were absent.



Fig.	1:	West Feuerstein Ridge from the Wilder Freiger, Stubai, 23.8.81
Fig.	2:	On the Aperer Feuerstein, Stubai, 25.8.81
Fig.	3:	The Oberwalder Hütte, Kärnten, 2.9.81.
Fig.	4:	The Grossglockner Ridge from the snowfields below the Johannis-
		berg, Kärnten, 2.9.81.

References

BRUCE, W.S. in THORNLEY, A.	1896	On some Coleoptera from the summit of Ben Nevis, collected by Mr.W.S.Bruce. Ann. Scott. Nat. Hist., 5, 28-37.
DUELLI, P.	1980	Adaptive dispersal and appetitive flight in the green lacewing, <u>Chrysopa carnea</u> . Ecol. Ent., 5(3), 215-220.
JANETSCHEK, H.	1949	Tierische Sukzessionen auf hochalpinem Neu- land. Ber. natmed. Ver. Innsbruck, 48/49, 1-215.
MASUTTI, L.	1978	Insetti e nevi stagionale. Riflessioni su reperti relativi alle Alpi Carniche e Giulie. Boll. Ist. Ent. Univ. Bologna, 34, 75-94.
REZBANYAI, L.	1979	Nachtaktive Wanderfalter (Macroheterocera) in den Zentralschweizer Alpen 1972-1975: Fangergebnisse einer Lichtfalle am Brisen- Haldigrat NW 1920m ü.M. Atalanta, 10, 245-279.
-	1980	Die Insektenfauna des Hochmoores Balmoos bei Hasle, Kanton Luzern. II. Macrolepidop- tera (Grosschmetterlinge). Ent. Berichte Luzern, 3, 15-76.
Author Addresses:		A.D. Liston 99 Clermiston Road Edinburgh EH12 6UU Scotland

A.D.Leslie 15 Old Church Lane

Edinburgh EH 15 Scotland