

3. Recurrences.

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **4 (1958)**

Heft 1: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

powers of Ψ are lowered into subscripts; thus $\Psi^{(1)}$ would then be written $\Psi_1(t)$.

If (18) and (19) are substituted in (13), there results the following umbral identity:

$$(20) \quad \Psi (1 - \cos \Psi_s) + 2 \frac{\partial}{\partial t} \left(\frac{1 - \cos \Psi_s}{\Psi} \right) = 2 \sin \Psi_s * \sin \Psi_s ,$$

where the asterisk (*) indicates umbral multiplication.

For the cases $r = 2, 3$, rather extensive calculations show that umbral identities of the same form exist. We may therefore state the following result which is implied by the non-linear equation (13).

Theorem 1: "Let Ψ, X, Φ be respectively the umbrae of the sequences $\{\Psi_{2k-1}(t)\}$, $\{X_{2k-1}(t)\}$, and $\{\Phi_{2k-1}(t)\}$. If γ is one of these umbrae, then the following umbral identity holds:

$$(21) \quad \gamma (1 - \cos \gamma_s) + 2 \frac{\partial}{\partial t} \left(\frac{1 - \cos \gamma_s}{\gamma} \right) = 2 \sin \gamma_s * \sin \gamma_s .$$

3. RECURRENCES.

It is clear that (20) implies a recurrence relation for the functions $\Psi_j(t)$, and indeed, Theorem 1 yields the following.

Theorem 2: "Let $\gamma_j(t)$ be $\Psi_j(t)$, $X_j(t)$ or $\Phi_j(t)$; then the following recurrence holds:

$$(22) \quad \frac{d}{dt} \gamma_{2n-1}(t) + \frac{1}{2} \gamma_{2n+1}(t) = \sum_{k=0}^{n-1} \binom{2n}{2k+1} \gamma_{2k+1}(t) \gamma_{2n-2k-1}(t) ,$$

and hence $\gamma_{2n+1}(t)$ is a polynomial in $\gamma_1(t)$ and its derivatives up to order n ."

This result, in turn, implies the following

Theorem 3: "Let $\rho_{2k-1}(n)$ be either of the arithmetical functions $\beta_{2k-1}(n)$ or $\xi_{2k-1}(n)$ defined by (8) and (10) respectively; then $\rho_{2k-1}(n)$ satisfies a recurrence relation of the form:

$$(23) \quad \rho_{2k+1}(n) - n \rho_{2k-1}(n) = 2 \sum_{s=0}^{k-1} \sum_{j=1}^{n-1} \binom{2k}{2s+1} \rho_{2s+1}(j) \rho_{2k-2s-1}(n-j) ,$$

for all n and $k \geq 1$. Moreover, the arithmetical function $\zeta_{2k-1}(n)$ defined by (9) satisfies the recurrence

$$(24) \quad \zeta_{2k+1}(n) - 2n \zeta_{2k-1}(n) = 2 \sum_{s=0}^{k-1} \binom{2k}{2s+1} \left\{ U_{k-s} \zeta_{2s+1}(n) + \right. \\ \left. + U_{s+1} \zeta_{2k-2s-1}(n) + \sum_{j=1}^{n-1} \zeta_{2s+1}(j) \zeta_{2k-2s-1}(n-j) \right\}$$

where U_k is defined by (11) and $n, k \geq 1$."

Incidentally, the comparison of coefficients which yields (24) also gives:

$$(25) \quad U_{n+1} = 2 \sum_{k=0}^{n-1} \binom{2n}{2k+1} U_{k+1} U_{n-k}, \quad n \geq 1,$$

which is equivalent to a result given by NIELSEN [7].

Finally, the case $r = 1$, has been discussed by VAN DER POL [1] who finds an expression analogous to (22) as follows:

$$(26) \quad \frac{d}{dt} h_{2n-1}(t) + \frac{2n+3}{4n+2} h_{2n+1}(t) = \sum_{k=0}^{n-1} \binom{2n}{2k+1} h_{2k+1}(t) h_{2n-2k-1}(t), \quad n \geq 1,$$

where,

$$(27) \quad h_{2n-1}(t) = \frac{(-1)^n B_n}{4n} \alpha_{2n-1}(t),$$

$\alpha_{2n-1}(t)$ being defined by (2).

We find the analogue of (24) for this case to be:

$$(28) \quad \frac{2k+3}{2k+1} \sigma_{2k+1}(n) - 2n \sigma_{2k-1}(n) = 2 \sum_{s=0}^{k-1} \binom{2k}{2s+1} \left\{ V_{k-s} \sigma_{2s+1}(n) + \right. \\ \left. + V_{s+1} \sigma_{2k-2s-1}(n) + \sum_{j=1}^{n-1} \sigma_{2s+1}(j) \sigma_{2k-2s-1}(n-j) \right\}, \quad n \geq 1.$$

Corresponding to (25), we find

$$(29) \quad V_{n+1} = \frac{4n+2}{2n+3} \sum_{k=0}^{n-1} \binom{2n}{2k+1} V_{k+1} V_{n-k},$$

which is equivalent to a known recurrence for the BERNOLLI numbers [8].

4. THE FUNCTIONS $\Psi_{2k-1}(t)$, $X_{2k-1}(t)$, $\Phi_{2k-1}(t)$ AS DOUBLE SUMS.

The results which are stated as (4), (5), (6) follow readily from (1) and (2) which are known to be equivalent (see [1], [2]). It is to be observed first that a comparison of (4) and (5) with (1) taking into account (27) gives the relations:

$$(30) \quad \Psi_{2k-1}(t) = h_{2k-1}(t/2) - h_{2k-1}(t) = V_k(\alpha_{2k-1}(t/2) - \alpha_{2k-1}(t)),$$

$$(31) \quad X_{2k-1}(t) = 2^{2k} h_{2k-1}(2t) - h_{2k-1}(t) = V_k(2^{2k} \alpha_{2k-1}(2t) - \alpha_{2k-1}(t)).$$

From (4) and (6) we also have,

$$(32) \quad \Phi_{2k-1}(t) = 2^{2k} \Psi_{2k-1}(2t) - \Psi_{2k-1}(t).$$

By (30), we may write

$$(33) \quad \Phi_{2k-1}(t) = -V_k(\alpha_{2k-1}(t/2) - (2^{2k} + 1)\alpha_{2k-1}(t) + 2^{2k}\alpha_{2k-1}(2t)).$$

Thus, our functions (4), (5), (6) are expressed in terms of $\alpha_{2k-1}(u)$. These relations in conjunction with (1) and (2) identify them with $(4)_1$, $(5)_1$, and $(6)_1$ respectively.

It is of interest to note that (31) with $k = 2$ permits, with the aid of a result of VAN DER POL [1], the deduction of Jacobi's famous theorem on the number of representations $r_8(n)$ of the integer n as the sum of eight squares. Thus,

$$(34) \quad 240 X_3(t) = 16 \alpha_3(2t) - \alpha_3(t) = 15 \theta_0^8(0, q)$$

where $q = \exp(-t)$. Hence,

$$\theta_0^8(0, q) = 16 X_3(t) = 1 + 16 \sum_{n=1}^{\infty} q^n \zeta_3(n),$$

and

$$\theta_3^8(0, q) = 1 + 16 \sum_{n=1}^{\infty} (-1)^n q^n \zeta_3(n).$$