

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 5 (1959)
Heft: 3: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SPECIAL CASE OF KUMMER'S CONGRUENCE
Autor: Carlitz, L.
DOI: <https://doi.org/10.5169/seals-35487>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A SPECIAL CASE OF KUMMER'S CONGRUENCE
 (mod 2^e)

by L. CARLITZ, Duke University, Durham

(Reçu le 30 janvier 1959.)

Put

$$\frac{x}{\sinh x} = \sum_1^{\infty} D_{2n} \frac{x^{2n}}{(2n)!}, \quad D_{2n+1} = 0, \quad (1)$$

where [3, p. 27-28]

$$(D + 1)^n - (D - 1)^n = \begin{cases} 2 & (n = 1) \\ 0 & (n < 1); \end{cases}$$

also

$$D^n = (2B + 1)^n = (2 - 2^n) B_n \quad (2)$$

and B_n is defined by

$$\frac{x}{e^x - 1} = \sum_0^{\infty} B_n \frac{x^n}{n!}.$$

The first few values of D_{2n} are

$$D_0 = 1, D_2 = -\frac{1}{3}, \quad D_4 = \frac{7}{15}, \quad D_6 = \frac{31}{21}, \quad D_8 = \frac{127}{15}.$$

If we define

$$\Delta_n = (D^2 - 1)^n = \sum_{r=0}^n (-1)^{n-r} \binom{n}{r} D_{2r} \quad (3)$$

we find that

$$\Delta_0 = 1, \quad \Delta_1 = \frac{2^2}{3}, \quad \Delta_2 = \frac{2^5}{15}, \quad \Delta_3 = \frac{2^9}{105}, \quad \Delta_4 = \frac{2^{11}}{105}.$$

(Compare [1, p. 821].)

By the second part of (2), both numerator and denominator of D_{2n} (in reduced form) are odd. Moreover by the Staudt-Clausen theorem for the Bernoulli numbers B_n , we have

$$p D_{2n} \equiv \begin{cases} -1 & (p \equiv 1 \pmod{2n}) \\ 0 & (p \equiv 1 \pmod{2n}) \end{cases},$$

where p is any odd prime. Thus the denominator of Δ_n can be determined (see below). The numerators of Δ_n for $0 \geq n \geq 4$ are all powers of 2. However this is really too good to be true; indeed we find that

$$\Delta_5 = -\frac{2^{15}}{3 \cdot 7 \cdot 11}, \quad \Delta_6 = \frac{2^{17} \cdot 191}{3 \cdot 5 \cdot 7 \cdot 11 \cdot 13}.$$

It may be of interest to determine the highest power of 2 dividing Δ_n . A result of this sort is analogous to Kummer's congruences for the Bernoulli or Euler numbers [2, Chapter 14]; however the standard results on Kummer's congruences are restricted to odd moduli.

It is convenient to first transform Δ_n . We have, using the symbolic notation,

$$\Delta_n = (D^2 - 1)^n = ((2B + 1)^2 - 1)^n = (4B^2 + 4B)^n = 4^n (B^2 + B)^n$$

and therefore

$$\Delta_n = 2^{2n} \sum_{r=0}^n \binom{n}{r} B_{n+r}. \quad (4)$$

Since $2B_{2r} \equiv 1 \pmod{2}$ it follows readily that the denominator of

$$\sum_{r=0}^n \binom{n}{r} B_{n+r}$$

(in reduced form) is odd. Consequently

$$\Delta_n \equiv 0 \pmod{2^{2n}}. \quad (5)$$

Incidentally (4) is a bit more convenient for computation than (2). For example

$$\Delta_5 = 2^{10} (5B_6 + 10B_8 + B_{10}) = -\frac{2^{15}}{3 \cdot 7 \cdot 11},$$

$$\begin{aligned} \Delta_6 &= 2^{12} (B_6 + 15B_8 + 15B_{10} + B_{12}) \\ &= 2^{12} \left(\frac{1}{42} - \frac{15}{30} + \frac{75}{66} - \frac{691}{2730} \right) = \frac{2^{17} \cdot 191}{3 \cdot 5 \cdot 7 \cdot 11 \cdot 13}. \end{aligned}$$

The congruence (5) can be improved. We make use of the explicit formula

$$B_n = \sum_{r=0}^n \frac{1}{r+1} \sum_{s=0}^r (-1)^s \binom{r}{s} s^n,$$

which yields

$$(B^2 + B)^n = 2^n \sum_{r=0}^{2n} \frac{1}{r+1} \sum_{s=0}^r (-1)^s \binom{r}{s} \left(\frac{1}{2}s(s+1)\right)^n. \quad (6)$$

However allowance must be made for the denominator $r+1$. Clearly the highest power of 2 contained in $r+1$, where $0 \leq r \leq 2n$, is at most 2^e , where e is determined by

$$2^e \leq 2n < 2^{e+1}. \quad (7)$$

Hence by (4) and (6),

$$\Delta_n \equiv 0 \pmod{2^{3n-e}}. \quad (8)$$

This result also can be improved for $n \geq 2$. Suppose first that n is even and let

$$2^k \mid n, 2^{k+1} + n \quad (k \geq 1). \quad (9)$$

Then for arbitrary odd u we have

$$u^n \equiv 1 \pmod{2^{k+2}}. \quad (10)$$

Put

$$S_j = S_j^{(r)} = \sum_{s \equiv j \pmod{4}} \binom{r}{s} \quad (0 \leq j \leq 3).$$

It is easily verified that

$$2(S_1 - S_2) = (1+i)^{r-1} + (1-i)^{r-1}.$$

Hence for r odd

$$S_1 - S_2 = \begin{cases} 0 & (r \equiv 3 \pmod{4}) \\ (-1)^{1/4(r-1)} 2^{1/2(r-1)} & (r \equiv 1 \pmod{4}) \end{cases}$$

Since $s(s+1)/2$ is odd if and only if $s \equiv 1$ or $2 \pmod{4}$, it follows from (10) that ($n > 2$)

$$S = \sum_{s=0}^r (-1)^s \binom{r}{s} \left(\frac{1}{2}s(s+1)\right)^n \equiv S_2 - S_1 \pmod{2^{k+2}},$$

and therefore

$$S \equiv 0 \pmod{2^{k+2}} \quad (r \equiv 3 \pmod{4}), \quad (11)$$

$$S \equiv 0 \pmod{(2^{k+2}, 2^{1/2(r-1)})} \quad (r \equiv 1 \pmod{4}). \quad (12)$$

Now when $r \equiv 1 \pmod{4}$, $r+1$ is divisible by 2 but not by 2^2 ; hence the term $S/(r+1)$ is integral ($\pmod{2}$) provided

$r > 1$. When $r \equiv 3 \pmod{4}$, let 2^{e_r} denote the highest power of 2 dividing $r + 1$. Then it follows from (11) that $S/(r + 1)$ is of the form $2^{k+2-e_r}A$, where A is integral $\pmod{2}$. The least favorable case arises when e_r has its maximum value e as determined by (7). We have therefore

$$\Delta_n \equiv 0 \pmod{2^{3n-e+k+2}} \quad (13)$$

provided $k + 2 > e$; when $k + 2 \leq e$ we have

$$\Delta_n \equiv 0 \pmod{2^{3n-1}} \quad (14)$$

since for example the term $r = 1$ has denominator 2.

In the next place for n odd, let

$$2^k \mid n - 1, \quad 2^{k+1} \nmid n - 1 \quad (k \geq 1);$$

Then (10) is replaced by

$$u^{n-1} \equiv 1 \pmod{2^{k+2}},$$

where u is odd. Put

$$\begin{aligned} T_j = T_j^{(r)} &= \frac{1}{2} \sum_{s \equiv j \pmod{4}} \binom{r}{s} s(s+1) \\ &= \binom{r}{2} S_{j-2}^{(r-2)} + r S_{j-1}^{(r-1)}. \end{aligned}$$

Thus

$$\begin{aligned} T_1 - T_2 &= \binom{r}{2} (S_3^{(r-2)} - S_0^{(r-2)}) + r (S_0^{(r-1)} - S_0^{(r-1)}) \\ &= \frac{i}{4} \binom{r}{2} \{(1+i)^{r-1} - (1-i)^{r-1}\} + \frac{r}{4} \{(1+i)^r + (1-i)^r\}. \end{aligned}$$

Simplifying we get (for $r \geq 3$)

$$T_1 - T_2 = \begin{cases} \frac{1}{4}r(r+1)2^{\frac{1}{2}(r-1)}i^{\frac{1}{2}(r+1)} & (r \equiv 3 \pmod{4}) \\ \frac{1}{2}r(2i)^{\frac{1}{2}(r-1)} & (r \equiv 1 \pmod{4}). \end{cases}$$

Also ($n > 3$)

$$T = \sum_{s=0}^r (-1)^s \binom{r}{s} \left(\frac{1}{2}s(s+1)\right)^n \equiv T_2 - T_1 \pmod{2^{k+2}}$$

and therefore in particular

$$T \equiv 0 \pmod{(2^{\frac{1}{2}(r-1)}, 2^{k+2})} \quad (r \equiv 3 \pmod{4}). \quad (15)$$

As in the case n even, it will suffice to take $r \equiv 3 \pmod{4}$. Let 2^{e_r} denote the highest power of 2 dividing $r + 1$, and consider $2^{-e_r} T$. For $e_r \leq 2$, we get the exponent -1 ; for $e_r \geq 3$ it follows that $\frac{1}{2}(r - 1) \geq e_r$. Hence if $k + 2 < e_r$, we get the exponent $k + 2 - e_r$, while if $k + 2 \geq e_r$ we again get -1 (at most). Consequently

$$\Delta_n \equiv 0 \pmod{2^{3n-e+k+2}} \quad (k + 2 < e_r), \quad (16)$$

$$\Delta_n \equiv 0 \pmod{2^{3n-1}} \quad (k + 2 \geq e_r). \quad (17)$$

Comparing (16) and (17) with (13) and (14) we may accordingly state the following

THEOREM. *Let $2^e \leq 2n < 2^{e+1}$ and let 2^k denote the highest power of 2 dividing n or $n - 1$ according as n is even or odd. Then Δ_n as defined by (1) and (3) satisfies (16) and (17).*

Applying the Staudt-Clausen theorem, it follows from (3) that

$$p\Delta_n \equiv - \sum_{r>0} (-1)^{n-rm} \binom{n}{rm} \pmod{p},$$

where $p = 2m + 1$ is an odd prime. It can be shown that

$$\sum_{r>0} (-1)^{n-rm} \binom{n}{rm} \equiv \begin{cases} 2 & (p - 1 \mid n) \\ \binom{m}{k} & (p - 1 \mid n + k, 0 < k \leq m) \\ 0 & \text{otherwise.} \end{cases}$$

In particular the prime factors of the denominator of Δ_n are simple and cannot exceed $2n + 1$.

In connection with formula (4) above it may be of interest to cite the formula [2, p. 189]

$$(-1)^n \sum_{r=0}^n \binom{n}{r} \frac{B_{n+r+1}}{n+r+1} + \frac{n! n!}{2(2n+1)!} = 0.$$

REFERENCES

1. G. FROBENIUS, Über die Bernoullischen Zahlen und die Eulerschen Polynome, *Sitzungsberichte der Preussischen Akademie der Wissenschaften* (1910), pp. 809-847.
 2. N. NIELSEN, *Traité élémentaire des nombres de Bernoulli*. Paris, 1923.
 3. N. E. NÖRLUND, *Vorlesungen über Differenzenrechnung*. Berlin, 1924.
- Duke University.