

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 7 (1961)  
**Heft:** 1: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** VECTOR FIELDS ON SPHERES AND ALLIED PROBLEMS  
**Autor:** Bott, Raoul

#### Bibliographie

**DOI:** <https://doi.org/10.5169/seals-37127>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

proof which possibly goes more to the heart of the matter emerges from the point of view of Atiyah and Hirzebruch. They define the groups:

$$KU^i(X) = \pi[E^{-i}X; Z \times B_U] \quad i \leq 0$$

where  $\pi[A, B]$  denotes homotopy classes of maps. In this terminology the periodicity formula:  $\Omega^2(Z \times B_U) = Z \times B_U$  is expressed by:

$$KU^i(X) = KU^{i+2}(X) \quad i \leq -2.$$

Now Atiyah and Hirzebruch use this recurrence to define  $KU^i(X)$  for all integers  $i$ , and then show that the resulting functor  $X \rightarrow \{KU^i(X)\}$  satisfies all the axioms of a cohomology theory—except for the dimension axiom. Further they observe that the uniqueness theorem of Eilenberg-Steenrod can be generalized to yield a spectral sequence relating  $E_2 = H^*(X; KU^*(p))$  to  $KU^*(X)$ . (Here  $KU^i(p)$ —the  $KU$ —theory of a point—is  $Z$  if  $i$  is even and 0 otherwise.) This sequence immediately implies the proposition. (See [8].)

To return to our bundles  $\xi^m$  on  $P_n$ . By the proposition just discussed the restriction of  $\xi^m$  to  $P_{n-k}$  will be trivial if  $m \geq n - k$ . By trivializing this element on  $P_{n-k}$  we obtain bundles  $\xi^m$  on  $P_{n,k}$  which under the projection  $\pi: P_n \rightarrow P_{n,k}$  go over into  $\xi^m$ ,  $m \geq n - k$ . In particular,  $\pi^* ch(\xi^m) = (e^x - 1)^m$ . Thus in any case we obtain these criteria the  $S$ -reducibility of  $P_{n,k}$ :

$P_{n,k}$  is  $S$ -reducible only if the coefficient of  $x^{n-1}$  in  $(e^x - 1)^m$ ,  $m \geq n - k$ , is an integer.

This is the number theoretical condition from which Atiyah and Todd deduce theorem III. Their result is the best possible one obtainable from the test-space  $B_U$ , because one can show quite easily, with the spectral sequence alluded to earlier, that the elements 1,  $\xi^m$ ;  $n - 1 \geq m \geq n - k$ ; form a base of  $KU(P_{n,k})$ .

#### BIBLIOGRAPHY

- [1] ADAMS, J., On the nonexistence of elements of Hopf invariant one.  
*Ann. of Math.*, 72 (1960), 20-104.
- [2] ATIYAH, M., *Thom Complexes* (to be published).

- [3] ATIYAH, M., and F. HIRZEBRUCH, Riemann-Roch theorems for differentiable manifolds. *Bull. Amer. Math. Soc.*, 65 (1959), 276-284.
- [4] —— and J. TODD, On complex Stiefel Manifolds. *Proc. Camb. Phil. Soc.* (1960).
- [5] BOTT, R., The stable homotopy of the classical groups. *Ann. of Math.*, 70 (1959).
- [6] —— Quelques remarques sur les théorèmes de périodicité. *Bull. Soc. Math. France* (1959), 293-310.
- [7] —— and J. MILNOR, On the parallelizability of the spheres. *Bull. Amer. Math. Soc.*, 64 (1958), 87-89.
- [8] ECKMANN, B., Gruppentheoretischer Beweis des Satzes von Hurwitz-Radon über die Komposition quadratischer Formen. *Comm. Math. Helv.*, 15 (1943), 358-366.
- [9] HIRZEBRUCH, F., A Riemann-Roch theorem for differentiable manifolds. *Séminaire Bourbaki* (1959), no 177.
- [10] HURWITZ, A., Ueber die Komposition des quadratischen Formen. *Math. Ann.*, 88 (1923), 1-25.
- [11] —— Ueber die Komposition quadratischer Formen von beliebig vielen Variabeln. *Nachr. Ges. d. Wiss. Göttingen* (1898), 309-316.
- [12] JAMES, I., The intrinsic join. *Proc. London Math. Soc.* (3), 8 (1958), 507-35.
- [13] —— Cross sections of Stiefel manifolds. *Ibid.*, 536-47.
- [14] —— Spaces associated with Stiefel Manifolds. *Ibid.* (3), 9 (1959), 115-40.
- [15] KERVAIRE, M., Nonparallelizability of the  $n$ -sphere for  $n > 7$ . *Proc. Nat. Acad. Sci. U.S.A.* (1958), 283-283.
- [16] RADON, J., Lineare Scharen orthogonaler Matrizen. *Abh. Sem. Hamburg*, I (1923), 1-14.
- [17] SPANIER, E. and J. H. C. WHITEHEAD, Duality in homotopy theory. *Mathematica*, 2 (1955), 56-80.

Department of Mathematics  
 Harvard University  
 Cambridge, Mass. U.S.A.