Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 8 (1962)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: NEW FIXED POINT THEOREM FOR CONTINUOUS MAPS OF THE CLOSED n-CELL

Autor: Abian, Alexander / Brown, Arthur B.

Kapitel: 4. Preliminary Lemmas

DOI: https://doi.org/10.5169/seals-37951

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 16.10.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

R^n, and $(n-1)$-spheres oriented with orientations induced by their interiors.

Symbols c^{n-1}, g^{n-1}, ... denote oriented $(n-1)$-cycles in R^n; D^{n-1}, V^{n-1}, ... denote $(n-1)$-spheres in R^n. E^n denotes a closed solid n-sphere in R^n, and the boundary of E^n is denoted by S^{n-1}. η^n denotes a closed n-cell in R^n and the boundary of η^n is denoted by σ^{n-1}.

In this paper η^n is assumed to be the image of E^n under homeomorphism θ, and η^n and σ^{n-1} obtain their orientations from E^n and S^{n-1} respectively.

3. The Turning Index

Let c^{n-1} be an $(n-1)$-cycle in R^n and g a continuous map of c^{n-1} into R^n having no fixed point. Let D^{n-1} be an $(n-1)$-sphere with center 0, called a *direction sphere* [2]. Let c^{n-1} be mapped on D^{n-1} as follows. To a point $c \varepsilon c^{n-1}$ there corresponds a point $d \varepsilon D^{n-1}$ such that the line segment from 0 to d has the same sense and direction as that from c to $g(c)$. The resulting $(n-1)$-cycle g^{n-1} on D^{n-1} is called, in the sequel, *the $(n-1)$-cycle g^{n-1} resulting from g applied to c^{n-1}*, and the degree of the resulting map, that is, the multiple of D^{n-1} which is homologous to g^{n-1} (which is clearly independent of the radius of D^{n-1} and the location of 0) is called the *turning index* of c^{n-1} under g.

If p is a point not on c^{n-1}, the *index of p relative to c^{n-1}* is defined as the turning index of the map which maps every point of c^{n-1} into p. (For odd n, this is the negative of the corresponding definition given in [3], as shown by Theorem 1.5, page 105).

4. Preliminary Lemmas

Lemma 1. *Let g and h be two continuous maps into R^n of an $(n-1)$-cycle c^{n-1}, such that neither leaves any point of c^{n-1} fixed, and, for no point $c \varepsilon c^{n-1}$ are the directions from c to $g(c)$ and from c to $h(c)$ exactly opposite. Then the turning indices of c^{n-1} under g and h are equal.*

Proof. For each $c \varepsilon c^{n-1}$, the directions of the two vectors $\overline{c,g(c)}$ and $\overline{c,h(c)}$ are not opposite and hence, if not identical,

determine a 2-plane P in which they make an angle of less than π radians. As a parameter p varies from 0 to 1, let the direction of $\overline{c,h}\,(c)$ change in P so that the angle between the two vectors $\overline{c,h}\,(c)$ and $\overline{c,g}\,(c)$ decreases uniformly to zero while their lengths remain fixed. If the angle is zero at the start, no change in direction takes place. For each value of p, $0 \leq p \leq 1$, the corresponding mapping as determined above in the definition of turning index, maps c^{n-1} on the direction sphere D^{n-1}, and the result, as p varies from 0 to 1, is to deform the $(n-1)$-cycle h^{n-1} on D^{n-1} resulting from h applied to c^{n-1} into the $(n-1)$-cycle g^{n-1} resulting from g applied to c^{n-1}. Hence h^{n-1} is homologous to g^{n-1}, and therefore to the same multiple of D^{n-1}, so that the turning indices under consideration are equal. Thus Lemma 1 is proved.

LEMMA 2. *Let g be a continuous map into R^n of an $(n-1)$-cycle c^{n-1},* such that c^{n-1} and $g\,(c^{n-1})$ *are contained in different half-spaces into which R^n is separated by some $(n-1)$-plane. Then the turning index of c^{n-1} under g is zero.*

Proof. Since the $(n-1)$-cycle g^{n-1} resulting from g applied to c^{n-1} is clearly entirely on one hemisphere of D^{n-1}, we conclude that c^{n-1} cannot be homologous to any multiple of D^{n-1} other than zero. Thus Lemma 2 is proved.

LEMMA 3. *Let σ^{n-1} be the boundary of a closed n-cell $\eta^n \subset R^n$. Let e be a point in the inside of σ^{n-1}. Then the index of e relative to σ^{n-1} is 1 or -1.*

While this result is given in [3], page 109, Theorem 4.1, the following proof is given as shorter and obtained independently.

Proof. Let η^n and σ^{n-1} be respectively the homeomorphic images (under homeomorphism θ) of the closed solid n-sphere E^n with boundary S^{n-1}, i.e., $\eta^n = \theta\,(E^n)$ and $\sigma^{n-1} = \theta\,(S^{n-1})$. By use of the invariance of regionality, it is easy to show that $\eta^n = \theta\,(E^n)$ contains no point outside σ^{n-1} and contains every point inside σ^{n-1}.

Let V^{n-1} be an $(n-1)$-sphere with center at e, so small that V^{n-1} and its interior are inside σ^{n-1}, hence composed of points of η^n. Let $\beta^{n-1} = \theta^{-1}\,(V^{n-1})$ and $d = \theta^{-1}\,(e)$.

For each point $b \varepsilon \beta^{n-1}$ let the half-line beginning at d and passing through b intersect S^{n-1} at b'.

Now, for every t, with $0 \leq t \leq 1$, let $\beta^{n-1}(t)$ be the $(n-1)$-cycle determined as follows. For each point $b \varepsilon \beta^{n-1}$ there corresponds a point $b(t)$ of $\beta^{n-1}(t)$ on the closed segment from b to b' such that the distance from b to $b(t)$ is t times the distance from b to b'.

Let $V^{n-1}(t) = \theta[\beta^{n-1}(t)], \qquad 0 \leq t \leq 1.$

As t varies from 0 to 1, the cycle $V^{n-1}(t)$ undergoes a deformation from initial position $V^{n-1}(0) = V^{n-1}$ to final position $V^{n-1}(1)$. Since $V^{n-1}(1)$ is on σ^{n-1}, there is an integer x such that

$$(1) \qquad V^{n-1}(1) \sim x\,\sigma^{n-1} \qquad \text{on } \sigma^{n-1},$$

where $\sim$ stands for " is homologous to ".

For each t, let $k(t)$ be the mapping which maps every point of $V^{n-1}(t)$ into e, and let V^{n-1} serve as the direction sphere. As t varies from 0 to 1, the $(n-1)$-cycle $k^{n-1}(0)$ resulting from $k(0)$ applied to V^{n-1} is deformed on the direction sphere V^{n-1} into the $(n-1)$-cycle $k^{n-1}(1)$ resulting from $k(1)$ applied to $V^{n-1}(1)$. Thus these two $(n-1)$ cycles are homologous on V^{n-1}. Therefore the index of e relative to V^{n-1} equals the index of e relative to $V^{n-1}(1)$. However, since $k(0)$ maps every point of V^{n-1} into e, we derive that ([4], page 92)

$$(2) \qquad \text{the index of } e \text{ relative to } V^{n-1}(1) = (-1)^n.$$

Let y be the index of e relative to σ^{n-1}. From (1) we infer that xy is the index of e relative to $V^{n-1}(1)$. Hence, by (2), $xy = (-1)^n$. Consequently, $y = 1$ or $y = -1$. Thus Lemma 3 is proved.

LEMMA 4. *If a continuous map f of a closed n-cell $\eta^n \subset R^n$ into R^n has no fixed point, then the turning index of the boundary σ^{n-1} of η^n under f is zero.*

Proof. Let, as in the proof of Lemma 3, $\eta^n = \theta(E^n)$ and $\sigma^{n-1} = \theta(S^{n-1})$ be respectively the images under the homeomorphism θ of the closed solid n-sphere E^n and its boundary S^{n-1}.

Let u be the center of S^{n-1}. Since f has no fixed point, it is clear that we can choose $d > 0$ so small that a closed solid n-sphere H_d^n of radius d with center at $\theta(u)$ is entirely in η^n, and H_d^n and its image $f(H_d^n)$ are contained in different half-spaces into which R^n is separated by some $(n-1)$-plane.

Now, let S^{n-1} undergo a deformation by uniform radial shrinking toward u till it reaches a position S_2^{n-1} whose image σ_2^{n-1} under θ is contained in the interior of H_d^n. By means of θ, there results a deformation of σ^{n-1} into σ_2^{n-1} which by means of the mapping f induces a deformation, on the direction sphere, of the $(n-1)$-cycle f^{n-1} resulting from f applied to σ^{n-1} into the $(n-1)$-cycle f_2^{n-1} resulting from f applied to σ_2^{n-1}.

Thus the turning index of σ^{n-1} under f equals the turning index of σ_2^{n-1} under f, which by Lemma 2 equals zero. Thus Lemma 4 is proved.

5. The Theorems

Theorem 1. *Let $\eta^n \subset R^n$ be a closed n-cell and f a continuous mapping of η^n into R^n such that f maps the boundary σ^{n-1} of η^n into η^n. Then f has at least one fixed point.*

Proof. Assume no fixed points. Let, as in the case of Lemma 3, η^n and σ^{n-1} be respectively the images (under the homeomorphism θ) of the closed solid n-sphere E^n with boundary S^{n-1}, i.e., $\eta^n = \theta(E^n)$ and $\sigma^{n-1} = \theta(S^{n-1})$.

Let u be the center of S^{n-1}. Consider the mapping f' of σ^{n-1} which maps every point $\sigma \varepsilon \sigma^{n-1}$ into the point $\theta(u)$. Since f' is the mapping which appears in the definition of the index of $\theta(u)$ relative to σ^{n-1}, we see by Lemma 3 that the turning index of σ^{n-1} under f' is non-zero.

By hypothesis, $f(\sigma) \varepsilon \eta^n$ for every $\sigma \varepsilon \sigma^{n-1}$. Hence we may deform $f(\sigma^{n-1})$ as follows. As a parameter p varies from 0 to 1, the point σ' moves in η^n along the path $\theta[\overline{\theta^{-1} f(\sigma), u}]$ starting from σ and ending at $\theta(u)$.

For $p = 1$, the above deformation yields the mapping f'. Therefore, the $(n\text{-}1)$-cycle resulting from f applied to σ^{n-1} is homologous on the direction sphere (as a consequence of a deformation) to the $(n-1)$-cycle resulting from f' applied to