

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	8 (1962)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	ON THE CONSTRUCTION OF RELATED EQUATIONS FOR THE ASYMPTOTIC THEORY OF LINEAR ORDINARY DIFFERENTIAL EQUATIONS ABOUT A TURNING POINT
Autor:	Langer, Rudolph E.
Kapitel:	4. TWO DIFFERENTIAL OPERATORS OF THE ORDERS p AND q.
DOI:	https://doi.org/10.5169/seals-37963

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

the factors $B(\chi)$ and $\Gamma(\chi)$ also in an alternative form. This is done as follows:

Let K designate the familiar matrix

$$K(z) = \begin{bmatrix} 0 & 0 & - & - & - & - & - & -b_p \\ 1 & 0 & - & - & - & - & - & -b_{p-1} \\ 0 & 1 & 0 & - & - & - & - & -b_{p-2} \\ - & - & - & - & - & - & - & - \\ - & - & - & - & - & - & - & - \\ 0 & - & - & - & - & 1 & -b_1 & \end{bmatrix} \quad (3.3)$$

The eigen-values of this are the roots x_i , $i = 1, 2, \dots, p$ of the equation $B(x) = 0$. If we designate by ξ_i an eigen-vector corresponding to x_i we have

$$K^h \xi_i = x_i^h \xi_i, \quad h = 1, 2, \dots, q,$$

and accordingly

$$\Gamma(K) \xi_i = \Gamma(x_i) \xi_i.$$

Thus $\Gamma(x_i)$ is an eigenvalue of the matrix $\Gamma(K)$, and, since the product of the eigenvalues is the determinant of the matrix, we have

$$|\Gamma(K)| = \prod_{i=1}^p \Gamma(x_i).$$

Observing that no factor on the right is zero, and giving to $\Gamma(K)$ its explicit form, we conclude with the result

$$|\sum_{j=0}^q c_j(z) K^{q-j}(z)| \neq 0 \quad (3.4)$$

4. TWO DIFFERENTIAL OPERATORS OF THE ORDERS p AND q .

Let the functions $\beta_j(z, \lambda)$ and $\gamma_i(z, \lambda)$ be taken to be polynomials of the degree $(r - 1)$ in $1/\lambda$, thus

$$\begin{aligned} \beta_j(z, \lambda) &= \sum_{v=0}^{r-1} \frac{\beta_{j,v}(z)}{\lambda^v}, \quad \beta_{j,0}(z) \equiv b_j(z); \quad j = 1, 2, \dots, p, \\ \gamma_i(z, \lambda) &= \sum_{v=0}^{r-1} \frac{\gamma_{i,v}(z)}{\lambda^v}, \quad \gamma_{i,0}(z) \equiv c_i(z); \quad i = 1, 2, \dots, q. \end{aligned} \quad (4.1)$$

As has been indicated, the terms of the zeroth degree are to be the coefficients which appear in the formulas (2. 7). The remaining terms, $\beta_{j,v}(z)$ and $\gamma_{i,v}(z)$, with $v \geq 1$, shall be analytic over the z -region, but beyond that shall be left, for the moment, unspecified. By l and m we shall designate the differential operators

$$\begin{aligned} l &= \sum_{j=0}^p \lambda^j \beta_j(z, \lambda) D^{p-j}, \quad \beta_0 \equiv 1, \\ m &= \sum_{i=0}^q \lambda^i \gamma_i(z, \lambda) D^{q-i}, \quad \gamma_0 \equiv 1. \end{aligned} \tag{4. 2}$$

The immediate objective will be to show that the unspecified terms in the formulas (4. 1) can be so chosen as to give the differential form $l(m(u))$ coefficients which differ from those of the form (2. 2) only by terms that are of at least the r^{th} degree in $1/\lambda$.

The k -fold differentiation of $m(y)$ yields the formula

$$D^k m(y) = \sum_{i=0}^q \sum_{s=0}^k \lambda^i \binom{k}{s} D^s \gamma_i D^{q-i+k-s} y,$$

in which the symbol $\binom{k}{s}$ denotes, as customarily, the coefficient of x^s in the binomial expansion of $(1+x)^k$. On using $i+s$ in place of i as the variable of summation, and observing that the terms which appear to have been gratuitously included are ones to which the value zero are to be assigned, we find that the formula may be written

$$D^k m(y) = \sum_{i=0}^{q+k} \sum_{s=0}^p \lambda^{i-s} \binom{k}{s} D^s \gamma_{i-s} D^{q+k-i} y, \quad k = 0, 1, 2, \dots, p. \tag{4. 3}$$

From this it follows that

$$l(m(y)) = \sum_{j=0}^p \sum_{i=0}^q \sum_{s=0}^{p-j} \lambda^{j+i-s} \binom{p-j}{s} \beta_j D^s \gamma_{i-s} D^{n-j-i} y.$$

This formula is again improved by using $i+j$ in place of i as the variable of summation. It becomes, then

$$l(m(y)) = \sum_{i=0}^n \lambda^i \Psi_i(z, \lambda) D^{n-i} y, \quad (4.4)$$

with

$$\Psi_i(z, \lambda) = \sum_{j=0}^p \sum_{s=0}^{p-j} \lambda^{-s} \binom{p-j}{s} \beta_j D^s \gamma_{i-j-s}. \quad (4.5)$$

The functions $\Psi_i(z, \lambda)$, inasmuch as they are combinations of those given in (4.1), are polynomials in $1/\lambda$. We may therefore write them in the form

$$\Psi_i(z, \lambda) = \sum_{\mu=0}^{r-1} \frac{\psi_{i,\mu}(z)}{\lambda^\mu} + \frac{\psi_{i,r}(z, \lambda)}{\lambda^r}. \quad (4.6)$$

A comparison of the terms in like powers of $1/\lambda$ in the relations (4.5) and (4.6) yields formulas for the functions $\psi_{i,\mu}(z)$. Those for which $\mu = 0$ are particularly easy to obtain. On setting $s = 0$ in (4.5), and replacing β_j and γ_{i-j} by their leading terms b_j and c_{i-j} , we find that

$$\psi_{i,0}(z) = \sum_{j=0}^p b_j(z) c_{i-j}(z).$$

Recourse to the relation (2.8) thus shows that

$$\psi_{i,0}(z) = p_{i,0}(z), \quad i = 1, 2, \dots, n. \quad (4.7)$$

At least to the extent of the leading terms of their coefficients, the forms (2.2) and (4.4) are, therefore, the same.

5. A DETERMINATION OF UNSPECIFIED COEFFICIENTS.

We propose now to deduce a formula for the general coefficient $\psi_{i,\mu}(z)$ in (4.6) by selecting the multiplier of the appropriate power of $1/\lambda$ from the formula (4.5). To begin with, it follows from the relations (4.1) that

$$\beta_j D^s \gamma_{i-j-s} = \sum_{\mu=0}^{2r-2} \sum_{k=0}^{\mu} \lambda^{-\mu} \beta_{j,k} D^s \gamma_{i-j-s, \mu-k}$$