

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 8 (1962)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE CONSTRUCTION OF RELATED EQUATIONS FOR THE ASYMPTOTIC THEORY OF LINEAR ORDINARY DIFFERENTIAL EQUATIONS ABOUT A TURNING POINT

Autor: Langer, Rudolph E.
Kapitel: 6. A SET OF FUNCTIONS $\eta_i(z, \lambda)$.
DOI: <https://doi.org/10.5169/seals-37963>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

6. A SET OF FUNCTIONS $\eta_i(z, \lambda)$.

It is the immediate purpose of this paper to show that under the hypotheses that have been made the construction of a related equation for the differential equation (2. 1) is referable to such constructions for equations of lower order. These latter will be precisely the differential equations (5. 6). In proceeding, we shall therefore suppose that the related equations

$$l^*(v) = 0, \tag{6. 1}$$

with

$$l^*(v) \equiv \sum_{j=0}^p \lambda^j \bar{\beta}_j(z, \lambda) D^{p-j} v, \tag{6. 2}$$

and

$$m^*(y) = 0, \tag{6. 3}$$

with

$$m^*(y) \equiv \sum_{i=0}^q \lambda^i \bar{\gamma}_i(z, \lambda) D^{q-i} y \tag{6. 4}$$

have been brought to hand. Their coefficients are of the forms

$$\begin{aligned} \bar{\beta}_j(z, \lambda) &= \beta_j(z, \lambda) + \frac{1}{\lambda^r} \bar{\beta}_{j,r}(z, \lambda), \\ \bar{\gamma}_j(z, \lambda) &= \gamma_j(z, \lambda) + \frac{1}{\lambda^r} \bar{\gamma}_{j,r}(z, \lambda), \end{aligned} \tag{6. 5}$$

the functions $\bar{\beta}_{i,r}$ and $\bar{\gamma}_{j,r}$ being bounded over the domains of z and λ . Since terms of a degree greater than the $(r-1)^{th}$ in $1/\lambda$ were never involved in the derivation of the relation (5. 5), it is clear that also

$$L(u) = l^*(m^*(u)) + \frac{1}{\lambda^r} \sum_{j=1}^n \lambda^j \epsilon_j(z, \lambda) D^{n-j} u, \tag{6. 6}$$

with certain bounded coefficients ϵ_j .

As for all related equations, by definition, the solutions of the equations (6. 1) and (6. 3) are known. We shall denote complete

sets for these solutions respectively by $v_i(z, \lambda)$, $i = 1, 2, \dots, p$; and $y_j(z, \lambda)$, $j = 1, 2, \dots, q$. Our method, as will be seen, requires a modification of one of these solution sets. That is the purpose of the replacement of the functions $v_i(z, \lambda)$ by corresponding ones $\eta_i(z, \lambda)$ to which we proceed.

Let the functions $\alpha_j^{(0)}(z, \lambda)$ be polynomials of the degree $(r - 1)$ in $1/\lambda$, namely

$$\alpha_j^{(0)}(z, \lambda) = \sum_{v=0}^{r-1} \frac{\alpha_{j,v}^{(0)}(z)}{\lambda^v}, \tag{6.7}$$

with coefficients $\alpha_{j,v}^{(0)}(z)$ that are analytic but, for the time being, unspecified. Then consider the formulas

$$\eta_i(z, \lambda) = \sum_{j=1}^p \lambda^{1-j} \alpha_j^{(0)}(z, \lambda) D^{j-1} v_i(z, \lambda), \quad i = 1, 2, \dots, p. \tag{6.8}$$

The differentiation of this, and the subsequent elimination of $D^p v_i$ by use of the equation (6.1), yields a corresponding formula for $D\eta_i$, and repetitions of the procedure yield more generally that

$$D^k \eta_i = \lambda^k \sum_{j=1}^p \lambda^{i-j} \alpha_j^{(k)}(z, \lambda) D^{j-1} v_i(z, \lambda), \quad k = 0, 1, 2, \dots, q. \tag{6.9}$$

In this the coefficients are recursively given by the relation

$$\alpha_j^{(k)} = \alpha_{j-1}^{(k-1)} - \bar{\beta}_{p+1-j} \alpha_p^{(k-1)} + \frac{1}{\lambda} D \alpha_j^{(k-1)} \tag{6.10}$$

The relations (6.9) yield the formula

$$m^*(\eta_i) = \lambda^q \sum_{j=1}^p \lambda^{1-j} \sigma_j^{(0)}(z, \lambda) D^{j-1} v_i. \tag{6.11}$$

with

$$\sigma_j^{(0)}(z, \lambda) = \sum_{k=0}^q \gamma_k \alpha_j^{(q-k)}. \tag{6.12}$$

These latter coefficients $\sigma_j^{(0)}(z, \lambda)$ are evidently expressible in powers of $1/\lambda$, and are therefore of the form

$$\sigma_j^{(0)}(z, \lambda) = \sum_{v=0}^{r-1} \frac{\sigma_{j,v}^{(0)}(z)}{\lambda^v} + \frac{\sigma_{j,r}^{(0)}(z, \lambda)}{\lambda^r} \tag{6.13}$$

with each $\sigma_{j,v}^{(0)}(z)$ analytic, and $\sigma_{j,r}^{(0)}(z, \lambda)$ bounded. We shall show that the elements $\alpha_{j,v}^{(0)}(z)$ in (6.7) may be so specified as to yield

$$\sigma_{j,v}^{(0)}(z) \equiv \begin{cases} 1 & \text{when } (j, v) = (1, 0), \quad v = 0, 1, 2, \dots, (r-1) \dots \\ 0 & \text{when } (j, v) \neq (1, 0). \end{cases} \quad (6.14)$$

The effect of this will be to give the formula (6.11) the form

$$m^*(\eta_i) = \lambda^q \left\{ v_i(z, \lambda) + \frac{1}{\lambda^r} \sum_{j=1}^p \lambda^{1-j} \sigma_{j,r}^{(0)}(z, \lambda) D^{j-1} v_i \right\}. \quad (6.15)$$

7. ANOTHER DETERMINATION OF COEFFICIENTS.

The dependence of the functions (6.12) upon the unspecified ones $\alpha_{j,v}^{(0)}(z)$ of (6.7) is advantageously set forth in terms of vector-matrix notation. To this end, let a column vector with the components $\varphi_i, i = 1, 2, \dots, p$, be denoted by (φ) and let the vector whose components are the terms in $1/\lambda^v$ of (φ) , namely with the components $\varphi_{i,v} i = 1, 2, \dots, p$, be denoted by $(\varphi)_v$. Also let H designate the square matrix

$$H = \begin{bmatrix} \lambda^{-1}D & 0 & 0 & - & - & -\bar{\beta}_p \\ 1 & \lambda^{-1}D & 0 & - & - & -\bar{\beta}_{p-1} \\ 0 & 1 & \lambda^{-1}D & - & - & -\bar{\beta}_{p-2} \\ - & - & - & - & - & - \\ - & - & - & - & - & - \\ 0 & 0 & - & - & - & -\bar{\beta}_1 + \lambda^{-1}D \end{bmatrix} \quad (7.1)$$

the elements of which are in part functions of z and λ , and in part the indicated differential operator. Again let H_v designate the matrix that is obtainable from (7.1) by replacing its elements by their terms in $1/\lambda^v$. The relations (6.10) are then seen at once to take the form

$$(\alpha^{(k)}) = H(\alpha^{(k-1)}).$$

With iteration defined in the manner

$$H^{[k]}(\varphi) = H(H^{[k-1]}(\varphi)), \quad H^{[1]} = H, \quad (7.2)$$