

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	8 (1962)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	ON THE CONSTRUCTION OF RELATED EQUATIONS FOR THE ASYMPTOTIC THEORY OF LINEAR ORDINARY DIFFERENTIAL EQUATIONS ABOUT A TURNING POINT
Autor:	Langer, Rudolph E.
Kapitel:	7. ANOTHER DETERMINATION OF COEFFICIENTS.
DOI:	https://doi.org/10.5169/seals-37963

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

with each $\sigma_{j,v}^{(0)}(z)$ analytic, and $\sigma_{j,r}^{(0)}(z, \lambda)$ bounded. We shall show that the elements $\alpha_{j,v}^{(0)}(z)$ in (6.7) may be so specified as to yield

$$\sigma_{j,v}^{(0)}(z) \equiv \begin{cases} 1 & \text{when } (j, v) = (1, 0), v = 0, 1, 2, \dots, (r-1) \\ 0 & \text{when } (j, v) \neq (1, 0). \end{cases} \quad (6.14)$$

The effect of this will be to give the formula (6.11) the form

$$m^*(\eta_i) = \lambda^q \left\{ v_i(z, \lambda) + \frac{1}{\lambda^r} \sum_{j=1}^p \lambda^{1-j} \sigma_{j,r}^{(0)}(z, \lambda) D^{j-1} v_i \right\}. \quad (6.15)$$

7. ANOTHER DETERMINATION OF COEFFICIENTS.

The dependence of the functions (6.12) upon the unspecified ones $\alpha_{j,v}^{(0)}(z)$ of (6.7) is advantageously set forth in terms of vector-matrix notation. To this end, let a column vector with the components $\varphi_i, i = 1, 2, \dots, p$, be denoted by (φ) and let the vector whose components are the terms in $1/\lambda^v$ of (φ) , namely with the components $\varphi_{i,v}, i = 1, 2, \dots, p$, be denoted by $(\varphi)_v$. Also let H designate the square matrix

$$H = \begin{bmatrix} \lambda^{-1}D & 0 & 0 & - & - & - & -\bar{\beta}_p \\ 1 & \lambda^{-1}D & 0 & - & - & - & -\bar{\beta}_{p-1} \\ 0 & 1 & \lambda^{-1}D & - & - & - & -\bar{\beta}_{p-2} \\ - & - & - & - & - & - & - \\ - & - & - & - & - & - & - \\ 0 & 0 & - & - & - & - & -\bar{\beta}_1 + \lambda^{-1}D \end{bmatrix} \quad (7.1)$$

the elements of which are in part functions of z and λ , and in part the indicated differential operator. Again let H_v designate the matrix that is obtainable from (7.1) by replacing its elements by their terms in $1/\lambda^v$. The relations (6.10) are then seen at once to take the form

$$(\alpha^{(k)}) = H(\alpha^{(k-1)}).$$

With iteration defined in the manner

$$H^{[k]}(\varphi) = H(H^{[k-1]}(\varphi)), \quad H^{[1]} = H, \quad (7.2)$$

and with $H^{[0]}$ signifying the unit matrix, it is then easily seen that

$$(\alpha^{(k)}) = H^{[k]}(\alpha^{(0)}). \quad (7.3)$$

The relation (6.12) may thus be written in the form

$$(\sigma^{(0)}) = J(\alpha^{(0)}), \quad (7.4)$$

with J standing for the matrix

$$J = \sum_{k=0}^q \bar{\gamma}_k H^{[q-k]}. \quad (7.5)$$

The evaluations

$$(\sigma^{(0)})_v = \sum_{j=0}^v J_j (\alpha^{(0)})_{v-j},$$

$$J_j = \sum_{k=0}^q \sum_{i=0}^j \bar{\gamma}_{k,i} H^{[q-k]}_{j-i},$$

evidently combine to yield the formula

$$(\sigma^{(0)})_v = \sum_{k=0}^q \sum_{j=0}^v \sum_{i=0}^j \bar{\gamma}_{k,i} H^{[q-k]}_{j-i} (\alpha^{(0)})_{v-j}. \quad (7.6)$$

In connection with this, certain observations are apropos. To begin with, the index value $j = 0$ implies $i = 0$, whereas by (6.5) and (4.1), $\bar{\gamma}_{k,0} = c_k(z)$. Further when $i = j$ the matrix $H^{[q-k]}_{j-i}$ reduces to precisely $K^{q-k}(z)$, with $K(z)$ as given in (3.3). On the basis of these facts the equation (7.6) may be arranged into the form

$$\sum_{k=0}^q c_k(z) K^{q-k}(z) (\alpha^{(0)})_v = (\sigma^{(0)})_v - \sum_{k=0}^q \sum_{j=1}^v \sum_{i=0}^j \bar{\gamma}_{k,i} H^{[q-k]}_{j-i} (\alpha^{(0)})_{v-j} \quad (7.7)$$

This is a vector equation for $(\alpha^{(0)})_v$, which we shall consider for successive values of v , assuming that the values (6.14) have been assigned.

When $v = 0$, the triple sum on the right of the equality in (7.7) vanishes, and the right-hand member is, therefore, the vector $(\sigma^{(0)})_0$ whose first component is 1 and whose other components are 0. The equation is therefore a non-homogeneous

one, and accordingly admits of an analytic solution for $(a^{(0)})_0$ provided the matrix multiplier of this vector on the left is non-singular. This condition is assured by the relation (3. 4).

Now we may proceed by induction. Assuming that the vectors $(a^{(0)})_j$ for $j = 1, 2, \dots, (v-1)$, have been determined and are analytic, the right-hand member of the equation (7. 7) is known. As in the case $v = 0$, so now, the equation is analytically solvable. The solutions for the successive values $v = 0, 1, 2, \dots, (r-1)$, yield the coefficients (6. 7) for which the functions $\eta_i(z, \lambda)$, as given by the formulas (6. 8), fulfill the relations (6. 5).

8. ON LINEAR INDEPENDENCE.

With the functions $a_j^{(0)}(z, \lambda)$ now at hand, we have at our disposal the n known functions $y_j(z, \lambda)$, $j = 1, 2, \dots, q$, which are the solutions of the differential equation (6. 3), and $\eta_i(z, \lambda)$, $i = 1, 2, \dots, p$, which are given by the formulas (6. 8). We shall show that these functions are linearly independent.

Let the Wronskians of the entire set and of the respective sub-sets be denoted respectively by W_n , $W_q(y)$ and $W_p(\eta)$. If the usual form

$$W_n = \begin{vmatrix} y_1 & - & - & - & y_q & \eta_1 & - & - & - & \eta_p \\ Dy_1 & - & - & - & Dy_q & D\eta_1 & - & - & - & D\eta_q \\ - & - & - & - & - & - & - & - & - & - \\ - & - & - & - & - & - & - & - & - & - \\ D^{n-1}y_1 & - & - & - & D^{n-1}y_q & D^{n-1}\eta_1 & - & - & - & D^{n-1}\eta_p \end{vmatrix} \quad (8. 1)$$

is modified by adding to each of the last p rows suitable multiples of the preceding ones, the formula can be made to appear thus

$$\begin{vmatrix} y_1 & - & - & - & - & y_q & \eta_1 & - & - & - & \eta_p \\ Dy_1 & - & - & - & - & Dy_q & D\eta_1 & - & - & - & D\eta_p \\ - & - & - & - & - & - & - & - & - & - & - \\ D^{q-1}y_1 & - & - & - & - & D^{q-1}y_q & D^{q-1}\eta_1 & - & - & - & D^{q-1}\eta_p \\ m^*(y_1) & - & - & - & - & m^*(y_q) & m^*(\eta_1) & - & - & - & m^*(\eta_p) \\ Dm^*(y_1) & - & - & - & - & - & - & - & - & - & - \\ - & - & - & - & - & - & - & - & - & - & - \\ D^{p-1}m^*(y_1) & - & - & - & - & D^{p-1}m^*(y_q) & D^{p-1}m^*(\eta_1) & - & - & - & D^{p-1}m^*(\eta_p) \end{vmatrix} \quad (8. 2)$$