

<b>Zeitschrift:</b>	L'Enseignement Mathématique
<b>Herausgeber:</b>	Commission Internationale de l'Enseignement Mathématique
<b>Band:</b>	8 (1962)
<b>Heft:</b>	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
<b>Artikel:</b>	ON THE CONSTRUCTION OF RELATED EQUATIONS FOR THE ASYMPTOTIC THEORY OF LINEAR ORDINARY DIFFERENTIAL EQUATIONS ABOUT A TURNING POINT
<b>Autor:</b>	Langer, Rudolph E.
<b>Kapitel:</b>	8. On linear independence.
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-37963">https://doi.org/10.5169/seals-37963</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

one, and accordingly admits of an analytic solution for  $(a^{(0)})_0$  provided the matrix multiplier of this vector on the left is non-singular. This condition is assured by the relation (3. 4).

Now we may proceed by induction. Assuming that the vectors  $(a^{(0)})_j$  for  $j = 1, 2, \dots, (v-1)$ , have been determined and are analytic, the right-hand member of the equation (7. 7) is known. As in the case  $v = 0$ , so now, the equation is analytically solvable. The solutions for the successive values  $v = 0, 1, 2, \dots, (r-1)$ , yield the coefficients (6. 7) for which the functions  $\eta_i(z, \lambda)$ , as given by the formulas (6. 8), fulfill the relations (6. 5).

### 8. ON LINEAR INDEPENDENCE.

With the functions  $a_j^{(0)}(z, \lambda)$  now at hand, we have at our disposal the  $n$  known functions  $y_j(z, \lambda)$ ,  $j = 1, 2, \dots, q$ , which are the solutions of the differential equation (6. 3), and  $\eta_i(z, \lambda)$ ,  $i = 1, 2, \dots, p$ , which are given by the formulas (6. 8). We shall show that these functions are linearly independent.

Let the Wronskians of the entire set and of the respective sub-sets be denoted respectively by  $W_n$ ,  $W_q(y)$  and  $W_p(\eta)$ . If the usual form

$$W_n = \begin{bmatrix} y_1 & - & - & - & y_q & \eta_1 & - & - & - & \eta_p \\ Dy_1 & - & - & - & Dy_q & D\eta_1 & - & - & - & D\eta_q \\ - & - & - & - & - & - & - & - & - & - \\ - & - & - & - & - & - & - & - & - & - \\ D^{n-1}y_1 & - & - & - & D^{n-1}y_q & D^{n-1}\eta_1 & - & - & - & D^{n-1}\eta_p \end{bmatrix} \quad (8. 1)$$

is modified by adding to each of the last  $p$  rows suitable multiples of the preceding ones, the formula can be made to appear thus

$$= \begin{bmatrix} y_1 & - & - & - & - & y_q & \eta_1 & - & - & - & \eta_p \\ Dy_1 & - & - & - & - & Dy_q & D\eta_1 & - & - & - & D\eta_p \\ - & - & - & - & - & - & - & - & - & - & - \\ D^{q-1}y_1 & - & - & - & - & D^{q-1}y_q & D^{q-1}\eta_1 & - & - & - & D^{q-1}\eta_p \\ m^*(y_1) & - & - & - & - & m^*(y_q) & m^*(\eta_1) & - & - & - & m^*(\eta_p) \\ Dm^*(y_1) & - & - & - & - & - & - & - & - & - & - \\ - & - & - & - & - & - & - & - & - & - & - \\ D^{p-1}m^*(y_1) & - & - & - & - & D^{p-1}m^*(y_q) & D^{p-1}m^*(\eta_1) & - & - & - & D^{p-1}m^*(\eta_p) \end{bmatrix} \quad (8. 2)$$

In this, however, each of the elements occupying a position in one of the first  $q$  columns and in one of the last  $p$  rows is zero. The formula therefore reduces at once to

$$W_n = W_q(y) T, \quad (8.3)$$

with

$$T = \begin{bmatrix} m^*(\eta_1) & - & - & - & - & m^*(\eta_p) \\ Dm^*(\eta_1) & - & - & - & - & Dm^*(\eta_p) \\ - & - & - & - & - & - \\ - & - & - & - & - & - \\ D^{p-1}m^*(\eta_1) & - & - & - & - & D^{p-1}m^*(\eta_p) \end{bmatrix}. \quad (8.4)$$

Now  $m^*(\eta_j)$  is given by the formula (6.15). If this is repeatedly differentiated, and at each step the element  $D^p v_j$  is eliminated by use of the equation (6.1), the results are the formulas

$$D^i m^*(\eta_j) = \lambda^q D^i v_j + \lambda^{q+i-r} \sum_{\mu=0}^p \lambda^{1-\mu} \sigma_{\mu, r}^{(i)} D^{\mu-1} v_j, \quad i = 0, 1, 2, \dots . \quad (8.5)$$

We may write this also, with the use of the symbol  $\delta_{i,j}$  to denote 1 when  $j = i$  and 0 when  $j \neq i$ , in the form

$$D^{i-1} m^*(\eta_j) = \lambda^{q+i-1} \sum_{\mu=1}^p \left\{ \delta_{i,\mu} + \frac{\sigma_{\mu, r}^{(i-1)}}{\lambda^r} \right\} \frac{D^{\mu-1} v_j}{\lambda^{\mu-1}}. \quad (8.6)$$

This shows, now, at once, that the determinant  $T$  can be factored, thus

$$T = \lambda^{pq} E W_p(v) \quad (8.7)$$

in which  $E$  is the determinant whose element in the  $i^{\text{th}}$  row and  $j^{\text{th}}$  column is indicated thus

$$E = \left| \delta_{i,j} + \frac{\sigma_{j,r}^{(i-1)}}{\lambda^r} \right|. \quad (8.8)$$

It is clear that  $E$  differs from 1 by terms of at least the degree  $r$  in  $1/\lambda$ . Since  $W_p(v)$  and  $W_q(y)$  are non-vanishing, it follows from (8.3) and (8.7) that the same is true of  $W_n$ .

## 9. THE RELATED EQUATION.

We are prepared now to make the construction toward which this entire discussion has been directed.

Consider the equation

$$L^*(u) = 0 . \quad (9.1)$$

with

$$L^*(u) = \frac{1}{T} \begin{bmatrix} m^*(\eta_1) & - & - & - & - & - & m^*(\eta_p) & m^*(u) \\ Dm^*(\eta_1) & - & - & - & - & - & Dm^*(\eta_p) & Dm^*(u) \\ - & - & - & - & - & - & - & - \\ - & - & - & - & - & - & - & - \\ D^{p-1}m^*(\eta_1) & - & - & - & - & - & D^{p-1}m^*(\eta_p) & D^{p-1}m^*(u) \\ l^*(m^*(\eta_1)) & - & - & - & - & - & l^*(m^*(\eta_p)) & l^*(m^*(u)) \end{bmatrix} . \quad (9.2)$$

$T$  being the determinant given in (8.4). This is clearly a differential equation of the  $n^{\text{th}}$  order in  $u$ , for which each one of the functions  $y_j(z, \lambda)$  and  $\eta_i(z, \lambda)$  is a solution. For if  $\eta_i$  is substituted for  $u$  two of the columns of the determinant (9.2) are the same, and if  $u$  is replaced  $y_j$  every element of the last column vanishes. Because the  $n$  solutions thus produced are linearly independent the solutions of the equation (9.1) are completely known.

The co-factor of the element  $l^*(m(u))$  in the formula (9.2) is the determinant  $T$ . The expansion of the formula thus gives it the aspect

$$L^*(u) = l^*(m^*(u)) - \sum_{v=1}^p \frac{T_v}{T} D^{p-v} m^*(u) , \quad (9.3)$$

where  $T_v$  is the determinant that is obtainable from the formula (8.4) by replacing its elements  $D^{p-v} m^*(\eta_j)$  by  $l^*(m^*(\eta_j))$ .

From the formula (8.5) it is seen that

$$l^*(m^*(\eta_j)) = \lambda^n \sum_{v=1}^p \frac{\tau_v(z, \lambda)}{\lambda^r} \cdot \frac{D^{u-1} v_j}{\lambda^{u-1}} \quad (9.4)$$

with

$$\tau_v(z, \lambda) = \sum_{k=0}^p \bar{\beta}_k(z, \lambda) \sigma_{v, r}^{(p-k)}(z, \lambda) . \quad (9.5)$$

The replacements which change  $T$  to  $T_v$  are thus seen to be ones which replace

$$\lambda^{n-v} \left\{ \delta_{p-v, j} + \frac{\sigma_{j, r}^{(p-v)}}{\lambda^r} \right\} \text{ by } \lambda^n \frac{\tau_v}{\lambda^r}.$$

It follows that

$$\frac{T_v}{T} = \lambda^v \frac{\theta_v(z, \lambda)}{\lambda^r},$$

with some function  $\theta_v(z, \lambda)$  which is bounded over the  $z$  and  $\lambda$  domains. This gives to the relation (9.3) the form

$$L^*(u) = l^*(m^*(u)) - \frac{1}{\lambda^r} \sum_{v=1}^p \lambda^v \theta_v D^{p-v} m^*(u). \quad (9.7)$$

With the substitution of the expression for  $D^{p-v} m^*(u)$ , as it may be obtained from (4.3) by writing  $\bar{\gamma}_{i-s}$  in the place of  $\gamma_{i-s}$ , it is found that

$$L^*(u) = l^*(m^*(u)) - \frac{1}{\lambda^r} \sum_{j=1}^n \lambda^j \omega_j(z, \lambda) D^{n-j} u, \quad (9.8)$$

with

$$\omega_j(z, \lambda) = \sum_{v=1}^p \sum_{s=0}^p \lambda^{-s} \binom{p-v}{s} \theta_v D^s \bar{\gamma}_{\mu-v-s}.$$

A comparison of this with the earlier result (6.6) shows that

$$L^*(u) = L(u) - \frac{1}{\lambda^r} \sum_{j=1}^n \lambda^j \{ \epsilon_j(z, \lambda) + \omega_j(z, \lambda) \} D^{n-j} u. \quad (9.9)$$

The equation (9.1), whose solutions are completely known, thus has coefficients which differ from those of the given equation (2.1) only by terms that are of at least the  $r^{\text{th}}$  degree in  $1/\lambda$ . It is, therefore, by definition, a related equation.

#### REFERENCES

- [1] R. E. LANGER, The asymptotic solutions of ordinary linear differential equations of the second order with special reference to a turning point. *Trans. Amer. Math. Soc.*, vol. 67 (1949), pp. 461-490.
- [2] R. W. MCKELVEY, The solutions of second order linear ordinary differential equations about a turning point of order two. *Trans. Amer. Math. Soc.*, vol. 79 (1955), pp. 103-123.