Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 9 (1963)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON IMPLICIT FUNCTION THEOREMS AND THE EXISTENCE OF SOLUTIONS OF NON-LINEAR EQUATIONS

Autor: Ehrmann, Hans H.

Kapitel: 3. The implicit function theorem.

DOI: https://doi.org/10.5169/seals-38780

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.03.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Condition (2.4) is satisfied if (2.3) holds in the sphere

$$S: \| u - u_0 \| \leqq (1-l)^{-1} \| Vu_0 - u_0 \| . \tag{2.5}$$

Moreover, u is the limit of the sequence $\{ u_n \}$ where

$$u_{u+1} = Vu_n , \qquad n = 0, 1, 2, \dots ,$$

and there results the estimate

$$\| u - u_{n+1} \| \leqq l(1-l)^{-1} \| u_{n+1} - u_n \| \leqq l^{n+1} (1-l)^{-1} \| u_1 - u_0 \| . \tag{2.6}$$

3. The implicit function theorem.

Theorem 3.1. Let T^* be an operator with domain $D \subset B_1$ and range in B_2, let $S^* = S(u^*, r^*) \subset D$ and

$$T^* u^* = \theta . \tag{3.1}$$

We assume furthermore that there exists a linear operator K on S^* into B_2 with the following properties:

α) K has a bounded inverse, K^{-1}, defined on B_2 and
β) There exists a constant $m < \| K^{-1} \|^{-1}$ such that

$$\| T^* v - T^* u - K(v-u) \| \leqq m \| v - u \| \quad \text{for} \quad u, v \in S^* . \tag{3.2}$$

Then there exists an $\Omega = (u^*, r, a, b)$-neighborhood of T^*, such that for all $T \in \Omega$ the equation

$$Tu = \theta , \tag{3.1a}$$

has a unique solution $u = u(T)$ in $S(u^*, r)$. This solution is continuous in T at $T = T^*$ in the sense

$$\| u(T) - u^* \| \to 0 \quad \text{as} \quad \| Tu^* \| \to 0 . \tag{3.3}$$

In this theorem the operators T and K need not be continuous.

Proof. Let T lie in a (u^*, r, a, b)-neighborhood of T^* with $r \leqq r^*$. Then by (3.2), with $\Delta T = T - T^*$, we have

$$\begin{aligned} \| Tv - Tu - K(v-u) \| &\leqq \| \Delta Tv - \Delta Tu \| \\ + \| T^*v - T^*u - K(v-u) \| &\leqq (b+m) \cdot \| v-u \| \qquad (3.4) \\ \text{for} \quad u, v \in S(u^*, r) &\subset S^*, \end{aligned}$$

and the equation

$$u = Vu \equiv K^{-1}(K-T)u, \quad u \in S(u^*, r) = S, \qquad (3.5)$$

is equivalent to (3.1 a), $u \in S$.

For every $b \geqq 0$ with $l = (b+m) \| K^{-1} \| < 1$, (3.4) yields

$$\| Vu - Vv \| = \| K^{-1}[K(u-v) - Tu + Tv] \| \leqq l \| u-v \|,$$
$$l < 1, \qquad \text{for} \qquad u, v \in S(u^*, r).$$

If

$$\| Vu^* - u^* \| = \| K^{-1} Tu^* \| < (1-l) r, \qquad (3.6)$$

then the assumptions of the contraction mapping theorem [Section 2 f] are satisfied. Thus, under these conditions, there exists a unique solution $u = u(T)$ in S satisfying the condition

$$\| u - u^* \| \leqq (1-l)^{-1} \| K^{-1} Tu^* \| \leqq (1-l)^{-1} \| K^{-1} \| \cdot \| Tu^* \|. \qquad (3.7)$$

This implies the continuity (3.3).

The inequality (3.6) is satisfied if $T \in \Omega$ with

$$a = [\| K^{-1} \|^{-1} - (b+m)] r.$$

This completes the proof.

This proof also gives quantitative conditions for r, a, b which are sufficient for the existence of a unique and continuous solution u of (3.1 a) in $S(u^*, r)$.

Supplement. The assertion of Theorem 3.1 is true for each Ω-neighborhood of T^* with $0 < r \leqq r^*$ and a, b satisfying

$$a = [\| K^{-1} \|^{-1} - (b+m)] r > 0. \qquad (3.8)$$

Then, for the solution $u = u(T)$ in S, the estimate (3.7) holds.

A unique solution of (3.1 a) in $S(u^*, r)$ also exists for such r and b if (3.6) holds, but in (3.8) the sign " $>$ " cannot be replaced by " $\geqq$ ", nor can the constant a in (3.8) be replaced by any larger one.

The last statement can be proved by simple examples in the one-dimensional case and with an operator T which is linear in $S(u^*, r)$.

4. Inverse function theorems.

Under the conditions of the implicit function Theorem 3.1, the operator T has a local inverse defined in a neighborhood of a point w_0 for which

$$Tu = w. \tag{4.1}$$

has a solution u_0. This inverse has its range in a neighborhood of u_0. For the proof set $T^*u = Tu - w_0$ in Theorem 3.1. However, the conditions of this theorem are still not sufficient for the existence of a solution u of equation (4.1) for all w in B_2 even if T is defined on the whole Banach space B_1 and the conditions are satisfied at each point u of B_1.[1])

However, this actually is not necessary for the existence of at least one solution u of (4.1) for all $w \in B_2$ as is indicated by the following theorem.

Theorem 4.1. Let the operator T, mapping a non-empty domain $D \subset B_1$ into B_2, satisfy the following conditions:

For each $u \in D$ there exist a sphere $S(u, r) \subset D$, a linear operator K, and a constant m such that the following conditions hold:

α) K has a bounded inverse K^{-1} on $TS(u, r)$

β) $\| Tv - T\tilde{v} - K(v - \tilde{v}) \| \leqq m \| v - \tilde{v} \|$ for $v, \tilde{v} \in S(u, r)$

γ) $(\| K^{-1} \|^{-1} - m)\, r \geqq c > 0$ where the constant c is independent of $u \in D$.

---

1) Example: $Tu \equiv \arctan u = w$, with $B_1 = B_2 = \{$real numbers$\}$, is not solvable for all $w \in B_2$, although the conditions of Theorem 3.1 are satisfied at each point $(u, w = \arctan u)$ for $T^*u = Tu - w$.