

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 10 (1964)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: VERTEX POINTS OF FUNCTIONS
Autor: Amir-Moéz, Ali R.
Kapitel: 2. Tangent Quadric
DOI: <https://doi.org/10.5169/seals-39423>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 22.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

VERTEX POINTS OF FUNCTIONS

by Ali R. AMIR-MOÉZ

For f a real function of n variables, usually the Hessian matrix is studied in connection with Gaussian and mean curvatures of $f(x_1, \dots, x_n)$. In this paper we study other properties of f in a neighborhood of a point. In particular we get a method for obtaining vertex points of the function f . We also generalize the idea to some complex cases.

1. DEFINITIONS AND NOTATIONS

Let f a function of complex variables x_1, \dots, x_n be of class C'' in x_1, \dots, x_n , and $\bar{x}_1, \dots, \bar{x}_n$, in a neighborhood of a point. Then f is called unitarily analytic if

$$\frac{\partial^2 f}{\partial x_i \partial \bar{x}_j} = \left(\overline{\frac{\partial^2 f}{\partial \bar{x}_i \partial x_j}} \right).$$

Theorem: Let f be of class C'' in $x_1, \dots, x_n, \bar{x}_1, \dots, \bar{x}_n$ in a neighborhood of a point, and

$$\frac{\partial f}{\partial \bar{x}_k} = \left(\overline{\frac{\partial f}{\partial x_k}} \right).$$

Then f is unitarily analytic.

The proof is quite simple and we omit it. Note that the converse is not necessarily true.

2. TANGENT QUADRIC

Let f be unitarily analytic in a neighborhood of (c_1, \dots, c_n) .

Let, for example, $\frac{\partial f}{\partial c_1}$ be the value of $\frac{\partial f}{\partial x_1}$ at (c_1, \dots, c_n) , and

$f_c = f(c_1, \dots, c_n)$. Then

$$(x_1 - c_1 \dots x_n - c_n) \begin{bmatrix} \frac{\partial^2 f}{\partial c_1 \partial \bar{c}_1} & \dots & \frac{\partial^2 f}{\partial c_1 \partial \bar{c}_n} \left(\frac{\partial f}{\partial c_1} \right) \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial c_n \partial \bar{c}_1} & \dots & \frac{\partial^2 f}{\partial c_n \partial \bar{c}_n} \left(\frac{\partial f}{\partial c_n} \right) \\ \frac{\partial f}{\partial c_1} & \frac{\partial f}{\partial c_n} & f_c \end{bmatrix} \begin{bmatrix} x_1 - c_1 \\ \vdots \\ x_n - c_n \\ 1 \end{bmatrix} = 0 \quad (2.1)$$

is called the tangent quadric of f at (c_1, \dots, c_n) . We shall study only the cases that at least one of the first or second derivatives is not zero. It is clear that the tangent plane of (2.1) at (c_1, \dots, c_n) is the same as the tangent plane of $f = 0$ at this point.

Let the matrix of (2.1) be A , $\xi = (x_1 - c_1 \dots x_n - c_n)$, and $\eta = (0 \dots 0 \ 1)$. Then by section 8 of [1]

$$\xi A \eta^* = 0 \quad (2.2)$$

is the tangent plane of (2.1) at (c_1, \dots, c_n) . Here η^* is the conjugate transpose of η .

We easily see that (2.2) can be written as

$$\sum_{i=1}^n \frac{\partial f}{\partial c_i} (x_i - c_i) = 0. \quad (2.3)$$

3. MATRICES RELATED TO f

Besides A there are other matrices of some interest. We denote the matrix of the quadratic form of (2.1) by Q . The projection on the normal and tangent plane are of some interest. We denote the projection on the normal by P , and clearly $I - P$ is the projection on the tangent plane where I is the identity matrix. It is easy to see that $P = (P_{ij})$, where