

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 11 (1965)
Heft: 2-3: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON ERMAKOF'S CONVERGENCE CRITERIA AND ABEL'S FUNCTIONAL EQUATION.
Autor: Ostrowski, A. M.
Kapitel: I. Introduction
DOI: <https://doi.org/10.5169/seals-39968>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ON ERMAKOF'S CONVERGENCE CRITERIA AND ABEL'S FUNCTIONAL EQUATION. *)

A.M. OSTROWSKI

I. INTRODUCTION

1. We owe to V. Ermakof ([1], [2]) very remarkable criteria for the convergence or divergence of infinite series $\Sigma f(\varphi)$ ($f(x) > 0$) which uses the quotient

$$\frac{f(\Psi(x)) \Psi'(x)}{f(x)} \quad (1)$$

for continuously differentiable function $\Psi(x)$ with the properties

$$\Psi(x) > x, \Psi(x) \rightarrow \infty (x \rightarrow \infty).$$

As a matter of fact, the first discussion given by Ermakof [1] only established directly the connection with the convergence or the divergence of the integral

$$\int^{\infty} f(x) dx \quad (2)$$

so that in order to obtain the results concerning the infinite series we have to assume that $f(x)$ is monotonically decreasing or to make some analogous assumptions to permit the transition from the integral to the infinite series. We discuss some conditions of this kind in the sections 33-38.

2. In his second paper [2] Ermakof developed however a new and very ingenious method of proof using Abel's functional equation

$$\varphi(\Psi(x)) = \varphi(x) + 1.$$

*) This investigation was carried out under the contract DA-91-591-EUC-2824 of the Institute of Mathematics, University of Basle, with the US Department of the Army.

This method allows, under suitable regularity conditions on $\Psi(x)$, to connect directly the behavior of (1) with the convergence or divergence of the infinite series $\sum f(v)$, *without any monotony condition for $f(x)$* .¹⁾

But Ermakof only sketched his discussion and indicated as the sufficient additional condition to impose on $\Psi(x)$ that, in our notations, $\Psi'(x_0)$ is ≥ 1 , for a suitable x_0 .²⁾

It appears however that this additional condition is not sufficient to carry the discussion through. In a paper [5], published 1955, I showed that if beyond Ermakof's condition $\Psi'(x)$ is supposed monotonically increasing, the method can be carried through, indeed. If on the other hand $\Psi'(x)$ is supposed monotonically decreasing the method worked but Ermakof's additional condition was not necessary.

3. In this communication I develop a new method of proof which allows to avoid Abel's functional equation and to obtain the essential results for not necessarily monotonic $f(x)$. This gives a direct and very elementary way of proof as well for monotonically increasing as, (in the case of convergence), for monotonically decreasing $\Psi'(x)$. Beyond that, this method allows also to prove the convergence criteria in the case that $\lim_{x \rightarrow \infty} \Psi'(x)$ exists and is finite (Theorems 4-6).

4. As to the divergence criterion, here too, a new result in the case of monotonically decreasing $\Psi'(x)$ can be obtained (Theorem 7), however, with a different method which has more points of contact with Ermakof's second proof — here we have to form a minorant of $f(x)$, which can be interpreted as the derivative of a solution of Abel's functional equation —.

5. In the first sections of this paper we give 3 Theorems concerning the convergence and divergence of the integral (2) generalizing some results given in our first paper [5]. Finally, in the last part of the paper we discuss Pringsheim's treatment

¹⁾ Curiously enough, Abel's functional equation was also treated by Korkine in the note [4] where he gave another and direct proof of Ermakof's criteria for monotonic $f(x)$, without using, however, this functional equation.

²⁾ Ermakof says in his paper [2] in a footnote on p. 142: "C'est la seule condition pour que notre démonstration soit juste."

of the problem and prove generalized versions of Pringsheim's results.

This note brings therefore an improvement and simplification of the sections I-V and XI of [5], while I have nothing to add to the sections VI-X of [5].

II. ERMAKOF'S DIRECT METHOD

6. The form of the expression (4) makes it plausible that we will have to use the integral transformation formula

$$\int_a^b f(\Psi(x)) \Psi'(x) dx = \int_{\Psi(a)}^{\Psi(b)} f(x) dx. \quad (3)$$

In order to be able to use (3) we have in any case to assume that $f(x)$ is integrable in the integration interval and $\Psi(x)$ totally continuous between a and b . However, additional conditions are necessary and two such conditions are known either of which ensures the relation (3):

- J_1 : $|f(x)|$ is uniformly bounded in the integration interval;
- J_2 : $\Psi(x)$ is monotonically increasing or monotonically decreasing.

7. THEOREM 1. Assume that $\psi(x)$ and $\Psi(x)$ are totally continuous for $x \geq x_0$ and that we have for a sequence $b_v \geq x_0$ ($v = 1, 2, \dots$)

$$\psi(b_v) \leq \Psi(b_v), \quad \Psi(b_v) \rightarrow \infty \quad (v \rightarrow \infty). \quad (4)$$

Let $f(x)$ be ≥ 0 on no half-line $x \geq \xi$ almost everywhere $= 0$, and measurable in an interval J containing all values of $\psi(x)$ and $\Psi(x)$ for $x \geq x_0$. Assume further that for any finite subinterval of J the transformation formula (3) holds as well for $\psi(x)$ as for $\Psi(x)$. Then, if we have for almost all x with $x \geq x_0$ and for an α with $0 < \alpha < 1$:

$$f(\Psi(x)) \Psi'(x) \leq \alpha f(\psi(x)) \psi'(x) \quad (x \geq x_0), \quad 0 < \alpha < 1, \quad (5)$$

the integral (2) is convergent and we have for all $x \geq x_0$:

$$\Psi(x) > \psi(x) \quad (x \geq x_0). \quad (6)$$