Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	12 (1966)
Heft:	4: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	ON L(p,q) SPACES
Autor:	Hunt, Richard A.
Kapitel:	Section 5. References
DOI:	https://doi.org/10.5169/seals-40747

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Proof. From (2.7), we have

$$||f^*g||_{p1}^* = \sup_h B | \int f^*g(x) h(x) dm(x) |,$$

where $h^{*}(t) \leq t^{(1/p)-1}$. Let $I(h) = \int f^{*}g(x) h(x) dm(x)$.

$$|I(h)| \leq \int (\int |f(y)| \cdot |g(xy^{-1})| dm(y)| h(x) | dm(x)$$

= $\int |f(y)| (\int |g(xy^{-1})| \cdot |h(x)| dm(x)) dm(y).$

Hence, $|I(h)| \leq ||fk||_{11}^*$, where $k(y) = \int |g(xy^{-1})| \cdot |h(x)| dm(x)$. By the multiplication theorem it follows that $|I(h)| \leq B ||f||_{p_0q_0}^* ||k||_{p_0q_1}^*$, where $1/p_0 + 1/p'_0 = 1$. But $k = |\bar{g}|^* |h|$, where $\bar{g}(x) = g(x^{-1})$. Hence by Lemma 4.8,

$$|| k ||_{p'_0q_1}^* \leq B || \bar{g} ||_{p_1q_1}^* || h ||_{p'^{\infty}}^* \leq B || \bar{g} ||_{p_1q_1}^*.$$

Since (G, dm) is unimodular, we have $(\bar{g})^*(t) = g^*(t)$ and the lemma follows.

By applying the strong type interpolation theorem to the end point results of Lemma 4.8 and Lemma 4.9, we obtain

THEOREM 4.10. (Convolution theorem):

$$||f^*g||_{pq}^* \leq B||f||_{p_0q_0}^*||g||_{p_1q_1}^*,$$

where $0 < 1/p = 1/p_0 + 1/p_1 - 1 < 1$, $1 < p_0, p_1 < \infty$ and $0 \le 1/q = 1/q_0 + 1/q_1 \le 1$.

Section 5. REFERENCES

Various properties of L(p, q) spaces have appeared in many places, often as special cases of a more general theory. We will mention several places where related results and applications are found. The references given are not necessarily the first or the only place where the indicated result appears.

The principal references are [19] and [20], where G. G. Lorentz defines special cases of L(p, q) spaces and proves many of their properties. The notion of a non-increasing rearrangement of a function was used by Hardy Littlewood and Payley. (See [32].) A simple proof of the inequality $||f||_{pq_2}^*$ $\leq B ||f||_{pq_1}^*, q_1 \leq q_2$, is found in O'Neil [22]. The technique used in the proof

— 273 —

of Hardy's inequality is well known. A different proof of Hardy's inequality is found in [9, p. 245] or [32, Vol. I, p. 20]. The author learned the simple proof of inequality (1.9) from class notes from a course given by A. Zygmund in Chicago. A similar proof is given in [11, p. 278].

The L(p, q) spaces which are Banach spaces appear as intermediate spaces in the general interpolation theory of Calderón [1]. Peetre [24] identifies L(p, q) spaces as intermediate spaces for the interpolation theory of Lions and Peetre [18]. Many L(p, q) results are then contained in these general theories. In particular, there are results concerning density, inclusion, separability and duality of the spaces. A ** norm is used in these results. Riviere [25] generalized the results of Calderón [1] to include L(p, q), p, q > 0. Similarly, P. Kree and J. Peetre generalized the results of Lions and Peetre [18].

(2.5) is proved by Krein and Semenov [18] and is contained implicitly in Stein and Weiss [30]. Halperin [8] and [9] obtains general results on conjugate spaces and reflexitivity. Results on uniform convexity of some related spaces are found in Halperin [10]. The results concerning linear functionals on L(p, q), p < 1, correspond to results of Day [5] for L^p spaces 0 .

The weak type theorem of Section 3 restricted to linear operators on the L(p, q) spaces which are Banach spaces was proved by A. P. Calderón [2]. We learned that E. M Stein also obtain these results. Proof of these cases is found in Lions and Peetre [18], together with Peetre [24]. Also see Calderón [1] and Oklander [21]. Krein and Semonov [17] prove some special cases. The theorem is closely related to results of Stein and Weiss [29] and [30]. The weak type theorem for L(p, q), p, q > 0, is proved in Hunt [14]. These cases are also contained in work of P. Kree and J. Peetre.

The strong type theorem of Section 3 for linear operators on the L(p, q) spaces which are Banach spaces is found in Calderón [1]. These results are related to results of Hirschman [13], Stein [26] and Stein and Weiss [27]. The result for sublinear operators follows ideas found in Calderón [1], Calderón and Zygmund [3] and Weiss [31]. Rivière [25] obtains results for linear operators acting on L(p, q) spaces, p, q > 0.

Stein [26] proves an analogue of Theorem 4.3 for Fourier coefficients. The multiplication and convolution theorems are proved by a different method in O'Neil [22]. E. M. Stein also obtained these results. (See [22].)