

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 13 (1967)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: BOUNDEDNESS THEOREMS FOR SOLUTIONS OF $u''(t) + a(t)f(u)g(u') = 0$ (IV)
Autor: Wong, James S. W.
Kapitel: 1. Introduction
DOI: <https://doi.org/10.5169/seals-41539>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

BOUNDEDNESS THEOREMS FOR
SOLUTIONS OF $u''(t) + a(t)f(u)g(u') = 0$ (IV)

by James S. W. WONG

1. INTRODUCTION

In our previous work [1-3], we have presented rather fragmentary results concerning the boundedness of solutions to certain second order non-linear differential equations of the following form:

$$u''(t) + a(t)f(u)g(u') = 0 \quad (1.1)$$

where $a(t)$, $f(u)$ and $g(u')$ satisfy certain assumptions to be described below. The purpose of the present paper is to further extend these results and establish comparison theorems. Some of our results presented here may be considered as generalizations to the results of Zhang [4], where a special case of equation (1.1):

$$u''(t) + a(t)f(u) = 0 \quad (1.2)$$

was treated ¹⁾.

Throughout the discussion of this paper, we will need the following assumptions:

(A₁) $g(u')$ is a positive continuous function of u' ,

(A₂) $f(u)$ is a continuous function of u satisfying $uf(u) > 0$, if $u \neq 0$,

(A₃) $a(t)$ is continuous in t ,

$$(A_4) \lim_{|u| \rightarrow \infty} \int_0^u f(s) ds = \infty,$$

$$(A_5) \lim_{|v| \rightarrow \infty} \int_0^v \frac{g(s)}{s} ds = \infty.$$

We also list in the following a brief résumé of our previous results on boundedness.

¹⁾ For other boundedness result concerning (1.2), see [5]-[7], [13], [14].

Theorem (I). Suppose that assumptions $A_1, A_2, A_3,$ and A_4 hold and in addition that $a(t) > 0$ and $a'(t) \geq 0$ for $t \geq T$. Then all solutions of (1.1) are bounded.

Corollary. In addition to the hypothesis of Theorem (I), suppose that assumption A_5 also holds and that $\lim_{t \rightarrow \infty} a(t) = k > 0$; then all solutions of (1.1) and their derivatives are bounded.

Theorem (II). Suppose that assumptions A_1, A_2, A_3 and A_4 hold and in addition that $a'(t) \leq 0$ for $t \geq T$. Then all solutions of (1.1) are bounded.

Corollary. In addition to the hypothesis of Theorem (II), suppose that assumption A_5 also holds and $\lim_{t \rightarrow \infty} a(t) = k > 0$; then all solutions of (1.1) and their derivatives are bounded.

Theorem (III). Suppose that assumptions $A_1, A_2, A_3,$ and A_4 hold and in addition that $a(t) \geq a_0 > 0$ for $t \geq T$, and $\int_0^{\infty} |a'(t)| dt < \infty$. Then all solutions of (1.1) are bounded.

Corollary. In addition to the hypothesis of Theorem (III), suppose that assumption A_5 also holds; then all solutions of (1.1) and their derivatives are bounded.

The method of proof for the above results is based essentially on the well-known lemma of Gronwall [10], which is also known as the Bellman's lemma. In this paper, we use in addition to this fundamental lemma, its generalizations [11], [12], and techniques borrowed from Lyapunov's stability theory.

It might be of interest to note that quite a few results in [4] are incorrect; in particular Theorems 5 and 6. Also, Theorems 3 and 4 are stated incorrectly.

2. BOUNDEDNESS THEOREMS I

Theorem 1. Suppose that assumptions A_1, A_2, A_3 and A_4 hold and that $a(t) > 0$ for $t \geq T$ and there exists a non-negative function $\alpha(t)$ such that $-a'(t) \leq \alpha(t)a(t)$ with $\int_0^{\infty} \alpha(s) ds < \infty$. Then all solutions of (1.1) are bounded.