

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 13 (1967)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: BOUNDEDNESS THEOREMS FOR SOLUTIONS OF $u''(t) + a(t)f(u)g(u') = 0$ (IV)
Autor: Wong, James S. W.
Kapitel: 3. BOUNDEDNESS THEOREMS II
DOI: <https://doi.org/10.5169/seals-41539>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$(b) \int |a(s)| s^\alpha ds < \infty,$$

$$(c) 0 < g(v) \leq K \text{ for all } v;$$

then the derivative of any solution of (1.1) has a limit.

Proof. Proceeding as in the above proof, we obtain instead of (3.2) the following estimate:

$$\frac{|u(t)|}{t} \leq (|u(t_0)| + |u'(t_0)|) + \int_{t_0}^t s^\alpha KM |a(s)| h\left(\frac{|u(s)|}{s}\right) ds,$$

from which we conclude from a result of Bihari [14] that

$$\frac{|u(t)|}{t} \leq H^{-1} (H(|u(t_0)| + |u'(t_0)|) + KM \int_{t_0}^t |a(s)| s^\alpha ds)$$

which is bounded for t on account of assumption (a). The remaining proof follows verbatim that of Theorem 3.

3. BOUNDEDNESS THEOREMS II

Theorem 5. Suppose that assumptions A_1, A_2, A_3 and A_4 hold and in addition that

$$(i) a(t) > 0, \quad a'(t) \geq 0, \quad \text{for } t \geq T,$$

$$(ii) \frac{d}{dt} \left(\frac{b}{a} \right) \leq \beta(t) \left(1 + \frac{b}{a} \right), \quad \text{with } \int_0^\infty \beta(s) ds < \infty$$

and

$$\left(1 + \frac{b}{a} \right) \geq \varepsilon > 0;$$

then every solution of (1.1) with $(a(t) + b(t))$ replacing $a(t)$ is bounded.

Proof. Make the following substitution for the independent variable, $x = \int_0^t \sqrt{a(s)} ds$ which tends to infinity as $t \rightarrow \infty$, and obtain instead of (1.1) its transformed equation:

$$\frac{d^2 u}{dx^2} + \frac{1}{2} \left(\frac{a}{a^{3/2}} \right) \frac{du}{dx} + \left(1 + \frac{b}{a} \right) f(u) g(u') = 0 \quad (3.1)$$

where “dot” denotes differentiation with respect to t . Now write equation in its system form, letting $y_1 = u$:

$$\begin{cases} \frac{d y_1}{d x} = y_2 \\ \frac{d y_2}{d x} = -\frac{1}{2} \left(\frac{a}{a^{3/2}} \right) y_2 - \left(1 + \frac{b}{a} \right) f(y_1) g(\sqrt{a} y_2). \end{cases} \quad (3.2)$$

Define for (3.2) the following function:

$$V(x, y_1, y_2) = \left(1 + \frac{b}{a} \right) \int^{y_1} f(s) ds + \int^{y_2} \frac{s ds}{g(\sqrt{a} s)},$$

and observe:

$$\begin{aligned} \frac{dV}{dx} &\leq \frac{\beta(t)}{\sqrt{a(t)}} \left(1 + \frac{b}{a} \right) \int^{y_1} f(s) ds - \frac{1}{2} \frac{a}{a^{3/2}} y_2^2 \\ &\leq \frac{\beta(t)}{\sqrt{a(t)}} V. \end{aligned}$$

Hence we have

$$V(x, y_1, y_2) \leq V(x(T), y_1(x(T)), y_2(x(T))) \exp \int_T^t \beta(s) ds$$

which is finite. From (ii) we note that $V \rightarrow \infty$ as $y_1 \rightarrow \infty$ and $V > 0$ if $y_1^2 + y_2^2 \neq 0$. Thus, every solution of (1.1) is bounded.

Corollary. Suppose in addition to the hypothesis of Theorem 5 that assumption A_5 also holds and that $\lim_{t \rightarrow \infty} a(t) = a_1 < \infty$, then every solution of (1.1) and its derivative are bounded.

From the above result we may conclude for example that all solutions of the following equation:

$$u''(t) + (c_1 t^\alpha + c_2 t^\beta) u^\lambda(t) (1 + \exp u'(t) \sin u'(t)) = 0$$

are bounded for all $c_1, c_2 > 0$, $\alpha > \beta \geq 0$, and $\lambda > 0$.

We now consider the following inhomogeneous equation:

$$u''(t) + a(t) f(u) g(u') = h(t, u, u') \quad (3.3)$$

and assume that $|u' h(t, u, u')| \leq \gamma(t) g(u')$ where $\int_0^\infty \gamma(s) ds < \infty$.

Theorem 6. Suppose that assumptions A_1, A_2, A_3 and A_4 hold and in addition that $a(t) > 0$ and $a'(t) \geq 0$ for $t \geq T$; then all solutions of (3.3) are bounded.

Proof. Integrate (3.3) in the following manner:

$$G(u'(t)) - G(u'(t_0)) + a(t)F(u(t)) - a(t_0)F(u(t_0)) \\ = \int_{t_0}^t a'(s)F(u(s))ds + \int_{t_0}^t \frac{h(t, u, u')u'(s)ds}{g(u')} \quad (3.4)$$

where $G(v) = \int_0^v \frac{s ds}{g(s)}$ and $F(u) = \int_0^u f(s) ds$. Taking absolute values and noting that $G(v) \geq 0$ and $F(u) \geq 0$, we obtain

$$a(t)F(u(t)) \leq c_0 + c_1 + \int_{t_0}^t a'(s)F(u(s))ds \quad (3.5)$$

where $c_0 = G(u'(t_0)) + a(t_0)F(u(t_0))$ and $c_1 = \int_{t_0}^{\infty} \gamma(s) ds$ are non-negative constants. From (3.5) and A_4 it is now clear that every solution of (3.3) are bounded (cf. [1]).

Corollary. In addition to the hypothesis of Theorem 6, suppose that assumption A_5 also holds and that $\lim_{t \rightarrow \infty} a(t) = k > 0$; then all solutions of (3.3) and their derivatives are bounded.

We note that by setting $h(t, u, u') \equiv 0$, the above result again reduces to Theorem 1 and its corollary. Other comparison theorems may be formulated in a similar way as Theorem 6 by extending the corresponding result for the homogeneous equation. Since the procedure is clear, the statements and proofs of these results will be omitted.

RÉFÉRENCES

- [1] J. S. W. WONG, A note on boundedness theorems of certain second order differential equations. *SIAM Review*, 6 (1964), 175-176.
- [2] — and T. A. BURTON, Some properties of solutions of $u''(t) + a(t)f(u)g(u') = 0$ (II). *Monatshefte für Mathematik*, 69 (1965), 368-374.
- [3] J. S. W. WONG, Some properties of solutions of $u''(t) + a(t)f(u)g(u) = 0$ (III). *SIAM Journal*, 14 (1966), 209-214.
- [4] ZHANG BING-GEN (CHANG PING-KEN), Boundedness of solutions of ordinary differential equations of the second order. *Acta Math. Sinica*, 14 (1964), 128-137; English translation, *Chinese Mathematics*, 4 (1964), 139-148.