Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 14 (1968)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: NOTE ON ABSOLUTE SUMMABILITY FACTORS
Autor: Irwin, Ronald Lee
DOI: https://doi.org/10.5169/seals-42356

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 09.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

A NOTE ON ABSOLUTE SUMMABILITY FACTORS

by Ronald Lee IRWIN [1])

In this paper results which appear in [2] are extended, namely Theorem 3. The notation used here and explanation of the problems are given there. The notation $\varepsilon_\nu \in (|A|, |B|)_r$ means $\sum a_\nu \varepsilon_\nu \in |B|$ whenever $\sum a_\nu \in |A|$.

The following theorem is a special case of Theorem 3 in [2]. However, strictly speaking somewhat stronger, since it does not require the representation (4) in [2]. This is important later on.

THEOREM 1. If A is absolutely regular, normal, $A' \dot{\leqq} 0$ and $a_{\nu\nu} > 0\downarrow$ for $\nu\uparrow$ ($A' \dot{\leqq} 0$ and $a_{nn} > 0$ imply $A \dot{\geqq} 0$, see footnote 1) in [2]), then $\varepsilon_\nu = 0\,(a_{\nu\nu})$ implies $\varepsilon_\nu \in (|A|, |I|)_r$.

Proof. It suffices to prove $\sum_{n=\nu}^{\infty} |a'_{n\nu}\varepsilon_n| \leqq M$. Since $\sum_{n=\nu}^{\infty} |a'_{n\nu}| = \frac{2}{a_{\nu\nu}} - 1$ (see, Lemma 1 in [1]), and $|\varepsilon_n| \leqq K a_{nn}$, we have $\sum_{n=\nu}^{\infty} |a'_{n\nu}\varepsilon_n| \leqq 2K$ using $a_{nn}\downarrow$.

Proofs of the following lemmas can be found in [1]. The notation $|B| \subseteq |A|$ means $\sum a_\nu \in |A|$ whenever $\sum a_\nu \in |B|$.

Lemma 1. If A and B are absolutely regular and normal, and if $|B| \subseteq |A|$, then given a bounded sequence $c = \{c_\nu\}$ there exists a bounded sequence $c' = \{c'_\nu\}$ such that $\varepsilon_\nu(A, c) = \varepsilon_\nu(B, c')$.

Lemma 2. Let $A = BP$, P a weighted mean. Then,

$$[\varepsilon_{\nu+1}(A, c) - \varepsilon_\nu(A, c)]\frac{P_{\nu-1}}{p_\nu} = \varepsilon_\nu(A, c) - \varepsilon_\nu(B, c).$$

We now extend Theorem 1 to matrices of the form AP^k where k is a positive integer and P is a weighted mean. If P is a weighted mean, then

$$p_{n\nu} = \frac{p_n}{P_n P_{n-1}} P_{\nu-1} \; (p_{00} = 1).$$

[1]) This research was supported by NSF contract GP-9926.

THEOREM 2. Suppose P is a weighted mean with $p_\nu > 0$, $a_{\nu\nu} > 0\downarrow$ $(\nu\uparrow)$, and $p_{\nu+1} = 0(p_\nu)$. If $\varepsilon_\nu = 0(a_{\nu\nu})$ implies $\varepsilon_\nu \in (|A|, |I|)_r$, then $\delta_\nu = 0\left(\frac{a_{\nu\nu}\, p_\nu}{P_\nu}\right)$ implies $\delta_\nu \in (|AP|, |I|)_r$.

Proof. It suffices to prove $\sum |a_\nu a_{\nu\nu} \frac{p_\nu}{P_\nu}| < \infty$ whenever $\sum a_\nu \in |AP|$. Using the identity

$$a_\nu = \frac{1}{P_{\nu-1}} \sum_{k=1}^{\nu} P_{k-1} a_k - \frac{1}{P_{\nu-1}} \sum_{k=1}^{\nu-1} P_{k-1} a_k \, (\nu \geqq 1)$$

we conclude

$$\sum_{\nu=1}^{\infty} |a_\nu \frac{a_{\nu\nu}\, p_\nu}{P_\nu}| \leqq \sum_{\nu=1}^{\infty} |a_{\nu\nu} P_\nu(a)| + \sum_{\nu=1}^{\infty} |\frac{a_{\nu+1,\nu+1}}{a_{\nu\nu}} \cdot \frac{p_{\nu+1}}{p_\nu} \cdot a_{\nu\nu} P_\nu(a)|$$

($P_\nu(a)$ denotes the P-transform of $\sum a_\nu$.) Since, $\sum P_\nu(a) \in |A|$ we have $\sum |a_{\nu\nu} P_\nu(a)| < \infty$ by hypothesis. The same holds true for the second sum, since all of the additional factors are bounded. This completes the proof. To conclude the results of Theorem 2 with AP^k (k an integer, $k > 1$) in place of AP we need only show $a_{\nu+1,\nu+1} = 0(a_{\nu\nu})$ implies $(AP^k)_{\nu+1,\nu+1} = 0((AP^k)_{\nu\nu})$. This is done inductively using $p_{\nu+1} = 0(p_\nu)$. Thus we have by induction.

THEOREM 3. Suppose P is a weighted mean which satisfies the hypotheses of Theorem 2. If $\varepsilon_\nu = 0(a_{\nu\nu})$ implies $\varepsilon_\nu \in (|A|, |I|)_r$, then $\delta_\nu = 0\left(a_{\nu\nu}\left(\frac{p_\nu}{P_\nu}\right)^k\right)$ implies $\delta_\nu \in (|AP^k, |I|)_r$.

We now state and prove the main result of the paper.

THEOREM 4. Suppose A and B are normal, and absolutely regular. Let P be a weighted mean with $p_\nu > 0$ and $|AP| \supseteq |A|$. If

(1) $$\delta_\nu = 0(a_{\nu\nu}) \textit{ implies } \delta_\nu \in (|A|, |I|)_r$$

and

(2) $$\varepsilon_\nu(A, c) = 0\left(\frac{a_{\nu\nu}}{b_{\nu\nu}}\right) \textit{ implies } \varepsilon_\nu \in (|A|, |B|)_r,$$

then

$$\varepsilon_\nu(AP, c) = 0\left(\frac{a_{\nu\nu}\, p_\nu}{b_{\nu\nu} P_\nu}\right) \textit{ implies } \varepsilon_\nu(AP, c) \in (|AP|, |B|)_r$$

whenever

$$b_{n\nu} \uparrow \textit{ for } \nu \uparrow (\nu \leqq n),$$

$$a_{\nu+1,\,\nu+1} = 0(a_{\nu\nu}),\ b_{\nu\nu} = 0(b_{\nu+1,\,\nu+1}),\ \textit{and}\ p_{\nu+1} = 0(p_\nu).$$

Proof. If $\sum a_\nu \in |AP|$, then $\sum P_\nu(a) \in |A|$. Since $p_\nu > 0$, we have $a_\nu = \sum_{\mu=0}^{\nu} p'_{\nu\mu}\,\alpha_\mu$ where $\sum \alpha_\mu \in |A|$. Using this we have

$$\beta_n = \sum_{\nu=0}^{n} b_{n\nu}\,\varepsilon_\nu(AP, c)\,a_\nu = \sum_{\mu=0}^{n} \alpha_\mu \sum_{\nu=\mu}^{n} b_{n\nu}\,\varepsilon_\nu(AP, c)\,p'_{\nu\mu}.$$

The matrix $P' = (p'_{\nu\mu})$ is given by

$$p'_{\nu\nu} = \frac{P_\nu}{p_\nu},\ p'_{\nu,\nu-1} = -\frac{P_{\nu-2}}{p_{\nu-1}},\ p_{\nu\mu} = 0 \ \textit{otherwise}.$$

Introducing the inverse we have

$$\beta_n = \sum_{\mu=0}^{n} \alpha_\mu\, b_{n\mu}\,\varepsilon_\mu(AP, c) + \sum_{\mu=0}^{n} \alpha_\mu \frac{P_{\mu-1}}{p_\mu}\big(b_{n\mu}\,\varepsilon_\mu(AP, c) - b_{n\mu+1}\,\varepsilon_{\mu+1}(AP, c)\big).$$

Write

$$b_{n\mu}\,\varepsilon_\mu(AP, c) - b_{n,\mu+1}\,\varepsilon_{\mu+1}(AP, c) = b_{n\mu}(\varepsilon_\mu(AP, c) - \varepsilon_{\mu+1}(AP, c)) + \varepsilon_{\mu+1}(AP, c)(b_{n\mu} - b_{n\mu+1}).$$

From this it follows by Lemmas 1 and 2 that

$$\sum_{\nu=0}^{n} b_{n\nu}\,\varepsilon_\nu(AP, c)\,a_\nu = \sum_{\mu=0}^{n} b_{n\mu}\,\varepsilon_\mu(A, c')\,\alpha_\mu + \sum_{\mu=0}^{n} \alpha_\mu \frac{P_{\mu-1}}{p_\mu}\,\varepsilon_{\mu+1}(AP, c)(b_{n\mu} - b_{n\mu+1}), \tag{3}$$

where $\varepsilon_\nu(A, c') = 0\left(\dfrac{a_{\nu\nu}}{b_{\nu\nu}}\right)$ by Lemma 2. Thus by (2) the first term is absolutely convergent. Write the second term of (3) in the form $\sum_{\mu=0}^{n} a_{\mu\mu}\,\alpha_\mu\,A_{n\mu}$, where

$$A_{n\mu} = \frac{1}{a_{\mu\mu}} \frac{P_{\mu-1}}{p_\mu} \times \varepsilon_{\mu+1}(AP, c)(b_{n\mu} - b_{n\mu+1}).$$

$$\sum_{n=\mu}^{\infty} |A_{n\mu}| = 0\left(\frac{b_{\mu\mu}}{a_{\mu\mu}} \frac{P_{\mu-1}}{P_\mu} \frac{a_{\mu+1,\mu+1}}{b_{\mu+1,\mu+1}} \frac{p_{\mu+1}}{p_\mu}\right)$$

By hypothesis (1) $\sum | a_{\mu\mu} \alpha_\mu | < \infty$. Hence the second term of (3) is absolutely convergent if $\sum_{n=\mu}^{\infty} | A_{n\mu} | \leqq M$. Since $b_{n\mu}\uparrow$ for $\mu\uparrow$ we have

which is 0 (1).

THEOREM 5. Let A, B be normal and absolutely regular with $A \dot{>} 0$, $B \dot{>} 0$, and $A' \dot{\leqq} 0$. Furthermore, assume

$$a_{n\nu}\uparrow, \quad \nu\uparrow (\nu \leqq n) \tag{4}$$

$$b_{n\nu}\uparrow, \quad \nu\uparrow (\nu \leqq n) \tag{5}$$

$$\frac{a_{n\nu}}{b_{n\nu}} \downarrow, \quad \nu\uparrow (\nu \leqq n) \tag{6}$$

$$\frac{b_{n\nu}}{a_{n\nu}} (a_{kn} - a_{k\nu}) \downarrow, \quad \nu\uparrow \textit{ for all } n \leqq k \tag{7}$$

$$a_{\nu+1,\nu+1} = 0 (a_{\nu\nu}) \tag{8}$$

$$b_{\nu\nu} = 0 (b_{\nu+1,\nu+1}) \tag{9}$$

$$p_{\nu+1} = 0 (p_\nu) . \tag{10}$$

When these conditions are satisfied

$$\varepsilon_\nu (AP^k, c) = 0 \left(\frac{a_{\nu\nu}}{b_{\nu\nu}} \left(\frac{p_\nu}{P_\nu} \right)^k \right)$$

implies

$$\varepsilon_\nu (AP^k, c) \in (| AP^k |, | B |)_r .$$

Proof. With $k = 0$ these condition imply $\varepsilon_\nu (A,c) \in (|A|, |B|)_r$ (see Theorem 3 in [2]. Hence, the theorem follows by induction from Theorem 3 and 4.

Theorem 5 extends Theorem 3 in [2] to include all Cesaro methods $A = (C, \alpha)$, $B = (C, \beta)$ where $\alpha \geqq 0$, and $0 \leqq \beta \leqq 1$.

BIBLIOGRAPHY

[1] IRWIN, R. L., Absolute Summability Factors I, *Tohoku Math. Journal, 1.*, 247-254 (1966).

[2] —— and A. PEYERIMHOFF, On Absolute Summability Factors, to appear in *L'Enseignement Mathématique.*

(Reçu le 10 avril 1969.)

University of Missouri-St. Louis