

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 14 (1968)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ANALYTIC SPACES
Autor: Malgrange, Bernard
Kapitel: 4.1. Stein spaces
DOI: <https://doi.org/10.5169/seals-42341>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ly, let \mathcal{A} be a sheaf of \mathcal{O}_Y -algebras, which is coherent as sheaf of \mathcal{O}_Y -modules. Then there exists an analytic space (X, \mathcal{O}_X) and a finite morphism $f : (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ such that $f_*(\mathcal{O}_X)$ is isomorphic with \mathcal{A} as sheaf of \mathcal{O}_Y -algebras ; the triple (X, \mathcal{O}_X, f) is unique up to an isomorphism.

We do not prove this proposition here and refer to Houzel [6] or Narasimhan [9] for this proof. We note also that a proof of the direct part can be given along the same lines as theorem 3.1.3, combined with the fact that direct images under finite morphism preserve exact sequences of sheaves of \mathcal{O}_X -modules (in other words, that higher direct images are zero). We note also that, for *proper* morphisms (not necessarily finite), a much deeper result has been proved by Grauert [2], [3].

Finally, we remark that, in the real case, proposition 3.3.2. is false (take, for instance, X the submanifold of \mathbf{R}^2 defined by $x_2 - x_1^2 = 0$, $Y = \mathbf{R}$ and $f =$ the projection on the x_2 -axis ; $f_*(\mathcal{O}_X)$ has support $x_2 \geq 0$, which is not an analytic subset of \mathbf{R} , hence $f_*(\mathcal{O}_X)$ cannot be coherent!)

CHAPTER 4.

THE FINITENESS THEOREM

In this chapter, we consider only *complex* analytic spaces, separated and having a countable basis of open sets.

4.1. Stein spaces

Let (X, \mathcal{O}_X) be an analytic space, and K a subset of X ; we denote, as usual by \hat{K} the set

$$\left\{ x \in X \mid \forall f \in \Gamma(X, \mathcal{O}_x) : |f(x)| \leq \sup_{y \in K} |f(y)| \right\}$$

Definition 4.1.1. a) (X, \mathcal{O}_X) is called holomorphically convex if, for any K compact $\subset X$, \hat{K} is compact ;
 b) (X, \mathcal{O}_X) is called a Stein space if it is holomorphically convex, and if, for any $x \in X$, there exist sections $f_1, \dots, f_p \in \Gamma(X, \mathcal{O}_X)$ with $f_i(x) = 0$, such that x is an isolated point of the counter-image of 0 in the morphism $(X, \mathcal{O}_X) \rightarrow \mathbf{C}^p$ defined by f_1, \dots, f_p , (This last property can also be expressed as the fact that the morphism of germs : $(X, \mathcal{O}_X, x) \rightarrow (\mathbf{C}^p, 0)$ defined by f_1, \dots, f_p is finite).

If X is a Stein space, X_{red} is obviously also a Stein space. The converse is also true (see Grauert [2]).

Theorem 4.1.2. (“Theorems A and B” of Cartan-Oka). Let F be an analytic coherent sheaf over a Stein space (X, \mathcal{O}_X) . Then

- 1) For any $x \in X$, $\Gamma(X, F)$ generates F_x over $\mathcal{O}_{X,x}$
- 2) For $p \geq 1$, one has $H^p(X, F) = 0$

This theorem will not be proved here (see f.i. [5] for the reduced case ; the general case is similar). We will need here only the following special case :

Let (X, \mathcal{O}_X) be a closed analytic subspace of a domain of holomorphy $U \subset \mathbf{C}^n$; if F is an analytic coherent sheaf on X , let \tilde{F} be the trivial extension of F to U ; then \tilde{F} is a coherent sheaf of \mathcal{O}_U -modules, and theorems A and B are valid for \tilde{F} : therefore, they are true for F .

4.2. Topology on $\Gamma(X, F)$.

1. Let X be a closed analytic subspace of a domain of holomorphy $U \subset \mathbf{C}^n$; and, with the previous notations, suppose that \tilde{F} admits a *finite presentation* i.e. an exact sequence of sheaves of \mathcal{O}_U -modules

$$\mathcal{O}_U^q \xrightarrow{\alpha} \mathcal{O}_U^p \xrightarrow{\beta} \tilde{F} \rightarrow 0.$$

Applying theorem B to the exact sequences

$$0 \rightarrow \text{Im } \alpha \rightarrow \mathcal{O}_U^p \rightarrow \tilde{F} \rightarrow 0 \quad \text{and} \quad 0 \rightarrow \text{Ker } \alpha \rightarrow \mathcal{O}_U^q \rightarrow \text{Im } \alpha \rightarrow 0$$

we get an exact sequence

$$\Gamma(U, \mathcal{O}_U)^q \xrightarrow{\Gamma(U, \alpha)} \Gamma(U, \mathcal{O}_U)^p \xrightarrow{\Gamma(U, \beta)} \Gamma(U, \tilde{F}) \rightarrow 0.$$

The space $\Gamma(U, \mathcal{O}_U)$, with the topology of uniform convergence on compact sets is a Frechet space. And we claim that, for that topology, $\text{Im } \Gamma(U, \alpha)$ is closed. For, if f is adherent to $\text{Im } \Gamma(U, \alpha)$, it results easily from Krull's theorem (see Appendix) that, for $x \in U$, we have $f_x \in \text{Im } (\alpha_x)$, hence $f \in \Gamma(U, \text{Im } \alpha)$; but, according to theorem B, the mapping $\Gamma(U, \mathcal{O}_U)^q \rightarrow \Gamma(U, \text{Im } \alpha)$ is surjective.

Now, with the quotient topology, $\Gamma(X, F) \simeq \Gamma(U, \tilde{F}) \simeq \Gamma(U, \mathcal{O}_U)/\text{Im } \Gamma(U, \alpha)$ is a Frechet space. This topology does not depend on the given presentation of \tilde{F} (in fact, it does not even depend on the imbedding $X \rightarrow U$, but we shall not need it here). For, suppose we have a second presentation