Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 14 (1968)

Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: MEROMORPHIC MAPPINGS

Autor: Stein, K.

Kapitel: 1. Correspondences

DOI: https://doi.org/10.5169/seals-42342

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 19.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

MEROMORPHIC MAPPINGS

by K. STEIN

Introduction

We study meromorphic mappings of complex spaces. The notion of meromorphic mapping we use was introduced by Remmert [9], [11]¹. Some part of the material dealt with in these lectures is contained in [15] and we shall therefore not give proofs for all statements.

The first sections are preliminary. The concept of correspondence is discussed and used to define meromorphic mappings (these are not mappings in the usual sense). Extension problems are studied in Section 4. Essential use is made of the extension theorem for analytic sets first proved by Thullen [21] in a special case and later generalized by Remmert and Stein [13]. The final section deals with maximal meromorphic mappings.

1. Correspondences

Let X and Y be sets. A correspondence, denoted $f: X \to Y$, assigns to each $x \in X$ a subset $f(x) \subset Y$, which may be empty. $f: X \to Y$ is called empty if $f(x) = \emptyset$ for all $x \in X$. For $A \subset X$ we set $f(A) = \bigcup_{x \in A} f(x)$. A mapping $\varphi: X \to Y$ is looked upon as a special correspondence (we do not distinguish between a set consisting of one element and the element).

Each correspondence $f: X \to Y$ can be characterized by its graph $G_f = \{(x, y) \mid x \in X, y \in f(x)\} \subset X \times Y$. The projection maps of G_f into X and Y are denoted by $f: G_f \to X$ and $f: G_f \to Y$. Then, we have $f(x) = \hat{f}(f^{-1}(x))$. If $f: X \to Y$, $f': X \to Y$ are correspondences, we say that f is contained in f' if $G_f \subset G_{f'}$. For a subset $A \subset X$ we define the restriction

¹⁾ Another notion of meromorphic mapping and related concepts were defined by W. Stoll [16], [17].

 $f \mid A:A \xrightarrow{k} Y$ by setting $G_{f \mid A} = G_f \cap (A \times Y)$. To every correspondence $f: X \xrightarrow{k} Y$ there is associated the *inverse correspondence* $f^{-1}: Y \xrightarrow{k} X$ whose graph is $G_{f-1} = \{(y,x) \mid (x,y) \in G_f\}$. We have the rule $(f^{-1})^{-1} = f$. The Cartesian product $f \times f_1: X \times X_1 \xrightarrow{k} Y \times Y_1$ of two correspondences $f: X \xrightarrow{k} Y$ and $f_1: X_1 \xrightarrow{k} Y_1$ assigns, by definition, to $(x, x_1) \in X \times X_1$ the set $f(x) \times f_1(x_1)$. If $X = X_1$, we define the junction $(f, f_1): X \xrightarrow{k} Y \times Y_1$ by $(f, f_1)(x) = f(x) \times f_1(x)$. The product $g \circ f: X \xrightarrow{k} Z$ of the correspondences $f: X \xrightarrow{k} Y$ and $g: Y \xrightarrow{k} Z$ is defined by $g \circ f(x) = g(f(x))$. Hence, $(f, f_1) = (f \times f_1) \circ (I_X, I_X)$ where I_X is the identity mapping of X. We have the following rules: $(f \times f_1)^{-1} = f^{-1} \times f_1^{-1}$, $h \circ (g \circ f) = (h \circ g) \circ f$, $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$.

Definition 1. Let X and Y be topological spaces. A correspondence $f: X \to Y$ is continuous at $x \in X$ if

- 1) f(x) is quasicompact, and
- 2) given a neighborhood V of f(x), there exists a neighborhood U of x such that $f(U) \subset V$.

The correspondence f is continuous if it is continuous at every $x \in X$.

Proposition 1. Let $f: X \to Y$ be a correspondence such that f(x) is quasicompact for all $x \in X$. Then f is continuous if and only if f^{-1} is closed (in the sense that the images of closed sets are closed).

Proof. Let f be continuous and let N be a closed set in Y. Assume that $f^{-1}(N)$ is not closed, then there is a point $x \in \overline{f^{-1}(N)} \cap (X-f^{-1}(N))$. We have $f(x) \in Y-N$ since $x \in X-f^{-1}(N)$, hence Y-N is a neighborhood of f(x). Because of the continuity of f there exists a neighborhood f of f such that $f(f) \subset f$ is closed. Let f be a point of f and f and f an open neighborhood of f and f is closed. Let f be a point of f and f an open neighborhood of f because of f and f and f an open neighborhood of f because f is closed and does not contain f therefore f is continuous.

Remark. The statement of Proposition 1 becomes false if "closed" is replaced by "open" as can be seen by simple examples.

¹⁾ This definition and some of the following developments are due to K. Wolffhardt [22].

Proposition 2. If $f: X \to Y$ is a continuous correspondence, f(K) is quasicompact for every quasicompact set $K \subset X$.

Proof. Consider a covering of f(K) by open sets V_i . For each $x \in K$ there are finitely many V_i which cover f(x), let \tilde{V}_x be the union of those V_i . f is continuous at x, hence there is a neighborhood U_x of x such that $f(U_x) \subset \tilde{V}_x$. Now finitely many U_x , say $U_{x_1}, ..., U_{x_n}$, cover K, hence $\tilde{V}_{x_1} \cup ... \cup \tilde{V}_{x_n} \supset f(K)$, therefore finitely many V_i cover f(K).

Proposition 3. Let $f: X \to Y$, $f_1: X_1 \to Y_1$, $f_1': X \to Y_1$, $g: Y \to Z$ be continuous correspondences. Then $f \times f_1$, (f, f_1') , $g \circ f$ are continuous.

Proof. For $g \circ f$, the assertion follows by applying propositions 1 and 2, Furthermore, $f \times f_1(x, x_1) = f(x) \times f_1(x_1)$ is quasicompact for all $x \in X$. $x_1 \in X_1$ since f(x) and $f_1(x_1)$ are quasicompact. Let V be a neighborhood of $f(x) \times f_1(x_1)$; V contains a neighborhood $W \times W_1$ of $f(x) \times f_1(x_1)$ where W, W_1 are neighborhoods of f(x) resp. $f_1(x_1)$. There are neighborhoods U, U_1 of x resp. x_1 such that $f(U) \subset W$, $f_1(U_1) \subset W_1$, then $f \times f_1(U \times U_1) \subset W \times W_1$; hence $f \times f_1$ is continuous. As for (f, f_1) one has $(f, f_1) = (f \times f_1) \circ (I_X, I_X)$, therefore (f, f_1) is continuous because $f \times f_1$ and (I_X, I_X) are continuous.

Proposition 4. A correspondence $f: X \to Y$ is continuous if and only if $f^{-1}: X \to G_f$ is continuous.

Proof. Since $f = \hat{f} \circ f^{-1}$, the continuity of f^{-1} implies that of f by proposition 3. Let f be continuous and let f be a point of f. Since $f^{-1}(f)$ is homeomorphic to f(f), it is quasicompact. Let $f^{-1}(f)$ be open in $f^{-1}(f)$. We can cover $f^{-1}(f)$ by a finite number of sets of the form $f^{-1}(f)$ and $f^{-1}(f)$ open in $f^{-1}(f)$ and there exists a neighborhood $f^{-1}(f)$ of $f^{-1}(f)$ such that $f^{-1}(f)$ If $f^{-1}(f)$ and $f^{-1}(f)$ implies that of $f^{-1}(f)$ by a finite number of $f^{-1}(f)$ be open in $f^{-1}(f)$ and there exists a neighborhood $f^{-1}(f)$ open in $f^{-1}(f)$. If $f^{-1}(f)$ is $f^{-1}(f)$ in $f^{-1}(f)$ in

It follows that a correspondence f is continuous if and only if the projection f is a proper map, that is, f is continuous, closed, and $f^{-1}(x)$ is always quasicompact.

Proposition 5. If $f: X \to Y$ is continuous and Y is a Hausdorff space, G_f is closed in $X \times Y$.

Definition 2. A correspondence f is proper if f and f^{-1} are continuous.¹

Proposition 6. If $f: X_{\xrightarrow{k}} Y$, $f_1: X_{1\xrightarrow{k}} Y_1$, $g: Y_{\xrightarrow{k}} Z$ are proper, then $f \times f_1$ and $g \circ f$ are proper.

The junction of two proper correspondences need not, however, be proper. The diagonal mapping (I_X, I_X) serves as an example if X is not a Hausdorff space. If X is Hausdorff, the junction (f, f') of proper correspondences $f: X \to Y$ and $f': X \to Y'$ remains proper.

Proposition 7. Let $f: X_{\stackrel{\rightarrow}{k}} Y$, $f_1: X_{\stackrel{\rightarrow}{k}} Y_1$, $g: Y_{\stackrel{\rightarrow}{k}} Z$ be continuous where all the spaces are locally compact. Then we have:

- 1) If f is proper, then (f, f_1) and (f_1, f) are proper,
- 2) If $g \circ f$ is proper and g^{-1} surjective, then f is proper,
- 3) If $g \circ f$ is proper and f surjective, then g is proper.

2. Holomorphic correspondences

We consider reduced complex spaces (X, θ) where X is assumed Hausdorff and where the structure sheaf θ has no nilpotent elements. For the definition and related concepts we refer to [8]. The structure sheaf is usually omitted in the notation.

Definition 3. Let X and Y be complex spaces. A correspondence $f: X \xrightarrow{k} Y$ is called holomorphic if

- 1) f is continuous,
- 2) the graph G_f is an analytic set in $X \times Y$.

If only the condition 2) is fulfilled, f is said to be weakly holomorphic.

Let $f: X \to Y$ be weakly holomorphic. Then f^{-1} is weakly holomorphic; furthermore, if $A \subset X$ is analytic in X, $f \mid A$ is weakly holomorphic. Since $f^{-1}(x) = G_f \cap (\{x\} \times Y)$, $x \in X$, is analytic in G_f , $f(x) = \hat{f}(f^{-1}(x))$

¹⁾ Compare [3] where another notion of proper correspondence is defined.