

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 14 (1968)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FLATNESS AND PRIVILEGE
Autor: Douady, A.
Kapitel: I. Flat Morphisms
DOI: <https://doi.org/10.5169/seals-42343>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

FLATNESS AND PRIVILEGE

by A. DOUADY

I. FLAT MORPHISMS

§ 1. Analytic subspaces of an analytic space

Let Y_1 and Y_2 be closed analytic subspaces of an analytic space X , and let them be defined by the \mathcal{O}_X ideals J_1, J_2 .

Definition 1: We say that Y_1 is *analytically included* in Y_2 , and we write $Y_1 \subset Y_2$, when $J_1 \supset J_2$.

Remark: The analytic inclusion implies the set theoretic inclusion, but the converse is not true.

Example: $X = (\mathbf{C}, \mathcal{O}_{\mathbf{C}})$; $J_1 = (x)$, $J_2 = (x^2)$. The space Y_1 is a simple point, Y_2 is a double point, $Y_1 \not\subset Y_2$, while they have the same underlying set.

Definition 2: The subspace $Y_1 \cup Y_2$ is the smallest subspace of X containing Y_1 and Y_2 , and it is defined by $J_1 \cap J_2$. The subspace $Y_1 \cap Y_2$ is the biggest subspace of X contained in both Y_1 and Y_2 , and it is defined by $J_1 + J_2$.

Remark: The underlying set of $Y_1 \cup Y_2$ (Resp. $Y_1 \cap Y_2$) is the union (Resp. intersection) of the underlying sets of Y_1 and Y_2 . However \cup and \cap of analytic spaces do not satisfy the distributivity laws which hold in set-theory: $(Y_1 \cup Y_2) \cap Y_3$ contains $Y_1 \cap Y_3$ and $Y_2 \cap Y_3$, and therefore their union; similarly $(Y_1 \cap Y_2) \cup Y_3 \subset (Y_1 \cup Y_3) \cap (Y_2 \cup Y_3)$. In general the converse inclusions do not hold.

Example: Let $X = \mathbf{C}^2$ and Y_1, Y_2, Z be given by ideals $(x-y)$, $(x+y)$ and (x) respectively.

$(Y_1 \cup Y_2) \cap Z$ is $\{0\}$ provided with $\mathbf{C}\{y\}/(y^2)$, while $(Y_1 \cap Y_2) \cup (Y_2 \cap Z)$ is the reduced space $\{0\}$. On the other hand: $Y_1 \cap Y_2 \subset Z$, $(Y_1 \cap Y_2) \cup Z = Z$, while $(Y_1 \cup Z) \cap (Y_2 \cup Z)$ is the space defined by the ideal (x^2, xy) . Its local ring at the origin is $\mathbf{C}\{x, y\}/(x^2, xy)$ in which x is nilpotent.

Definition 3 : Let X', X be analytic spaces, Y a closed analytic subspace of X defined by J , and $f = (f_0, f^1) : X' \rightarrow X$ a morphism.

The inverse image of Y by $f, f^{-1}(Y)$, is the analytic subspace Y' of X' defined by the ideal $J' = f^1(J) \mathcal{O}_{X'}$.

The inverse image of a simple point x in X is called the f -fiber over x , and is denoted by $f^{-1}(x)$ or $X'(x)$.

Proposition 1 : If $f = (f_0, f^1) : X' \rightarrow X$ is a morphism of analytic spaces, and Y is a subspace of X , then $f^{-1}(Y) \simeq \underset{X}{Y \times X'}$.

Proof : Let T be any analytic space, and $g : T \rightarrow X'$ a morphism. Then g can be considered as a morphism from T to $f^{-1}(Y)$ if and only if $f \circ g$ can be considered as a morphism from T to Y . Thus $f^{-1}(Y)$ and $\underset{X}{X' \times X}$ are solutions of the same universal problem.

§ 2. Analytic pull-back

In the following we want to generalize the notion of inverse image of a subspace.

We shall first recall the basic properties of the tensor product $E \otimes_A F$, where A is a commutative ring and E, F are two A -modules.

(1°) $E \otimes A^n = E^n \quad (n \in \mathbb{N})$

(2°) If the sequence of A -modules $F' \rightarrow F \rightarrow F'' \rightarrow 0$ is exact, then also the sequence $E \otimes F' \rightarrow E \otimes F \rightarrow E \otimes F'' \rightarrow 0$ is exact. (Right exactness of the tensor product)

(3°) If $(F_i)_{i \in I}; f_{ij} : F_j \rightarrow F_i$ is an inductive system, then

$$E \otimes \lim_{\rightarrow} F_i = \lim_{\rightarrow} (E \otimes F_i).$$

On the other hand these properties characterize completely the functor \otimes .

Definition 1 : Let $f = (f_0, f^1) : X' \rightarrow X$ be a morphism of analytic spaces, and \mathcal{E} an \mathcal{O}_X -module. Then $f_0^* \mathcal{E}$ is an $f_0^* \mathcal{O}_X$ -module and $\mathcal{O}_{X'}$ is also an $f_0^* \mathcal{O}_X$ -module (by $f^1 : f_0^* \mathcal{O}_X \rightarrow \mathcal{O}_{X'}$).

The analytic pull-back $f^* \mathcal{E}$ of \mathcal{E} by f is defined by scalar extension:

$$f^* \mathcal{E} = f_0^* \mathcal{E} \otimes_{f_0^* \mathcal{O}_X} \mathcal{O}_{X'}$$

Remark : The inverse image is a particular case of the analytic pull-back.

In fact, if Y is a closed analytic subspace of X and $f : X' \rightarrow X$ is a morphism:

$$f^* \mathcal{O}_Y = f_0^* (\mathcal{O}_X / J_Y) \otimes_{f_0^* \mathcal{O}_X} \mathcal{O}_{X'} \simeq f_0^* \mathcal{O}_X / f_0^* J_Y \otimes_{f_0^* \mathcal{O}_X} \mathcal{O}_{X'} \\ \simeq \mathcal{O}_{X'} / f^1(J_Y). \mathcal{O}_{X'} \simeq \mathcal{O}_{f^{-1}(Y)}$$

(The third isomorphism follows from the fact, that $A/I \otimes_A E \simeq E/IE$).

Elementary properties of the analytic pull-back :

- (a) $(f^* \mathcal{E})_{x'} = (f_0^* \mathcal{E})_{x'} \otimes_{(f_0^* \mathcal{O}_X)_{x'}} \mathcal{O}_{X',x'} \simeq \mathcal{E}_x \otimes_{\mathcal{O}_{X,x}} \mathcal{O}_{X',x'}$ where $x = f_0(x')$ (since \otimes commutes with inductive limits).
- (b) $f^* (\mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{F}) = f^* \mathcal{E} \otimes_{\mathcal{O}_{X'}} f^* \mathcal{F}$, where \mathcal{E} and \mathcal{F} are \mathcal{O}_X -modules.
- (c) If \mathcal{E} is a coherent \mathcal{O}_X -module, then $f^* \mathcal{E}$ is a coherent $\mathcal{O}_{X'}$ -module.

In fact, \mathcal{E} has a locally finite presentation:

$$\mathcal{O}_X^q \rightarrow \mathcal{O}_X^p \rightarrow \mathcal{E} \rightarrow 0, \text{ and } f^* \text{ is compatible with cokernels, } f^* (\mathcal{O}_X^r) = \mathcal{O}_{X'}^r.$$

Special case : The pull-back of vector bundle. Let (E, π) be an analytic

$$\begin{array}{ccc} E \times X' & \xrightarrow{\bar{f}} & E \\ \downarrow \pi' & & \downarrow \pi \\ X' & \xrightarrow{f} & X \end{array}$$

vector bundle over the analytic space X , and $f : X' \rightarrow X$ a morphism of analytic spaces. The fiber product carries a unique structure of vector bundle over X' , such that \bar{f} is a bundle morphism. We call this bundle E' .

Proposition 1 : Let \mathcal{E} (Resp. \mathcal{E}') be the sheaf of analytic sections of E (Resp. E'). Then $\mathcal{E}' = f^* \mathcal{E}$.

Proof (Sketch) : We have a $f_0^* \mathcal{O}_X$ linear morphism $f_0^* \mathcal{E} \rightarrow \mathcal{E}'$, which extends to a morphism $f^* \mathcal{E} \rightarrow \mathcal{E}'$. We can prove that this is an isomorphism. Since the question is local with respect to X' , we can suppose that E is a trivial bundle over X with fiber \mathbf{C}^r , then $\mathcal{E} = \mathcal{O}_X^r$. Also $\mathcal{O}_{X'}^r = f^* \mathcal{O}_X^r$. Therefore $f^* \mathcal{E} = \mathcal{E}'$.

§ 3. Introduction to flatness by examples


Let S be an analytic space. By analytic space over s we mean an analytic space X provided with a morphism $\pi : X \rightarrow S$. Let S be a simple point in S , and consider $X(s) = f^{-1}(s)$.

The main purpose of these lectures is to give a precise meaning to the expression:

“ $X(s)$ depends nicely on s ”, and to give a criterion for the “ nice ” behaviour.

We begin with some examples.


Example 1: X is the closed subspace on \mathbf{C}^2 defined by $(y^2 - x)$, $S = \mathbf{C}$ and $\pi = 1st$ projection.



$$X(s) = \begin{cases} 2 \text{ simple points if } s \neq 0 \\ \text{double point if } s = 0. \end{cases}$$

Here the behaviour of $X(s)$ is nice.

Example 2: X is the closed subspace of \mathbf{C}^2 defined by (xy) , $S = \mathbf{C}$ and $\pi = 1st$ projection.



$X(s)$ is given by $(x-s, xy)$, and

$$(x-s, xy) = \begin{cases} (x-s, y) & \text{if } s \neq 0 \\ (x) & \text{if } s = 0. \end{cases}$$

The first case is a simple point, the second one the y -axis.

A similar example is the map of a point into \mathbf{C} .

In both of these examples the dimension of the fiber makes a jump at one point. We notice, however, that the exceptional point corresponds to an irreducible component of X , and after removing this component π behaves nicely.

This kind of removing is not possible in general, as the following example shows:

Example 3: X is given in \mathbf{C}^3 by $(xz - y)$, and π is the projection on the (x, y) -plane.

If $s = (x_0, y_0)$, then the fiber $X(s)$ is defined by

$$(x-x_0, y-y_0, xz-y) = \begin{cases} \left(x-x_0, y-y_0, z-\frac{y_0}{x_0}\right) & \text{if } x_0 \neq 0 \\ (x, y) & \text{if } x_0 = y_0 = 0 \\ (1) & \text{if } x_0 = 0, y_0 \neq 0. \end{cases}$$

The set of “ nice ” fibers is dense in X , so we cannot remove the z -axis and still get a closed subspace of \mathbf{C}_3 .

§ 4. Algebraic study of flatness

In the following all rings are commutative, with 1, and all modules are unitary.

Definition 1: An A -module E is *flat*, if for every exact sequence of A -modules

$$0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0,$$

the sequence $0 \rightarrow E \otimes F' \rightarrow E \otimes F \rightarrow E \otimes F'' \rightarrow 0$ is also exact. We can also say, because \otimes is right exact, that E is flat, if for every injective homomorphism $F' \rightarrow F$, $E \otimes F' \rightarrow E \otimes F$ is also injective.

Examples of modules which are not flat :

- (1) if $A = \mathbf{Z}$, $E = \mathbf{Z}_2 = \mathbf{Z}/2\mathbf{Z}$, $F = F' = \mathbf{Z}$; then the sequence $0 \rightarrow \mathbf{Z} \xrightarrow{2I} \mathbf{Z} (2I : x' \rightarrow 2x)$ is exact. But now $\mathbf{Z}_2 \otimes \mathbf{Z} = \mathbf{Z}_2$, and the homomorphism $\mathbf{Z}_2 \xrightarrow{2I} \mathbf{Z}_2$ is the zero homomorphism, which is not injective. So \mathbf{Z}_2 is not a flat \mathbf{Z} module.
- (2) If $A = \mathbf{C}\{x\}$, $E = \mathbf{C} = \mathbf{C}\{x\}/(x)$, $F = F' = \mathbf{C}\{x\}$, then the sequence $0 \rightarrow F \xrightarrow{xI} F' (xI : p(x) \rightarrow xp(x))$ is exact. But the homomorphism $E \xrightarrow{xI} E$ is not injective.

Proposition 1 : If A is an integral domain and E a flat A -module, then E is torsion-free.

Proof : Let $a \in A$, $a \neq 0$. Because A is an integral domain, the sequence $0 \rightarrow AA \xrightarrow{aI} (aI : x \rightarrow ax)$ is exact. Since E is flat, the sequence $0 \rightarrow E \xrightarrow{aI} E$ is also exact. In other words E has no torsion elements.

Proposition 2 : If A is a principal-ideal domain, then E is flat if and only if E is torsionfree.

Proof : See corollary of prop. 6.

Examples of flat modules :

- (1) The inductive limit of flat modules is flat, because the inductive limit preserves exactness, and it commutes with the tensor product.

- (2) Every free module is flat. In fact, if E is free and finite type, then $E = A^n$ and $E \otimes F = F^n$. If $F' \rightarrow F$ is injective, so is $F'^n \rightarrow F^n$ too.

If E is an arbitrary free module, then it is an inductive limit of free modules of finite type, and the flatness of E follows from (1).

- (3) Let S be a multiplicative system in A . Then the ring of fractions $S^{-1}A$ is a flat A -module. In fact the ring $S^{-1}A$ can be identified with an inductive limit of free modules, so it is flat ((1) (2)). We assume for simplicity that S has only regular elements. We can define in the set S a partial order in the following way:

$$s' \geq s \Leftrightarrow \exists t \in A, \quad ts = s' \quad (\text{such a } t \text{ is then unique}).$$

Let $E_s = A$ for every $s \in S$, and if $s' \geq s$ (i.e. $s' = ts$) then let $f_s^{s'}$ be the homomorphism $t \cdot I_A : E_s \rightarrow E_{s'}$. The family $(E_s)_{s \in S}$ with the homomorphisms $(f_s^{s'})$ is an inductive system.

Let $E = \lim_{\rightarrow} E_s$ be the inductive limit of this system, and φ_s the canonical homomorphism $E_s \rightarrow E$. We shall define an isomorphism $\psi : E \rightarrow S^{-1}A$.

We first define for every s a homomorphism $\psi_s : E_s = A \rightarrow S^{-1}A$; $x \rightarrow x/s$. Now if $s' \geq s$, then

$$(\psi_{s'} \circ f_s^{s'})(x) = \psi_{s'}(tx) = \frac{tx}{s'} = \frac{tx}{ts} = \frac{x}{s} = \psi_s(x).$$

Therefore there exists a homomorphism $\psi : E \rightarrow S^{-1}A$, satisfying $\psi_s = \psi \circ \varphi_s$ for every $s \in S$.

Because every element of $S^{-1}A$ has the form a/s , ψ is surjective. On the other hand if $\psi(\varphi_s(x)) = 0$, then $\psi_s(x) = x/s = 0$. Thus $x = 0$, and ψ is also injective.

The above proof can be extended to the general case, not assuming that the elements of S are regular. The extended proof involves the notion of inductive limit of an inductive system indexed by a category instead of an ordered set.

From (1) and (2) above, any module which is the inductive limit of free modules, is flat. Conversely:

Theorem 1 : (Daniel, Lazard)

Any flat module is a inductive limit of free modules.

For the proof: See *C.R. Acad. Sci. Paris*, 258 (1964), pp. 6313-6316.

Some elementary properties of flat modules :

- (1) If E and F are flat A -modules, then $E \otimes_A F$ is also flat. In fact, if $G' \rightarrow G$ is injective, then $F \otimes_A G' \rightarrow F \otimes_A G$ is injective, and also $E \otimes_A (F \otimes_A G') \rightarrow E \otimes_A (F \otimes_A G)$ is injective. The result follows from the associativity of the tensor product.
- (2) Let $\phi : A \rightarrow B$ be a ring homomorphism, and E a flat A -module. The module $B \otimes_A E$ is a flat B -module.

If F is a B -module, then $F \otimes_B (B \otimes_A E) = (F \otimes_B B) \otimes_A E = F \otimes_A E$ further if F' and F are B -modules, and $F' \rightarrow F$ an injective homomorphism of B -modules, we can consider this homomorphism as an injective homomorphism of A -modules. Because E is A -flat,

$$F' \otimes_A E \rightarrow F \otimes_A E \text{ is injective.}$$

- (3) Let $\phi : A \rightarrow B$ be a ring homomorphism, such that B is a flat A -module. If F is a flat B -module, then F is a flat A -module. In fact: if $E' \rightarrow E$ is injective, then $E' \otimes_A B \rightarrow E \otimes_A B$ is injective, and also $(E \otimes_A B) \otimes_B F' \rightarrow (E \otimes_A B) \otimes_B F$ is injective. But $(E' \otimes_A B) \otimes_B F' = E' \otimes_A F$; $(E \otimes_A B) \otimes_B F = E \otimes_A F$.

If an A -module E is not flat, we want to measure how far it is from being flat. For this purpose we introduce the functor Tor .

Definition 2 : A free resolution of E is an exact sequence: $\dots \rightarrow L_n \rightarrow L_{n-1} \rightarrow \dots \rightarrow L_1 \rightarrow L_0 \rightarrow E \rightarrow 0$, where all L_i are free A -modules.

The complex of the resolution is the sequence

$$(L.) \dots \rightarrow L_n \rightarrow L_{n-1} \rightarrow \dots \rightarrow L_1 \rightarrow L_0 \rightarrow 0.$$

Every module has a free resolution. Two resolutions are algebraically homotopy-equivalent. Forming the tensor products $L_i \otimes F$, we get

$$(L. \otimes F) \dots \rightarrow L_n \otimes F \rightarrow L_{n-1} \otimes F \rightarrow \dots \rightarrow L_1 \otimes F \rightarrow L_0 \otimes F \rightarrow 0.$$

Definition 3 :

$$\text{Tor}_n^A(E, F) = H_n(L. \otimes F) = \frac{\text{Ker}(L_n \otimes F \rightarrow L_{n-1} \otimes F)}{\text{Im}(L_{n+1} \otimes F \rightarrow L_n \otimes F)}$$

if $n \geq 1$, and $\text{Tor}_0^A(E, F) = \text{Coker}(L_1 \otimes F \rightarrow L_0 \otimes F) = E \otimes F$.

Basic properties of Tor :

- (1) $\text{Tor}_n(E, F)$ is independent of the choice of the resolution (up to a canonical isomorphism).

- (2) If we take a free resolution of F , we get $\text{Tor}_n(F, E) = \text{Tor}_n(E, F)$ (Symmetry of the Tor). We can also define $\text{Tor}_n(E, F)$ by taking two free resolutions, one of E and one of F .
- (3) If $0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$ is a short exact sequence, then we get a long exact sequence:

$$\begin{array}{ccccccc} \text{Tor}_n(E', F) & \rightarrow & \text{Tor}_n(E, F) & \rightarrow & \text{Tor}_n(E'', F) & \rightarrow & \\ \rightarrow & \text{Tor}_{n-1}(E', F) & \rightarrow & \text{Tor}_{n-1}(E, F) & \rightarrow & \text{Tor}_{n-1}(E'', F) & \rightarrow \\ \rightarrow & \text{---} & \rightarrow & \text{---} & \rightarrow & \text{---} & \rightarrow \\ \rightarrow & \text{Tor}_1(E', F) & \rightarrow & \text{Tor}_1(E, F) & \rightarrow & \text{Tor}_1(E'', F) & \rightarrow \\ \rightarrow & E' \otimes F & \rightarrow & E \otimes F & \rightarrow & E'' \otimes F & \rightarrow 0. \end{array}$$

- (4) Tor is compatible with inductive limit, i.e. if $E = \lim (E_i)$, then
- $$\text{Tor}_n(\lim E_i, F) = \lim (\text{Tor}_n(E_i, F)).$$

- (5) We can define $\text{Tor}_n(E, F)$ by taking a flat resolution of E .

Proposition 3: Let E be an A -module. Then the following conditions are equivalent:

- (a) E is flat.
 (b) For all A -modules F , and for all $n \geq 1$, $\text{Tor}_n(E, F) = 0$.
 (c) For all A -modules F , $\text{Tor}_1(E, F) = 0$.

Proof: (a) \Rightarrow (b). If $\dots \rightarrow L_n \rightarrow L_{n-1} \rightarrow \dots \rightarrow L_1 \rightarrow L_0 \rightarrow F \rightarrow 0$ is a free resolution of F , then the sequence

$$\dots \rightarrow E \otimes L_n \rightarrow E \otimes L_{n-1} \rightarrow \dots \rightarrow E \otimes L_1 \rightarrow E \otimes L_0 \rightarrow E \otimes F \rightarrow 0$$

is exact, thus $\text{Tor}_n(E, F) = 0$ for all $n \geq 1$.

(b) \Rightarrow (c) clear. (c) \Rightarrow (a): If the sequence $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$ is exact, so is also (by (3) above) $\text{Tor}_1(E, F'') \rightarrow E \otimes F' \rightarrow E \otimes F \rightarrow E \otimes F'' \rightarrow 0$. Now $\text{Tor}_1(E, F'') = 0$, thus E is flat.

Proposition 4: If I and J are two ideals in A , then $\text{Tor}_1^A(A/I, A/J) = I \cap J / I \cdot J$.

Proof: From the exact sequence $0 \rightarrow I \rightarrow A \rightarrow A/I \rightarrow 0$, we get the exact sequence:

$$\text{Tor}_1(A, A/J) \rightarrow \text{Tor}_1(A/I, A/J) \rightarrow I \otimes A/J \rightarrow A \otimes A/J \rightarrow A/I \otimes A/J \rightarrow 0.$$

But now $\text{Tor}_1(A, A/J) = 0$ (A being A -free), and $I \otimes A/J = I/I \cdot J$; $A \otimes A/J = A/J$. Therefore the sequence $0 \rightarrow \text{Tor}_1(A/I, A/J) \rightarrow I/I \cdot J \rightarrow A/J$ is exact, and $\text{Tor}_1(A/I, A/J) = \text{Ker}(I/I \cdot J \rightarrow A/J) = I \cap J / I \cdot J$.

Example: Let U be an open set in \mathbf{C}^n , and $x \in U$. Further let $X, Y \subset U$ be two hypersurfaces, defined by $I = (f)$ and $J = (g)$. Supposing that f and g do not have common factors: $I_x \cap J_x = I_x J_x$, and

$$\text{Tor}_1(\mathcal{O}_{X,x}, \mathcal{O}_{Y,x}) = \text{Tor}_1(\mathcal{O}_{U,x}/I_x, \mathcal{O}_{U,x}/J_x) = \frac{I_x \cap J_x}{I_x \cdot J_x} = 0.$$

Heuristic remark: The formula $\text{Tor}_1(\mathcal{O}_{X,x}, \mathcal{O}_{Y,x}) = 0$ expresses the fact that X and Y are “in general position”. If for example X and Y are two linear subspaces in \mathbf{C}^n of dimensions p and q , we have $\text{Tor}_1(\mathcal{O}_{X,x}, \mathcal{O}_{Y,x}) = 0$ if $\dim(X \cap Y) = p + q - n$, and $\text{Tor}_1(\mathcal{O}_{X,x}, \mathcal{O}_{Y,x}) \neq 0$ otherwise.

Next we shall prove an elementary flatness criterion.

Proposition 5: Let E be an A -module. The following conditions are equivalent:

- (a) E is flat.
- (b) For all finitely generated ideals I of A , $\text{Tor}_1(E, A/I) = 0$.
- (c) For all monogenous A -modules F , $\text{Tor}_1(E, F) = 0$.

Proof: (a) \Rightarrow (b), by prop. 3.

(b) \Rightarrow (c): Because Tor is compatible with inductive limit, we can suppose, that $\text{Tor}_1(E, A/I) = 0$ for an arbitrary ideal I of A . But every monogenous A -module F can be represented by A/I .

(c) \Rightarrow (a). By prop. 3 it is sufficient to prove that $\text{Tor}_1(E, F) = 0$ for any A -module F .

First consider the case, where F is finitely generated. We use induction, supposing that $\text{Tor}_1(E, F) = 0$, when F has n generators. Let F have $(n+1)$ generators x_1, \dots, x_n, x_{n+1} . If F' is the submodule generated by $\{x_1, \dots, x_n\}$, then $F' \subset F$ and $F'' = F/F'$ is monogenous. The exact sequence $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$ gives the exact sequence $\text{Tor}_1(E, F') \rightarrow \text{Tor}_1(E, F) \rightarrow \text{Tor}_1(E, F'')$. Now $\text{Tor}_1(E, F') = \text{Tor}_1(E, F'') = 0$, thus $\text{Tor}_1(E, F) = 0$. In the general case, F can be considered as an inductive limit of finitely generated modules, and because Tor is compatible with inductive limits, $\text{Tor}_1(E, F) = 0$.

Proposition 6: Let A be an integral domain, and E an A -module. Then E is torsionfree if and only if $\text{Tor}_1(E, A/(a)) = 0$, for any element $a \in A$.

Proof: If E is A -module, $a \in A$, then the exact sequence $0 \rightarrow A \xrightarrow{aI} A \rightarrow A/(a) \rightarrow 0$ gives the exact sequence $0 \rightarrow \text{Tor}_1(E, A/(a)) \rightarrow E \xrightarrow{aI} E$. In other words $\text{Tor}_1(E, A/(a)) = \{x \in E \mid ax = 0\}$, from which the result follows.

Corollary: Let A be a principal ideal domain. E is flat if and only if E is torsionfree.

Proof: We have already proved that, if E is flat, then it is torsion free. The converse follows from prop. 6 and prop. 5.

The first flatness criterion for noetherian local rings is the following:

Theorem 2: Let A be a noetherian local ring with maximal ideal m ; $k = A/m$, and E a finitely generated A -module. The following conditions are equivalent:

- (a) E is free.
- (b) E is flat.
- (c) $\text{Tor}_1^A(E, k) = 0$.

Proof: We have already proved $(a) \Rightarrow (b) \Rightarrow (c)$.

$(c) \Rightarrow (a)$: We recall first Nakayma's lemma. If A is a local ring with maximal ideal m ; $k = A/m$, and E is a finitely generated A -module, such that $k \otimes_A E = E/mE = 0$, then $E = 0$.

The module $\bar{E} = k \otimes_A E = E/mE$ is a finitely generated vector space over k . Let $\{\bar{x}_1, \dots, \bar{x}_r\}$ be a base of \bar{E} (over k), and $\{x_1, \dots, x_r\}$ E representatives of \bar{x}_i : s . Consider the homomorphism $\phi : A^r \rightarrow E$, $\phi(a_1, \dots, a_r) = \sum a_i x_i$. Denoting by R and Q the kernel and the cokernel of ϕ , we get an exact sequence:

$$(*) \quad 0 \rightarrow R \rightarrow A^r \rightarrow E \rightarrow Q \rightarrow 0$$

and R, Q are finitely generated A -modules. From $(*)$ we get the exact sequence

$$A^r \otimes_A k \rightarrow E \otimes_A k \rightarrow Q \otimes_A k \rightarrow 0.$$

But $\bar{E} = E \otimes_A k \simeq k^r = A^r \otimes_A k$, so $Q \otimes_A k = 0$, and by Nakayama's lemma $Q = 0$.

Therefore we have an exact sequence

$$0 \rightarrow R \rightarrow A^r \rightarrow E \rightarrow 0.$$

From this we get: $\text{Tor}_1(E, k) \rightarrow k \otimes_A R \rightarrow k^r \rightarrow \bar{E} \rightarrow 0$ (exact). Now: $\bar{E} \simeq k^r$, $\text{Tor}_1(E, k) = 0$ (by assumption). Therefore $k \otimes_A R = 0$, and once more by Nakayama's lemma $R = 0$, thus $E \simeq A^r$, i.e. E is free.

Proposition 7: Let $\phi : A \rightarrow B$ be a ring homomorphism, and let B be A -flat. If I is an ideal of A , we write $\bar{A} = A/I$, $\bar{B} = B/IB = \bar{A} \otimes_A B$. Let F be a B -module, then: $\text{Tor}_i^A(\bar{A}, F) = \text{Tor}_i^B(\bar{B}, F)$ ($i \geq 0$).

Proof: We choose first a B -free resolution of F

$$\rightarrow L_{n+1} \rightarrow L_n \rightarrow \dots \rightarrow L_1 \rightarrow L_0 \rightarrow F \rightarrow 0.$$

If $L.$ is the respective complex of resolution, then

$$\bar{B} \otimes_B L. = B/IB \otimes_B L. = \bar{A} \otimes_A (B \otimes_B L.) = \bar{A} \otimes_A L.$$

Because every L_i is B -free, and B is A -flat, every L_i is A -flat (Property 3 after Th. 1). Thus $L.$ is a flat A -resolution, and

$$\text{Tor}_i^A(\bar{A}, F) = H_i(\bar{A} \otimes_A L.) = H_i(\bar{B} \otimes_B L.) = \text{Tor}_i^B(\bar{B}, F).$$

We shall next state the second flatness criterion for noetherian local rings.

Theorem 3: Let A and B be two noetherian local rings, with maximal ideals $\underline{m}, \underline{n}; k = A/\underline{m}$. If $\phi : A \rightarrow B$ is a local homomorphism (i.e. $\phi(\underline{m}) \subset \underline{n}$), and F finitely generated B module then

$$F \text{ is } A\text{-flat} \Leftrightarrow \text{Tor}_1^A(k, F) = 0.$$

The proof of this theorem is much more difficult than that of th. 20 see for example:

Bourbaki: *Algèbre commutative*, Chapter III § 5, th1, (i) \Leftrightarrow (iii), p. 98.

The conditions in Bourbaki's theorem are here fulfilled:

- 1° A finitely generated module F over a noetherian local ring B is idealwise separated for \underline{n} . (*Ibid.*, § 5. 1. Ex. 1, p. 97.)
- 2° If $\phi : A \rightarrow B$ is a local homomorphism, F is also idealwise separated for \underline{m} . (*Ibid.*, § 5, prop. 2, p. 101.)
- 3° Also the flatness condition is fulfilled, because k is a field.

Remark: The main interest of the theorem lies in the fact, that it is true without any assumption of finiteness on B .

Corollary: If the assumptions are the same as in the theorem 3, and if moreover B is A -flat, then

$$F \text{ is } A\text{-flat} \Leftrightarrow \text{Tor}_1^B(\bar{B}, F) = 0,$$

where $\bar{B} = B/\underline{m}B$.

Proof: $\text{Tor}_1^A(k, F) = \text{Tor}_1^B(\bar{B}, F)$, by prop. 7.

§ 5. *Geometric applications of the flatness criterions*

A) *Flatness for finite morphisms*

Proposition 1: Let $\pi: X \rightarrow S$ be a finite morphism (i.e. proper with finite fibres) of analytic spaces. Then $\pi_*(\mathcal{O}_X)$ is a coherent analytic sheaf over S . The following conditions are equivalent:

- (a) π is flat (i.e. for every $x \in X$, $\mathcal{O}_{X,x}$ is a flat $\mathcal{O}_{S,s}$ -module, $s = \pi(x)$).
- (b) For every s , $(\pi_* \mathcal{O}_X)_s$ is a flat $\mathcal{O}_{S,s}$ -module.
- (c) $\pi_* \mathcal{O}_X$ is a locally free sheaf.

Proof: Because π is finite $\pi_*(\mathcal{O}_X)_s = \bigoplus_{x \in \pi^{-1}(s)} \mathcal{O}_{X,x}$, thus the only point to prove is (b) \Rightarrow (c).

Now if $\mathcal{O}_{X,x}$ is a flat $\mathcal{O}_{S,s}$ -module, then (by theorem 2) $\mathcal{O}_{X,x}$ is free, and a coherent sheaf whose fibers are free is a locally free sheaf.

Proposition 2: Let S be a reduced analytic space and \mathcal{E} a coherent \mathcal{O}_S -module. Let $E(s)$ be the finite dimensional vector space (over \mathbb{C}) $\mathcal{E}_s \otimes_{\mathcal{O}_{S,s}} \mathbb{C}_s$. \mathcal{E} is a locally free $\mathcal{O}_{S,s}$ -module if and only if $\dim_{\mathbb{C}} E(s)$ is locally constant.

Proof: If \mathcal{E} is locally free, then $\dim_{\mathbb{C}} E(s)$ is locally constant. Suppose now that $\dim_{\mathbb{C}} E(s)$ is locally constant in an open set $U \subset S$, and that $\mathcal{O}_U^p \xrightarrow{d} \mathcal{O}_U^q \rightarrow \mathcal{E}_U \rightarrow 0$ is exact. d is determined by a $p \times q$ matrix of analytic functions on U , so it gives a morphism $\mathbb{C}_U^p \xrightarrow{d} \mathbb{C}_U^q$ of trivial vector bundles over U .

From the exact sequence $\mathcal{O}_S^p \xrightarrow{d_s} \mathcal{O}_S^q \rightarrow \mathcal{E}_S \rightarrow 0$, we get (by making tensor-products with \mathbb{C}_s) the exact sequence:

$$\mathbb{C}_s^p \xrightarrow{d(s)} \mathbb{C}_s^q \rightarrow E(s) \rightarrow 0,$$

which shows that d has constant rank in U . Thus $\text{Ker } d$ and $\text{Im } d$ are vector bundles, and we can write

$$\mathbb{C}_U^p = F_1 \oplus G_1, \quad \mathbb{C}_U^q = F_0 \oplus G_0,$$

$$d : \begin{cases} F_1 \rightarrow 0 \\ G_1 \simeq F_0. \end{cases}$$

Now $\mathcal{E} \simeq$ the sheaf of analytic sections of G_0 , therefore \mathcal{E} is locally free.

Definition 1: Let $\pi : X \rightarrow S$ be a finite morphism of analytic spaces, and $s \in S$. For each $x \in X(s) = \pi^{-1}(s)$, $\mathcal{O}_{X(s),x} = \mathbf{C} \otimes_{\mathcal{O}_{S,s}} \mathcal{O}_{X,x}$ is finite dimensional vectorspace over \mathbf{C} . Denote its dimension by $v(x)$. Then the degree $v(s)$ of s is defined by $v(s) = \sum_{x \in X(s)} v(x)$.

Theorem 1: Let $\pi : X \rightarrow S$ be a finite morphism of analytic space and let S be a reduced space. Then X is flat over S if and only if $v(s)$ is locally constant function of s .

$$\begin{aligned} \text{Proof: } v(s) &= \sum_{x \in X(s)} \dim_{\mathbf{C}} \mathcal{O}_{X(s),x} = \dim_{\mathbf{C}} \left(\bigoplus_{x \in X(s)} \mathcal{O}_{X(s),x} \right) \\ &= \dim_{\mathbf{C}} \left(\bigoplus_{x \in X(s)} (\mathbf{C} \otimes_{\mathcal{O}_{S,s}} \mathcal{O}_{X,x}) \right) \\ &= \dim_{\mathbf{C}} \mathbf{C} \otimes_{\mathcal{O}_{S,s}} \pi_* (\mathcal{O}_X)_s = \dim_{\mathbf{C}} E(s). \end{aligned}$$

The theorem follows from propositions 1 and 2.

Examples of flat morphisms

Example 1: If $\pi : X \rightarrow S$ is a local isomorphism near x , then π is flat at x .

Example 2: Consider § 2, Ex. 1. Here $v(x) = 1$.

Examples of non-flat morphisms

Examples 1: If $X \subset S$ is a closed subspace, not open, $v(s)$ is not locally constant.

Example 2: Let X be a subspace of \mathbf{C}^4 defined by the ideal intersection of (x_3, x_4) and $(x_1 - x_1, x_4 - x_2)$ (which is equal to the product ideal) and let π be the projection onto the (x_1, x_2) -plane \mathbf{C}^2 . Then X is a union of two 2-planes in \mathbf{C}^4 , whose intersection is (0) . When $s \neq 0$, $X(s)$ consists of two simple points, so $v(s) = 2$. $X(0)$ is given by the ideal $(x_1, x_2, x_3^2, x_3x_4, x_4^2)$, thus $v(0) = 3$.

Example 3: Let $S = \{(u, v, w) \in \mathbf{C}^3 \mid v^2 = uw\}$ and $\pi : \mathbf{C}^2 \rightarrow S$ be the map $(x, y) \rightarrow (x^2, xy, y^2)$. This map identifies S with the quotient of \mathbf{C}^2 by the equivalence relation identifying (x, y) with $(-x, -y)$. However, π is not flat, since for $s \in S$, $v(s) = 2$ if $s \neq 0$ and $v(s) = 3$ if $s = 0$.

B) *Projection of a product of analytic spaces*

Theorem 2: Let S and X be analytic spaces. If $\pi : S \times X \rightarrow S$ is the projection morphism, then π is flat, i.e. $\mathcal{O}_{S \times X, (s,x)}$ is a flat $\mathcal{O}_{S,s}$ module for every $(s, x) \in S \times X$.

To prove this theorem we need first some homological algebra. Then we shall show it in the particular case, when S is a manifold, and finally in the general case.

(a) *Koszul complex*

Let A be a ring, M an A -module and h_1, \dots, h_n homomorphisms $M \rightarrow M$, which commute with each other, i.e. $h_i h_j = h_j h_i$ for every i, j .

If $1 \leq k \leq n$, set $Q_k = M/h_1(M) + \dots + h_k(M)$, and $Q_0 = M$, thus, in particular, $Q_n = Q = M / \sum_{i=1}^n h_i(M)$. Every h_k induces a map $\tilde{h}_k : Q_{k-1} \rightarrow Q_{k-1}$.

Definition 2: The sequence (h_1, \dots, h_n) is called regular if each of the mappings \tilde{h}_k ($1 \leq k \leq n$) is injective.

The Koszul complex of the module M and of the mappings h_k ($1 \leq k \leq n$) $K. = K. [M; h_1, \dots, h_n]$ is defined in the following way:

$$K_i = \wedge^{n+i} A^n \otimes M \simeq M^{\binom{n}{i}}, \quad 0 \leq i \leq n.$$

We define the homomorphisms $d_i : K_i \rightarrow K_{i-1}$ ($i > 0$) by $\lambda \otimes x \rightarrow \sum_i (e_i \wedge \lambda) \otimes \otimes h_i(x)$, where (e_i) is the natural base of A^n . We also define $\varepsilon : K_0 \rightarrow Q$ as the natural map $: K_0 = M \rightarrow M / \sum_{i=1}^n h_i(M) = Q$. Using the fact that h_1, \dots, h_n commute with each other, it is easy to verify that

$$(d_{i-1} \circ d_i)(\lambda \otimes x) = \sum_{i,j} (e_j \wedge e_i \wedge \lambda) \otimes h_j(h_i(x)) = 0;$$

also $\varepsilon d_1 = 0$. Thus $K.$ is really a complex.

Theorem 3 (Poincaré-Koszul).

If (h_1, \dots, h_n) is a regular sequence, then

$$H_i(K.) = \begin{cases} Q & \text{if } i = 0 \\ \cdot & \cdot \\ 0 & \text{if } i > 0 \end{cases}$$

If $h_i \in A$, it defines the map: $A \xrightarrow{h_i I} A$, which we denote also by h_i . We say that (h_1, \dots, h_n) is a regular sequence of elements if $(h_1 I, \dots, h_n I)$ is a regular sequence.

Corollary. If (h_1, \dots, h_n) is a regular sequence of elements, then the Koszul complex $K. = K. [A; h_1, \dots, h_n] = \{ \wedge^{n-1} A^n \simeq A^{(n)} \}$ is a free resolution of $Q = A/(h_i)$ ((h_i) is the ideal generated by h_1, \dots, h_n).

Example: If $A = \mathbf{C} \{x_1, \dots, x_n\}$; $h_i = x_i$, then $Q_k = A/(x_1, \dots, x_k) = \mathbf{C} \{x_{k+1}, \dots, x_n\}$ and $Q = Q_n = \mathbf{C}$. The complex $K. = K. [A; x_1, \dots, x_n]$ is a free resolution of \mathbf{C} .

(b) *Proof of theorem 2, when S is a complex manifold*

In this case we can take $\mathcal{O}_{S,S} = \mathbf{C} \{t_1, \dots, t_m\} = A$ and if $\mathcal{O}_{X,x} = \mathbf{C} \{x_1, \dots, x_n\}/(f_1, \dots, f_p)$, then

$$\mathcal{O}_{S \times X, (s,x)} = \mathbf{C} \{t_1, \dots, t_m, x_1, \dots, x_n\}/(f_1, \dots, f_p) = B.$$

B is an A -module in a natural way.

By the corollary of the Poincaré-Koszul theorem $K. = K. [A; t_1, \dots, t_m]$ in a free resolution of \mathbf{C} . We want to compute the modules $\text{Tor}_i^A(\mathbf{C}, B) = H_i(K. \otimes B)$ ($i > 0$).

It's easily seen, that we can consider the complex $K. \otimes B$ as a Koszul

complex $K'. = K. [B; t_1, \dots, t_m]$ (where $t_i : B \xrightarrow{t_i I} B$). But now the sequence (t_1, \dots, t_m) is regular, thus by the Poincaré-Koszul theorem $H_i[K'] = 0$ if $i > 0$.

In particular: $\text{Tor}_1^A(\mathbf{C}, B) = H_1[K. \otimes B] = H_1[K'] = 0$. By the second flatness criterion B is A -flat.

(c) *The general case*

The question being local, we can suppose that $S \subset W \subset \mathbf{C}^n$, where W is open, and S an analytic subspace of W . Let S be defined by g_1, \dots, g_r . Then $S \times X \subset W \times X$ and $\mathcal{O}_S = \mathcal{O}_W/(g_1, \dots, g_r)$. On the other hand $\mathcal{O}_{S \times X} = \mathcal{O}_{W \times X}/(g_1, \dots, g_r) = \mathcal{O}_S \otimes_{\mathcal{O}_W} \mathcal{O}_{W \times X}$. The last equality follows from

the fact, that if $\pi : X \rightarrow S$ is a morphism, and $S' \subset S$ a subspace, $X' = \pi^{-1}(S')$,

$$\text{then } \mathcal{O}_{X'} = \mathcal{O}_{S'} \otimes_{\mathcal{O}_S} \mathcal{O}_X.$$

Remark : This a particular case of the following proposition: if π and π' are two morphisms of which at least one is finite, then

$$\begin{array}{ccc} X & & Y \\ \pi \searrow & & \swarrow \pi' \\ & S & \end{array} \quad \mathcal{O}_{X \times_S Y} = \mathcal{O}_X \otimes_{\mathcal{O}_S} \mathcal{O}_Y.$$

We have proved that $\mathcal{O}_{W \times X}$ is \mathcal{O}_W -flat, so by scalar extension $\mathcal{O}_{S \times X}$ is \mathcal{O}_S flat.

Corollary : If X and S are two manifolds and $\pi : X \rightarrow S$ is a submersion, then π is flat.

III. PRIVILEGED POLYCYLINDERS

§ 1. Banach vector bundles over an analytic space

Let E be a Banach space and X an analytic space. We denote then by E_X the trivial bundle $X \times E$ over X .

To define bundle morphisms, we first define the sheaf $\mathcal{H}_X(E)$ of germs of analytic morphisms from X to E . If $U \subset \mathbb{C}^n$ is open, then the set $\mathcal{H}(U, E)$ of analytic morphisms from U into E consists of all functions $g : U \rightarrow E$ having at every point $x \in U$ a converging power series expansion.

Let now X' be a local model for X , i.e. X' is the support of the quotient sheaf \mathcal{O}_U/J , where $U \subset \mathbb{C}^n$ is open and J is a coherent sheaf of ideals of \mathcal{O}_U , then $\mathcal{H}_{X'}(E)$ is the sheaf associated to the presheaf $V \rightarrow \mathcal{H}(V, E)/J_V \cdot \mathcal{H}(V, E)$ ($V \subset U$, V -open).

Remark : If X' is reduced, the sections of $\mathcal{H}_{X'}(E)$ are just the functions from X' to E which are locally induced by analytic functions on open sets in U .

The sheaf $\mathcal{H}_X(E)$ is constructed with help of the local models X' of X , i.e. $\mathcal{H}_X(E)|_{X'} = \mathcal{H}_{X'}(E)$, for every local model X' .

Definition 1 : The set of analytic morphisms from an analytic space X into a Banach space E is the set $\mathcal{H}(X; E)$ of sections of the sheaf $\mathcal{H}_X(E)$.

Let $\mathcal{L}(E, F)$ be the Banach space of all continuous linear mappings from the Banach space E into the Banach space F .

Definition 2 : An analytic vector bundle morphism from E_X into F_X is an analytic morphism from X into $\mathcal{L}(E, F)$.