

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 14 (1968)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FLATNESS AND PRIVILEGE
Autor: Douady, A.
Kapitel: §1. Analytic subspaces of an analytic space
DOI: <https://doi.org/10.5169/seals-42343>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

FLATNESS AND PRIVILEGE

by A. DOUADY

I. FLAT MORPHISMS

§ 1. Analytic subspaces of an analytic space

Let Y_1 and Y_2 be closed analytic subspaces of an analytic space X , and let them be defined by the \mathcal{O}_X ideals J_1, J_2 .

Definition 1: We say that Y_1 is *analytically included* in Y_2 , and we write $Y_1 \subset Y_2$, when $J_1 \supset J_2$.

Remark: The analytic inclusion implies the set theoretic inclusion, but the converse is not true.

Example: $X = (\mathbf{C}, \mathcal{O}_{\mathbf{C}})$; $J_1 = (x)$, $J_2 = (x^2)$. The space Y_1 is a simple point, Y_2 is a double point, $Y_1 \not\subset Y_2$, while they have the same underlying set.

Definition 2: The subspace $Y_1 \cup Y_2$ is the smallest subspace of X containing Y_1 and Y_2 , and it is defined by $J_1 \cap J_2$. The subspace $Y_1 \cap Y_2$ is the biggest subspace of X contained in both Y_1 and Y_2 , and it is defined by $J_1 + J_2$.

Remark: The underlying set of $Y_1 \cup Y_2$ (Resp. $Y_1 \cap Y_2$) is the union (Resp. intersection) of the underlying sets of Y_1 and Y_2 . However \cup and \cap of analytic spaces do not satisfy the distributivity laws which hold in set-theory: $(Y_1 \cup Y_2) \cap Y_3$ contains $Y_1 \cap Y_3$ and $Y_2 \cap Y_3$, and therefore their union; similarly $(Y_1 \cap Y_2) \cup Y_3 \subset (Y_1 \cup Y_3) \cap (Y_2 \cup Y_3)$. In general the converse inclusions do not hold.

Example: Let $X = \mathbf{C}^2$ and Y_1, Y_2, Z be given by ideals $(x-y)$, $(x+y)$ and (x) respectively.

$(Y_1 \cup Y_2) \cap Z$ is $\{0\}$ provided with $\mathbf{C}\{y\}/(y^2)$, while $(Y_1 \cap Y_2) \cup (Y_2 \cap Z)$ is the reduced space $\{0\}$. On the other hand: $Y_1 \cap Y_2 \subset Z$, $(Y_1 \cap Y_2) \cup Z = Z$, while $(Y_1 \cup Z) \cap (Y_2 \cup Z)$ is the space defined by the ideal (x^2, xy) . Its local ring at the origin is $\mathbf{C}\{x, y\}/(x^2, xy)$ in which x is nilpotent.

Definition 3 : Let X', X be analytic spaces, Y a closed analytic subspace of X defined by J , and $f = (f_0, f^1) : X' \rightarrow X$ a morphism.

The inverse image of Y by $f, f^{-1}(Y)$, is the analytic subspace Y' of X' defined by the ideal $J' = f^1(J) \mathcal{O}_{X'}$.

The inverse image of a simple point x in X is called the f -fiber over x , and is denoted by $f^{-1}(x)$ or $X'(x)$.

Proposition 1 : If $f = (f_0, f^1) : X' \rightarrow X$ is a morphism of analytic spaces, and Y is a subspace of X , then $f^{-1}(Y) \simeq \underset{X}{Y \times X'}$.

Proof : Let T be any analytic space, and $g : T \rightarrow X'$ a morphism. Then g can be considered as a morphism from T to $f^{-1}(Y)$ if and only if $f \circ g$ can be considered as a morphism from T to Y . Thus $f^{-1}(Y)$ and $\underset{X}{X' \times X}$ are solutions of the same universal problem.

§ 2. Analytic pull-back

In the following we want to generalize the notion of inverse image of a subspace.

We shall first recall the basic properties of the tensor product $E \otimes_A F$, where A is a commutative ring and E, F are two A -modules.

(1°) $E \otimes A^n = E^n$ ($n \in \mathbb{N}$)

(2°) If the sequence of A -modules $F' \rightarrow F \rightarrow F'' \rightarrow 0$ is exact, then also the sequence $E \otimes F' \rightarrow E \otimes F \rightarrow E \otimes F'' \rightarrow 0$ is exact. (Right exactness of the tensor product)

(3°) If $(F_i)_{i \in I}; f_{ij} : F_j \rightarrow F_i$ is an inductive system, then

$$E \otimes \lim_{\rightarrow} F_i = \lim_{\rightarrow} (E \otimes F_i).$$

On the other hand these properties characterize completely the functor \otimes .

Definition 1 : Let $f = (f_0, f^1) : X' \rightarrow X$ be a morphism of analytic spaces, and \mathcal{E} an \mathcal{O}_X -module. Then $f_0^* \mathcal{E}$ is an $f_0^* \mathcal{O}_X$ -module and $\mathcal{O}_{X'}$ is also an $f_0^* \mathcal{O}_X$ -module (by $f^1 : f_0^* \mathcal{O}_X \rightarrow \mathcal{O}_{X'}$).

The analytic pull-back $f^* \mathcal{E}$ of \mathcal{E} by f is defined by scalar extension:

$$f^* \mathcal{E} = f_0^* \mathcal{E} \otimes_{f_0^* \mathcal{O}_X} \mathcal{O}_{X'}$$