Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 14 (1968)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FLATNESS AND PRIVILEGE
Autor: Douady, A.
Kapitel: §2. Analytic pull-back
DOI: https://doi.org/10.5169/seals-42343

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Definition 3 : Let X', X be analytic spaces, Y a closed analytic subspace of X defined by J, and $f = (f_0, f^1) : X' \to X$ a morphism.

The inverse image of Y by f, $f^{-1}(Y)$, is the analytic subspace Y' of X' defined by the ideal $J' = f^1(J)\,\mathcal{O}_{X'}$.

The inverse image of a simple point x in X is called the f-fiber over x, and is denoted by $f^{-1}(x)$ or $X'(x)$.

Proposition 1 : If $f = (f_0, f^1) : X' \to X$ is a morphism of analytic spaces, and Y is a subspace of X, then $f^{-1}(Y) \simeq Y \underset{X}{\times} X'$.

Proof : Let T be any analytic space, and $g : T \to X'$ a morphism. Then g can be considered as a morphism from T to $f^{-1}(Y)$ if and only if $f \circ g$ can be considered as a morphism from T to Y. Thus $f^{-1}(Y)$ and $X' \underset{X}{\times} X$ are solutions of the same universal problem.

§ 2. *Analytic pull-back*

In the following we want to generalize the notion of inverse image of a subspace.

We shall first recall the basic properties of the tensor product $E \underset{A}{\otimes} F$, where A is a commutative ring and E, F are two A-modules.

(1°) $E \otimes A^n = E^n \ (n \in N)$

(2°) If the sequence of A-modules $F' \to F \to F'' \to 0$ is exact, then also the sequence $E \otimes F' \to E \otimes F \to E \otimes F'' \to 0$ is exact. (Right exactness of the tensor product)

(3°) If $(F_i)_{i \in I}$; $f_{ij} : F_j \to F_i$ is an inductive system, then

$$E \otimes \varinjlim F_i = \varinjlim (E \otimes F_i).$$

On the other hand these properties characterize completely the functor $\otimes$.

Definition 1 : Let $f = (f_0\ f^1) : X' \to X$ be a morphism of analytic spaces, and $\mathcal{E}$ an $\mathcal{O}_X$-module. Then $f_0^* \mathcal{E}$ is an $f_0^* \mathcal{O}_X$-module and $\mathcal{O}_{X'}$ is also an $f_0^* \mathcal{O}_X$-module (by $f^1 : f_0^* \mathcal{O}_X \to \mathcal{O}_{X'}$).

The analytic pull-back $f^* \mathcal{E}$ of $\mathcal{E}$ by f is defined by scalar extension:

$$f^* \mathcal{E} = f_0^* \mathcal{E} \underset{f_0^* \mathcal{O}_X}{\otimes} \mathcal{O}_{X'}$$

Remark : The inverse image is a particular case of the analytic pull-back.

In fact, if Y is a closed analytic subspace of X and $f : X' \to X$ is a morphism:

$$f^* \mathcal{O}_Y = f_0^*(\mathcal{O}_X/J_Y) \underset{f_0^* \mathcal{O}_X}{\otimes} \mathcal{O}_{X'} \simeq f_0^* \mathcal{O}_X/f_0^* J_Y \underset{f_0^* \mathcal{O}_X}{\otimes} \mathcal{O}_{X'}$$

$$\simeq \mathcal{O}_{X'}/f^1(J_Y).\, \mathcal{O}_{X'} \simeq \mathcal{O}_{f^{-1}(Y)}$$

(The third isomorphism follows from the fact, that $A/I \underset{A}{\otimes} E \simeq E/IE$).

Elementary properties of the analytic pull-back :

(*a*) $(f^* \mathcal{E})_{x'} = (f_0^* \mathcal{E})_{x'} \otimes_{(f_0^* \mathcal{O}_X)_{x'}} \mathcal{O}_{X',x'} \simeq \mathcal{E}_x \otimes_{\mathcal{O}_{X,x}} \mathcal{O}_{X',x'}$ where $x = f_0(x')$ (since $\otimes$ commutes with inductive limits).

(*b*) $f^*(\mathcal{E} \underset{\mathcal{O}_X}{\otimes} \mathcal{F}) = f^* \mathcal{E} \underset{\mathcal{O}_{X'}}{\otimes} f^* \mathcal{F}$, where $\mathcal{E}$ and $\mathcal{F}$ are $\mathcal{O}_X$-modules.

(*c*) If $\mathcal{E}$ is a coherent $\mathcal{O}_X$-module, then $f^* \mathcal{E}$ is a coherent $\mathcal{O}_{X'}$-module.

In fact, $\mathcal{E}$ has a locally finite presentation:
$\mathcal{O}_X^q \to \mathcal{O}_X^p \to \mathcal{E} \to 0$, and f^* is compatible with cokernels, $f^*(\mathcal{O}_X^r) = \mathcal{O}_X^r$.

Special case : The pull-back of vector bundle. Let (E, π) be an analytic vector bundle over the analytic space X, and $f: X' \to X$ a morphism of analytic spaces. The fiber product carries a unique structure of vector bundle over X', such that $\bar{f}$ is a bundle morphism. We call this bundle E'.

$$\begin{tikzcd} E \times_X X' \arrow[r, "\bar{f}"] \arrow[d, "\pi'"] & E \arrow[d, "\pi"] \\ X' \arrow[r, "f"] & X \end{tikzcd}$$

Proposition 1 : Let $\mathcal{E}$ (Resp. $\mathcal{E}'$) be the sheaf of analytic sections of E (Resp. E'). Then $\mathcal{E}' = f^* \mathcal{E}$.

Proof (Sketch) : We have a $f_0^* \mathcal{O}_X$ linear morphism $f_0^* \mathcal{E} \to \mathcal{E}'$, which extends to a morphism $f^* \mathcal{E} \to \mathcal{E}'$. We can prove that this is an isomorphism. Since the question is local with respect to X', we can suppose that E is a trivial bundle over X with fiber $\mathbf{C}^r$, then $\mathcal{E} = \mathcal{O}_X^r$. Also $\mathcal{O}_{X'}^r = f^* \mathcal{O}_X^r$. Therefore $f^* \mathcal{E} = \mathcal{E}'$.

§ 3. *Introduction to flatness by examples*

Let S be an analytic space. By analytic space over s we mean an analytic space X provided with a morphism $\pi : X \to S$. Let S be a simple point in S, and consider $X(s) = f^{-1}(s)$.