

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 14 (1968)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FLATNESS AND PRIVILEGE
Autor: Douady, A.
Kapitel: §3. Introduction to flatness by examples
DOI: <https://doi.org/10.5169/seals-42343>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Remark: The inverse image is a particular case of the analytic pull-back.

In fact, if Y is a closed analytic subspace of X and $f : X' \rightarrow X$ is a morphism:

$$\begin{aligned} f^* \mathcal{O}_Y &= f_0^*(\mathcal{O}_X/J_Y) \otimes_{f_0^* \mathcal{O}_X} \mathcal{O}_{X'} \simeq f_0^* \mathcal{O}_X / f_0^* J_Y \otimes_{f_0^* \mathcal{O}_X} \mathcal{O}_{X'} \\ &\simeq \mathcal{O}_{X'} / f^1(J_Y). \mathcal{O}_{X'} \simeq \mathcal{O}_{f^{-1}(Y)} \end{aligned}$$

(The third isomorphism follows from the fact, that $A/I \underset{A}{\otimes} E \simeq E/IE$).

Elementary properties of the analytic pull-back:

- (a) $(f^* \mathcal{E})_{x'} = (f_0^* \mathcal{E})_{x'} \otimes_{(f_0^* \mathcal{O}_X)_{x'}} \mathcal{O}_{X', x'} \simeq \mathcal{E}_x \otimes_{\mathcal{O}_{X, x}} \mathcal{O}_{X', x'}$ where $x = f_0(x')$ (since \otimes commutes with inductive limits).
- (b) $f^* (\mathcal{E} \otimes_{\mathcal{O}_X} \mathcal{F}) = f^* \mathcal{E} \otimes_{\mathcal{O}_{X'}} f^* \mathcal{F}$, where \mathcal{E} and \mathcal{F} are \mathcal{O}_X -modules.
- (c) If \mathcal{E} is a coherent \mathcal{O}_X -module, then $f^* \mathcal{E}$ is a coherent $\mathcal{O}_{X'}$ -module.

In fact, \mathcal{E} has a locally finite presentation:

$\mathcal{O}_X^q \rightarrow \mathcal{O}_X^p \rightarrow \mathcal{E} \rightarrow 0$, and f^* is compatible with cokernels, $f^* (\mathcal{O}_X^r) = \mathcal{O}_{X'}^r$.

Special case: The pull-back of vector bundle. Let (E, π) be an analytic vector bundle over the analytic space X , and $f : X' \rightarrow X$ a morphism of analytic spaces. The fiber product carries a unique structure of vector bundle over X' , such that \bar{f} is a bundle morphism. We call this bundle E' .

Proposition 1: Let \mathcal{E} (Resp. \mathcal{E}') be the sheaf of analytic sections of E (Resp. E'). Then $\mathcal{E}' = f^* \mathcal{E}$.

Proof (Sketch): We have a $f_0^* \mathcal{O}_X$ linear morphism $f_0^* \mathcal{E} \rightarrow \mathcal{E}'$, which extends to a morphism $f^* \mathcal{E} \rightarrow \mathcal{E}'$. We can prove that this is an isomorphism. Since the question is local with respect to X' , we can suppose that E is a trivial bundle over X with fiber \mathbf{C}^r , then $\mathcal{E} = \mathcal{O}_X^r$. Also $\mathcal{O}_{X'}^r = f^* \mathcal{O}_X^r$. Therefore $f^* \mathcal{E} = \mathcal{E}'$.

§ 3. Introduction to flatness by examples


Let S be an analytic space. By analytic space over s we mean an analytic space X provided with a morphism $\pi : X \rightarrow S$. Let S be a simple point in S , and consider $X(s) = f^{-1}(s)$.

The main purpose of these lectures is to give a precise meaning to the expression:

“ $X(s)$ depends nicely on s ”, and to give a criterion for the “nice” behaviour.

We begin with some examples.


Example 1: X is the closed subspace on \mathbf{C}^2 defined by $(y^2 - x)$, $S = \mathbf{C}$ and $\pi = 1\text{st}$ projection.



$$X(s) = \begin{cases} 2 \text{ simple points if } s \neq 0 \\ \text{double point if } s = 0. \end{cases}$$

Here the behaviour of $X(s)$ is nice.

Example 2: X is the closed subspace of \mathbf{C}^2 defined by (xy) , $S = \mathbf{C}$ and $\pi = 1\text{st}$ projection.



$X(s)$ is given by $(x-s, xy)$, and

$$(x-s, xy) = \begin{cases} (x-s, y) & \text{if } s \neq 0 \\ (x) & \text{if } s = 0. \end{cases}$$

The first case is a simple point, the second one the, y -axis.

A similar example is the map of a point into \mathbf{C} .

In both of these examples the dimension of the fiber makes a jump at one point. We notice, however, that the exceptional point corresponds to an irreducible component of X , and after removing this component π behaves nicely.

This kind of removing is not possible in general, as the following example shows:

Example 3: X is given in \mathbf{C}^3 by $(xz-y)$, and π is the projection on the (x, y) -plane.

If $s = (x_0, y_0)$, then the fiber $X(s)$ is defined by

$$(x-x_0, y-y_0, xz-y) = \begin{cases} \left(x-x_0, y-y_0, z-\frac{y_0}{x_0}\right) & \text{if } x_0 \neq 0 \\ (x, y) & \text{if } x_0 = y_0 = 0 \\ (1) & \text{if } x_0 = 0, y_0 \neq 0. \end{cases}$$

The set of “ nice ” fibers is dense in X , so we cannot remove the z -axis and still get a closed subspace of \mathbf{C}_3 .

§ 4. Algebraic study of flatness

In the following all rings are commutative, with 1, and all modules are unitary.

Definition 1: An A -module E is *flat*, if for every exact sequence of A -modules

$$0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0,$$

the sequence $0 \rightarrow E \otimes F' \rightarrow E \otimes F \rightarrow E \otimes F'' \rightarrow 0$ is also exact. We can also say, because \otimes is right exact, that E is flat, if for every injective homomorphism $F' \rightarrow F$, $E \otimes F' \rightarrow E \otimes F$ is also injective.

Examples of modules which are not flat :

- (1) if $A = \mathbf{Z}$, $E = \mathbf{Z}_2 = \mathbf{Z}/2\mathbf{Z}$, $F = F' = \mathbf{Z}$; then the sequence $0 \rightarrow \mathbf{Z} \xrightarrow{2I} \mathbf{Z}$ ($2I : x' \rightarrow 2x$) is exact. But now $\mathbf{Z}_2 \otimes \mathbf{Z} = \mathbf{Z}_2$, and the homomorphism $\mathbf{Z}_2 \xrightarrow{2I} \mathbf{Z}_2$ is the zero homomorphism, which is not injective. So \mathbf{Z}_2 is not a flat \mathbf{Z} module.
- (2) If $A = \mathbf{C}\{x\}$, $E = \mathbf{C} = \mathbf{C}\{x\}/(x)$, $F = F' = \mathbf{C}\{x\}$, then the sequence $0 \rightarrow F \xrightarrow{xI} F'$ ($xI : p(x) \rightarrow xp(x)$) is exact. But the homomorphism $E \xrightarrow{xI} E$ is not injective.

Proposition 1: If A is an integral domain and E a flat A -module, then E is torsion-free.

Proof: Let $a \in A$, $a \neq 0$. Because A is an integral domain, the sequence $0 \rightarrow AA \xrightarrow{aI} (aI : x \rightarrow ax)$ is exact. Since E is flat, the sequence $0 \rightarrow E \xrightarrow{aI} E$ is also exact. In other words E has no torsion elements.

Proposition 2: If A is a principal-ideal domain, then E is flat if and only if E is torsionfree.

Proof: See corollary of prop. 6.

Examples of flat modules :

- (1) The inductive limit of flat modules is flat, because the inductive limit preserves exactness, and it commutes with the tensor product.