

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 14 (1968)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FLATNESS AND PRIVILEGE
Autor: Douady, A.
Kapitel: §2. Privileged polycylinders
DOI: <https://doi.org/10.5169/seals-42343>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$g|_W : \begin{cases} F_2 \rightarrow 0 \\ G_2 W \simeq F_3 \end{cases} .$$

If $p: E_{2W} \rightarrow F_2$ is the projection with kernel G_{2W} , the map, $p \circ f: E_{1W} \rightarrow F_2$ is a split epimorphism in x_0 . Again by prop. 2 we have over an open neighbourhood $U \subset W$ of x_0 a decomposition $E_{1U} = F_1 \oplus G_{1U}$ (with $F_1 = \text{Ker } p \circ f$)

$$(p \circ f)|_U : \begin{cases} F_1 \rightarrow 0 \\ G_{1U} \xrightarrow{\sim} F_{2U} \end{cases} .$$

The image $f|_U(F_1)$ is contained in G_{2U} . But $g|_U \circ f|_U = 0$ and $g|_{G_{2U}}$ is a monomorphism hence $f|_U: F_1 \rightarrow 0$. We get finally (restricting all our morphisms to U)

$$f|_U : \begin{cases} F_{1U} \rightarrow 0 \\ G_{1U} \simeq F_{2U} \end{cases} \quad g|_U : \begin{cases} F_{2U} \rightarrow 0 \\ G_{2U} \xrightarrow{\sim} F_{3U} \end{cases} .$$

§ 2. Privileged polycylinders

Definition 1: A polycylinder in \mathbf{C}^n is a compact set K of the form $K = K_1 \times \dots \times K_n$ where each K_i is a compact, convex subset of \mathbf{C} , with nonempty interior. If each K_i is a disc, then K is a polydisc. We first recall the following theorem of Cartan.

Theorem 1: Let K be a polycylinder contained in an open subset U of \mathbf{C}^n . Let \mathcal{F} be a coherent analytic sheaf on U .

- (A) There exists an open neighbourhood of K over which \mathcal{F} admits a finite free resolution

$$0 \rightarrow \mathcal{L}_n \rightarrow \dots \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow \mathcal{F} \rightarrow 0 .$$

- (B) $H^q(K, \mathcal{F}) = 0$ for $q > 0$.

(Reference: For instance Gunning and Rossi.)

We have the following consequences of this theorem:

- 1) Given a finite free resolution

$$0 \rightarrow \mathcal{L}_n \rightarrow \dots \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow \mathcal{F} \rightarrow 0$$

of a coherent sheaf \mathcal{F} , the sequence

$$0 \rightarrow \mathcal{L}_n(K) \rightarrow \dots \rightarrow \mathcal{L}_0(K) \rightarrow \mathcal{F}(K) \rightarrow 0$$

is an $\mathcal{O}_U(K)$ -free resolution of $\mathcal{F}(K)$.

2) Given a short exact sequence of coherent sheaves

$$0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0,$$

then the sequence

$$0 \rightarrow \mathcal{F}'(K) \rightarrow \mathcal{F}(K) \rightarrow \mathcal{F}''(K) \rightarrow 0 \quad \text{is exact.}$$

Let \mathcal{F} be a coherent analytic sheaf on U , and let $K \subset U$ be a polycylinder. If V is an open neighbourhood of K , then $\mathcal{F}(V)$ can be equipped with a Fréchet-space structure (see: Malgrange).

Hence we can give $\mathcal{F}(K)$ the structure of inductive limit of Fréchet-spaces. It is however essential for certain purposes to have Banach-spaces. This can be obtained by choosing a space slightly different from $\mathcal{F}(K)$ and by choosing K in a “privileged” way.

Let $B(K) = \{f : K \rightarrow \mathbf{C} \mid f \text{ continuous on } K \text{ and analytic on } \overset{\circ}{K}\}$, then $B(K)$ is Banach algebra and $B(K) \subset C(K)$. The sections of \mathcal{O}_U over K are elements of $B(K)$, and $B(K)$ is in fact the uniform closure of $\mathcal{O}_U(K)$ in $C(K)$.

If $\mathcal{L} = \mathcal{O}_U^r$, we define $B(K, \mathcal{L}) = B(K)^r$. Then $B(K; \mathcal{L})$ is a free $B(K)$ -module, and since $\mathcal{L}(K) = \mathcal{O}_U(K)^r$, we have $B(K; \mathcal{L}) = B(K) \otimes_{\mathcal{O}_U(K)} \mathcal{L}(K)$.

We now assume that \mathcal{F} is a coherent sheaf on U , where $U \subset \mathbf{C}^n$ is open. Consider a free resolution

$$(R) \quad 0 \rightarrow \mathcal{L}_n \rightarrow \dots \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow \mathcal{F} \rightarrow 0 \quad \text{of } \mathcal{F}.$$

From (R) we get an $\mathcal{O}_U(K)$ -free resolution of $\mathcal{F}(K)$

$$(R') \quad 0 \rightarrow \mathcal{L}_n(K) \rightarrow \dots \rightarrow \mathcal{L}_1(K) \rightarrow \mathcal{L}_0(K) \rightarrow \mathcal{F}(K) \rightarrow 0.$$

Taking the tensorproduct $B(K) \otimes_{\mathcal{O}_U(K)}$ we get the complex

$$B(K; \mathcal{L}_\cdot) : 0 \rightarrow B(K; \mathcal{L}_n) \rightarrow \dots \rightarrow B(K; \mathcal{L}_1) \rightarrow B(K; \mathcal{L}_0).$$

Definition 2: The polycylinder K is called \mathcal{F} -privileged if the complex $B(K; \mathcal{L}_\cdot)$ is split-exact in every degree > 0 .

Remark: The property of being \mathcal{F} -privileged is independent of the resolution (R) .

The exactness of $B(K; \mathcal{L})$ can be expressed by $\text{Tor}_i^{\mathcal{O}(K)}(B(K), \mathcal{F}(K)) = 0$, for every $i > 0$, and Tor is independent of the resolution (R) . It is a little

more complicated to show, that the splitting property is independent of (R) , and this is omitted.

Since $B(K; \mathcal{L}_i)$ is a Banach space, the image and its complement are thus Banach spaces if K is \mathcal{F} -privileged. In this case we define $B(K; \mathcal{F}) = \text{Coker } (B(K, \mathcal{L}_1) \rightarrow B(K, \mathcal{L}_0)) = B(K) \otimes_{\mathcal{O}_U} \mathcal{F}(K)$ and we get a $B(K)$ -module, which is a Banach-space.

Warning : In the definition of split-exactness, the subspaces are splitting vector spaces, but they are not splitting $B(K)$ -modules in general.

We have the following important theorem about the existence of privileged polycylinders:

Theorem 2 : Let U be an open subset of \mathbf{C}^n , and let \mathcal{F} be a coherent analytic sheaf on U . For any $x \in U$ there exists a fundamental system of neighbourhoods of x in U , which are \mathcal{F} -privileged polycylinders.

For the proof, see Douady: § 7, 4, th 1.

Example : (Curves in \mathbf{C}^2) Let $U \subset \mathbf{C}^2$ be an open connected neighbourhood of the origin, and let $h: U \rightarrow \mathbf{C}$ be analytic and $h \neq 0$.

Let X be the curve given by h , that is $X = h^{-1}(0)$, $\mathcal{O}_X = \mathcal{O}_{U,I}/(h)$. We have an exact sequence $0 \rightarrow \mathcal{O}_U \rightarrow \mathcal{O}_U \rightarrow \mathcal{O}_X \rightarrow 0$. Consider a polycylinder $K = K_1 \times K_2 \subset U$. By definition K is \mathcal{O}_X -privileged if and only if $h: B(K) \rightarrow B(K)$ is a split monomorphism.

Let K_j denote the boundary of K_j , and define $\ddot{K} = \dot{K}_1 \times \dot{K}_2$ (\ddot{K} is called the Šilov Boundary of K).

Proposition 1 : (a) The following conditions are equivalent:

- (i) $h: B(K) \rightarrow B(K)$ is a monomorphism.
- (i') $\exists a > 0$ such that $\|hf\| \geq a\|f\|$, $\forall f \in B(K)$.
- (ii) $X \cap \ddot{K} = \emptyset$.

(b) If $(K_1 \times K_2) \cap X = \emptyset$, then h is a split monomorphism (i.e. K is \mathcal{O}_X privileged).

Proof: (a) (i) \Leftrightarrow (i') is a well known fact from the theory of normed vector spaces.

(ii) \Rightarrow (i'). Assume $X \cap \ddot{K} = \emptyset$. If $f \in B(K)$, then it follows from the maximum principle that $\|f\| = \sup_K |f(x)| = \sup_{\ddot{K}} |f(x)|$. Since $h(x) \neq 0$

whenever $x \in K$, we get $a = \inf_K |h(x)| > 0$. Hence $\|hf\| = \sup_K |hf(x)| \geq \geq a \sup_K |f(x)| = a \|f\|$.

(i') \Rightarrow (ii). Suppose that $X \cap K \neq \emptyset$ and $x = (x_1, x_2) \in X \cap K$. We choose an analytic function $f_1 : U_1 \rightarrow \mathbf{C}$, where $U_1 \supset K_1$, and U_1 is open, such that $f_1(x_1) = 1$, $|f_1(z)| < 1$ if $z \in K_1$, $z \neq x_1$. Similarly we choose an analytic function $f_2 : U_2 \rightarrow \mathbf{C}$, with the same properties. Consider the function $f \in B(K) : (z_1, z_2) \mapsto f_1(z_1)f_2(z_2)$. Since $h(x) = 0$ it follows that the sequence $\{hf^n\}$ converges pointwise to 0 in K .

Applying Dini's theorem we get $\|hf^n\| \rightarrow 0$. From the inequality $a \|f^n\| \leq \|hf^n\|$ we get $\|f^n\| \rightarrow 0$, which is a contradiction, because for every $n : f^n(x) = 1$.

(b) Use the Weierstrass preparation theorem (extended form).

Question. Does the condition (ii) imply that $h : B(K) \rightarrow B(K)$ is a split monomorphism?

IV. FLATNESS AND PRIVILEGE

§ 1. *Morphisms from an analytic space into $B(K)$*

Let S be an analytic space and K a polycylinder in an open set $U \subset \mathbf{C}^n$. We want to construct an \mathcal{O}_S -algebra homomorphism $\phi : \mathcal{O}_{S \times U}(S \times U) \rightarrow \mathcal{H}(S; B(K))$.

- (a) Consider first $S = U' \subset \mathbf{C}^m$, U' -open. If $h \in \mathcal{O}_{U' \times U}(U' \times U)$ and $s \in U'$, $x \in K$, define $(\phi(h)(s))(x) = h(s, x)$. Using the Cauchy integral, one can show that $\phi(h)$ is analytic. On the other hand its obvious that ϕ is an $\mathcal{O}_{U'}$ -algebra homomorphism.
- (b) Let S have a special model in the polydisc Δ in \mathbf{C}^m , defined by a sheaf \mathcal{J} of ideals of \mathcal{O}_Δ , and let \mathcal{J} be generated by f_1, \dots, f_p , V -a polycylinder neighbourhood of K in U . By Cartan's theorem B for a polycylinder,

the sequence $0 \rightarrow \mathcal{J}(\Delta \times V) \xrightarrow{\pi} \mathcal{O}(\Delta \times V) \rightarrow \mathcal{O}(S \times V) \rightarrow 0$ is exact. If we denote by $\tilde{\pi}$ the projection $\mathcal{H}(\Delta, B(K)) \rightarrow \mathcal{H}(S, B(K))$, $(f_1, \dots, f_p) \cdot \mathcal{H}(\Delta, B(K)) \subset \subset \text{Ker } \tilde{\pi}$. Therefore, because π is surjection, there exists a unique

$\phi : \mathcal{O}(S \times V) \rightarrow \mathcal{H}(S, B(K))$, such that the diagram