

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 14 (1968)  
**Heft:** 1: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** COMPACT ANALYTICAL VARIETIES  
**Autor:** Narasimhan, Raghavan  
**Kapitel:** 1. Preliminaries  
**DOI:** <https://doi.org/10.5169/seals-42344>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# COMPACT ANALYTICAL VARIETIES

by Raghavan NARASIMHAN

## CONTENTS

Introduction

1. Preliminaries
  2. The vanishing theorem of Kodaira
  3. An imbedding theorem
  4. Line bundles associated to a divisor
  5. Meromorphic forms
  6. The Atiyah-Hodge theorem
  7. Lefschetz' theorem on hyperplane sections
- References.

## INTRODUCTION

These lectures deal with the vanishing theorem of Kodaira (cf. e.g. [2], p. 344) and some of its consequences, and with Lefschetz' theorem on hyperplane sections (cf. [1]). Only complex manifolds (and not complex spaces) are considered, but most of the results in the first part could be carried over to the more general case (with similar proofs).

## 1. PRELIMINARIES

We first give some definitions:

*Definition 1.1.* Let  $V$  be a complex manifold and  $D$  a relatively compact, open subset of  $V$ . Then  $D$  is *strongly pseudoconvex* if for every  $x_0 \in \partial D$  there exist a neighbourhood  $U$  of  $x_0$  and a real-valued  $C^2$ -function  $\varphi$  defined in  $U$  such that

$$(1) \quad d\varphi(x_0) \neq 0,$$

$$(2) \quad H(\varphi)(x_0) > 0 \text{ for all } \alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{C}^n - \{0\}.$$

(Here  $H(\varphi)$  is the complex Hessian form

$$\sum_{i,j=1}^n \frac{\partial^2 \varphi}{\partial z_i \partial \bar{z}_j} \alpha_i \bar{\alpha}_j$$

with respect to some system of local coordinates),

$$(3) \quad D \cap U = \{x \in U; \varphi(x) < 0\}.$$

It can be shown that strong pseudoconvexity of  $D$  is equivalent to the following property: For every  $x_0 \in \partial D$  there exist a neighbourhood  $U$  of  $x_0$  and a biholomorphic mapping  $f: U \rightarrow \Omega \subset \mathbb{C}^n$  such that  $f(U \cap D)$  has a strictly convex boundary (in the Euclidean sense).

*Definition 1.2.* Let  $V$  be a complex manifold and  $A$  a subset of  $V$ . We say that  $A$  "can be blown down to a point" if there exist an analytic space  $X$ , a point  $x_0 \in X$ , and a mapping  $f: V \rightarrow X$  such that  $f(A) = x_0$  and  $f: V - A \rightarrow X - \{x_0\}$  is an analytic isomorphism.

To give an example of sets which can be blown down to a point, we mention the following theorem (for a proof see [2], pp. 338 and 340):

*Theorem 1.3.* If  $D$  is strongly pseudoconvex, then  $D$  has a maximal compact analytic subset  $A$  whose dimension at any point is  $> 0$  and each component of  $A$  can be blown down to a point.

*Lemma 1.4.* If  $A$  can be blown down to a point, then  $A$  has a fundamental system of strongly pseudoconvex neighbourhoods.

*Proof.* Let  $X$ ,  $x_0$ , and  $f$  be as in Definition 1.2. The lemma follows from the fact that the inverse image of a strongly pseudoconvex neighbourhood of  $x_0$  is a strongly pseudoconvex neighbourhood of  $A$ .

We now introduce the concept of holomorphic line bundle.

*Definition 1.5.* Suppose  $X$  is a complex manifold. A holomorphic line bundle  $F$  on  $X$  is a complex manifold  $F$  together with a mapping  $\pi$  with the following properties:

- (i)  $\pi: F \rightarrow X$  is a holomorphic map (called projection) onto  $X$ .
- (ii) For  $x \in X$ ,  $\pi^{-1}(x)$  has the structure of a one-dimensional vector space over the complex numbers.
- (iii) For each  $x \in X$  there exist a neighbourhood  $U$  of  $x$  and a holomorphic mapping  $h$  of  $F|U = \pi^{-1}(U)$  onto  $U \times \mathbb{C}$  such that  $h^{-1}$  is holomorphic and  $h| \pi^{-1}(a)$  is a  $\mathbb{C}$ -isomorphism onto  $\{a\} \times \mathbb{C}$  for every  $a \in U$ .

Let  $\{U_i\}$  be an open covering of  $X$  such that for each  $i$  we have a mapping  $h_i$  of  $F|U_i$  onto  $U_i \times \mathbb{C}$  with the properties in (iii) above. If  $U_i \cap U_j \neq \emptyset$ , we get a mapping  $h_i \circ h_j^{-1}: (U_i \cap U_j) \times \mathbb{C} \rightarrow (U_i \cap U_j) \times \mathbb{C}$ . If  $(x, c) \in (U_i \cap U_j) \times \mathbb{C}$ , then the image of  $(x, c)$  under the mapping

$h_i \circ h_j^{-1}$  can be written  $(x, \gamma'(x, c))$  where  $\gamma'(x, c) \in \mathbf{C}$ . According to the last property in (iii), for fixed  $x \in U_i \cap U_j$  the mapping  $c \rightarrow \gamma'(x, c)$  is a  $\mathbf{C}$ -isomorphism of  $\mathbf{C}$  onto itself. Therefore

$$\gamma'(x, c) = g_{ij}(x) \cdot c, \text{ where } g_{ij}(x) \neq 0, \quad (1.1)$$

and it is easily seen that  $g_{ij}$  is holomorphic in  $U_i \cap U_j$ .

The functions  $g_{ij}$  obviously satisfy the cocycle conditions

$$g_{ij} g_{jk} g_{ki} = 1 \quad \text{on} \quad U_i \cap U_j \cap U_k, \quad (1.2)$$

$$g_{ij} g_{ji} = 1 \quad \text{on} \quad U_i \cap U_j. \quad (1.3)$$

The  $g_{ij}$  are called transition functions corresponding to the line bundle  $F$ .

Conversely, it is easy to prove (cf. [4], p. 135) that given an open covering  $\{U_i\}$  and functions  $g_{ij}$  without zeros in  $U_i \cap U_j$  which satisfy the cocycle conditions, we can construct a line bundle which has  $g_{ij}$  as transition functions.

Now, let  $F$  be a line bundle over a complex manifold  $X$ , and let  $\pi$  be the corresponding projection. We denote  $\pi^{-1}(a)$  by  $F_a$ . Let  $F_a^*$  be the  $\mathbf{C}$ -dual of  $F_a$ . Then

$$F^* = \bigcup_{a \in X} F_a^*$$

is in a natural way a holomorphic line bundle over  $X$ , which is called the dual bundle of  $F$ . If  $F$  has transition functions  $\{g_{ij}\}$ , then  $F^*$  has transition functions  $\{g_{ij}^{-1}\}$ .

*Definition 1.6.* Let  $F$  be a holomorphic line bundle over a compact complex manifold. Then  $F$  is *negative* if the zero cross section  $\mathfrak{o}$  of  $F$  can be blown down to a point.  $F$  is *positive* if the dual bundle is negative.

In the sequel we let  $\underline{F}$  denote the sheaf of germs of analytic sections of a line bundle  $F$ .

## 2. THE VANISHING THEOREM OF KODAIRA

This is the following theorem, which is our first main result:

*Theorem 2.1.* Let  $X$  be a compact connected complex manifold and  $F$  a positive line bundle on  $X$  and  $S$  a coherent analytic sheaf on  $X$ . Then there exists an integer  $k(S, F)$  such that for  $k > k(S, F)$  we have  $H^q(X, S \otimes \underline{F}^k) = 0$  ( $\forall q \geq 1$ ).

The proof uses the following finiteness theorem: