

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 14 (1968)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: COMPACT ANALYTICAL VARIETIES
Autor: Narasimhan, Raghavan
Kapitel: 5. Meromorphic forms
DOI: <https://doi.org/10.5169/seals-42344>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

on A . Since $f \in H^0(\mathbf{P}^n, F^m)$, this gives the desired homogeneous polynomial.

To prove the theorem, it now suffices to consider all homogeneous polynomials which vanish on A without being identically zero and apply the Hilbert basis theorem.

5. MEROMORPHIC FORMS

Let X be a complex manifold. A holomorphic differential form is a form which in local coordinates can be written as a finite sum

$$\omega = \sum a_{i_1 \dots i_k} dz_{i_1} \wedge \dots \wedge dz_{i_k} \quad (5.1)$$

with holomorphic coefficients $a_{i_1 \dots i_k}$.

A form is called meromorphic if it has locally the form (5.1) with coefficients that are meromorphic functions. Every meromorphic function can be written locally as $f\omega$ where f is a meromorphic function and ω a holomorphic form. The exterior differentiation d , satisfying $d^2 = 0$, extends naturally to meromorphic forms.

Let D be a divisor of X and let $\Omega^p(k, D) = \Omega^p(X, k, D)$ be the sheaf of germs of meromorphic p -forms on X with poles only on D and of order $\leq k$, and let $\Omega^p = \Omega^p(X)$ be the sheaf of germs of holomorphic p -forms on X .

Lemma 5.1. There is a natural isomorphism

$$\Omega^p(k, D) \simeq \Omega^p \otimes \underline{F^k}.$$

Proof. A germ in $\Omega^p(k, D)$ at $a \in X$ is represented by a form $f\omega$, where f is a meromorphic function in a neighbourhood U of a , with poles only on D and of order $\leq k$, and ω is a holomorphic form on U . Now to f corresponds biuniquely a section $s \in \Gamma(U, F^k)$ (see Sect. 4), which gives a germ $s_a \in \underline{F^k}_a$. Also ω defines a germ $\omega_a \in \Omega^p_a$.

The desired mapping $\Omega^p(K, D) \rightarrow \Omega^p \otimes \underline{F^k}$ is now uniquely defined by

$$f\omega \rightarrow \omega_a \otimes s_a.$$

To see that it is an isomorphism, it is sufficient to observe that the inverse mapping of $\Omega^p \otimes \underline{F^k}$ into $\Omega^p(k, D)$ is induced by the bilinear mapping $\Omega^p \oplus \underline{F^k} \rightarrow \Omega^p(k, D)$, which is given by

$$(\omega_a, s_a) \rightarrow (f\omega)_a, \quad (a \in X).$$

where f is the meromorphic function determined by s_a by the procedure described just before Th. 4.1.

Now let X be a compact submanifold of \mathbf{P}^n and consider hyperplanes H_c in \mathbf{P}^n , given in homogeneous coordinates z_0, \dots, z_n by equations

$$\sum_0^n c_j z_j = 0 \text{ where } c = (c_0, \dots, c_n) \neq 0.$$

Theorem 5.2. There is an open dense set Ω in \mathbf{C}^{n+1} such that if $c = (c_0, \dots, c_n) \in \Omega$, the hyperplane section $D_c = H_c \cap X$ is a non-singular analytic subset of X .

The proof is omitted here.

Let $D = H \cap X$ be a non-singular hyperplane section of X . To D is then associated a positive line bundle F on X (see Sect. 4). By Kodaira's vanishing theorem there is a k_0 such that

$$H^q(X, \Omega^p \otimes F^k) = 0, \quad (\forall q \geq 1, \forall k \geq k_0).$$

Using the isomorphism in Lemma 5.1, we have therefore proved.

Lemma 5.3. If D is a non-singular hyperplane section of a compact submanifold X of \mathbf{P}^n , then there exists k_0 such that

$$H^q(X, \Omega^p(k, D)) = 0, \quad (\forall q \geq 1, \forall k \geq k_0).$$

6. THE ATIYAH-HODGE THEOREM

We first recall two well-known theorems.

Let X be a paracompact C^∞ manifold and let \mathcal{E}^p be the sheaf of germs of C^∞ p -forms on X ($p=0, 1, \dots$).

Then the sequence

$$0 \rightarrow \mathbf{C} \xrightarrow{i} \mathcal{E}^0 \xrightarrow{d} \mathcal{E}^1 \xrightarrow{d} \mathcal{E}^2 \xrightarrow{d} \dots \tag{6.1}$$

is exact (Poincaré's lemma), and

$$H^q(X, \mathcal{E}^p) = 0, \quad (\forall q \geq 1, \forall p \geq 0), \tag{6.2}$$

because the \mathcal{E}^p are fine sheaves, i.e. they have partitions of unity. From (6.1) we get the sequence

$$0 \rightarrow \Gamma(X, \mathcal{E}^0) \rightarrow \Gamma(X, \mathcal{E}^1) \rightarrow \dots,$$

which need not be exact. Put