

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 14 (1968)  
**Heft:** 1: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** COMPACT ANALYTICAL VARIETIES  
**Autor:** Narasimhan, Raghavan  
**Kapitel:** 7. Lefschetz' theorem on hyperplane sections  
**DOI:** <https://doi.org/10.5169/seals-42344>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 12.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Now  $\omega$  takes the form

$$\omega - d\beta' = dz_1 \wedge \alpha'. \quad (6.23)$$

We distinguish the two cases  $k > 1$  and  $k = 1$ . In the first case we get from (6.23)

$$dz_1 \wedge \delta\alpha' = 0,$$

which implies that  $\delta\alpha' = 0$ . Since  $\alpha'$  is a form of type  $q-1 \geq 1$ , we can apply once again Lemma 6.8 and get  $\alpha' = \delta\alpha''$ . Thus  $dz_1 \wedge \alpha' = d(dz_1 \wedge \alpha'')$ , and we get  $\omega = d(\beta' + dz_1 \wedge \alpha'')$ . This proves that the cohomology under consideration is trivial for  $k > 1$ .

Finally, in the case  $k = 1$ ,  $\alpha'$  is a meromorphic function, independent of  $z_2, \dots, z_n$ . Thus by (6.23),  $\omega = d\gamma$  for some  $\gamma$  if and only if in the Laurent expansion of  $\alpha'$  the coefficient of  $z_1^{-1}$  is zero. Thus the cohomology in dimension 1 is generated by  $z_1^{-1} dz_1$ , which completes the proof of Theorem 6.4.

## 7. LEFSCHETZ' THEOREM ON HYPERPLANE SECTIONS

The Lefschetz theorem in the slightly more general setting proved by Andreotti and Frankel [1], is the following:

*Theorem 7.1.* Let  $V$  be a submanifold of  $\mathbf{P}^n$  of complex dimension  $d$  and let  $D$  be a hyperplane section of  $V$  (not necessarily non-singular). Then there are natural isomorphisms

$$H^q(V, \mathbf{Z}) \simeq H^q(D, \mathbf{Z}), \quad (\forall q < d-1),$$

and a natural injection

$$H^{d-1}(V, \mathbf{Z}) \rightarrow H^{d-1}(D, \mathbf{Z}).$$

*Proof.*  $X = V - D$  is a Stein manifold, since it is imbedded as a closed submanifold of  $\mathbf{C}^n$ . Now one knows that

$$H^q(V, D, \mathbf{Z}) \simeq H_c^q(X, \mathbf{Z}), \quad (7.1)$$

where the  $c$  indicates cohomology with compact support. On the other hand, since  $X$  is a topological manifold of dimension  $2d$ , Poincaré duality gives

$$H_c^q(X, \mathbf{Z}) \simeq H_{2d-q}(X, \mathbf{Z}). \quad (7.2)$$

Now we shall use the following theorem:

*Theorem 7.2.* Let  $X$  be a Stein manifold of dimension  $d$ . Then

$$H_r(X, \mathbf{Z}) = 0, \quad (\forall r > d). \quad (7.3)$$

Suppose this theorem is proved. Then (7.1) – (7.3) gives

$$H^q(V, D, \mathbf{Z}) = 0. \quad (\forall q < d). \quad (7.4)$$

Now we have the exact sequence

$$\dots \rightarrow H^q(V, D, \mathbf{Z}) \rightarrow H^q(V, \mathbf{Z}) \rightarrow H^q(D, \mathbf{Z}) \rightarrow H^{q+1}(V, D, \mathbf{Z}) \rightarrow \dots,$$

and using (7.4) we conclude that the mapping

$$H^q(V, \mathbf{Z}) \rightarrow H^q(D, \mathbf{Z})$$

is an isomorphism onto when  $q < d-1$  and an injection when  $q = d-1$ .

This proves Lefschetz' theorem.

The proof of Theorem 7.2 is based on *Morse theory*. Let  $X$  be a  $C^\infty$ -manifold with countable base. If  $f$  is a real-valued  $C^\infty$ -function on  $X$ , then a point  $a \in X$  is called *critical* for  $f$  if  $df(a) = 0$ . A critical point  $a$  is *non-degenerate*, if in local coordinates  $f(x) - f(a) = \sum a_{ij} (x_i - a_i)(x_j - a_j) + o(|x-a|^2)$ , where the symmetric matrix  $(a_{ij})$  is non-singular. It is non-degenerate of index  $r$  if  $(a_{ij})$  has  $r$  eigenvalues  $< 0$ . The non-degenerate critical points for  $f$  are necessarily isolated. We now quote some facts from Morse theory; for proofs, see [6].

*Lemma 7.3.* Suppose that  $f \in C^\infty(X)$ ,  $f \geq 0$ ,  $\alpha < \beta$ , and that  $X_\beta = \{x \in X; f(x) \leq \beta\}$  is compact.

(a) If  $f$  has no critical points in  $\{x \in X; \alpha \leq f(x) \leq \beta\}$ , then  $X_\alpha$  is a deformation retract of  $X_\beta$ , and hence

$$H_r(X_\beta, X_\alpha, \mathbf{Z}) = 0, \quad (\forall r \geq 0).$$

(b) If all critical points of  $f$  in  $\{x \in X; \alpha \leq f(x) \leq \beta\}$  are non-degenerate of index  $\leq d$ , then

$$H_r(X_\beta, X_\alpha, \mathbf{Z}) = 0, \quad (\forall r > d).$$

In particular, if all critical points of  $f$  in  $X_\beta$  are non-degenerate of index  $\leq d$ , then

$$H_r(X_\beta, \mathbf{Z}) = 0, \quad (\forall r > d).$$

In the proof of Theorem 7.2 we shall also use the following lemma of Morse:

*Lemma 7.4.* Let  $X$  be a  $C^\infty$ -manifold with countable base. Then every real function  $g \in C^\infty(X)$  can be approximated in the topology of  $C^\infty(X)$  by real functions  $f \in C^\infty(X)$ , whose critical points are all non-degenerate.

The topology of  $C^\infty(X)$  is the topology of uniform convergence of all derivatives on compact sets. Therefore the lemma explicitly means the following:

Let  $\varepsilon > 0$ , an integer  $r \geq 0$  and a compact set  $K \subset X$  be given, and let  $K = K_1 \cup \dots \cup K_k$ , where each  $K_j$  is compact and contained in an open set  $U_j$ , where we have a coordinate system. Then there is a function  $f$  of the prescribed type such that

$$\sup_j \sup_{|\alpha| \leq r} \sup_{x \in K_j} |D^\alpha f(x) - D^\alpha g(x)| < \varepsilon.$$

(Here  $D^\alpha$  means a derivative of order  $|\alpha|$  in the coordinates on  $U_j$ .)

To prove Lemma 7.4 we shall use a Lemma of Sard (see [8, Ch. I, §3, Th. 4]):

*Lemma 7.5.* Let  $\Omega$  be an open subset of  $\mathbf{R}^n$  and  $f: \Omega \rightarrow \mathbf{R}^n$  a  $C^1$ -mapping. Let  $A$  be the critical set of  $f$ , i.e. the set of  $a \in \Omega$  where  $\det(\partial f_i(a)/\partial x_j) = 0$ . Then  $f(A)$  has Lebesgue measure 0 in  $\mathbf{R}^n$ . In particular,  $f(A)$  is nowhere dense in  $\mathbf{R}^n$ .

*Proof of Lemma 7.4.* Suppose first that  $X$  is an open subset  $\Omega$  of  $\mathbf{R}^n$ . If  $g \in C^\infty(\Omega)$  is realvalued, consider the mapping

$$\varphi: \Omega \ni x \rightarrow (\partial g / \partial x_1, \dots, \partial g / \partial x_n) \in \mathbf{R}^n.$$

The critical set  $A$  of  $\varphi$  is the set in  $\Omega$  where

$$\det(\partial^2 g / \partial x_i \partial x_j) = 0.$$

The lemma of Sard, applied to  $\varphi$ , shows that there are arbitrarily small  $\varepsilon_1, \dots, \varepsilon_n \in \mathbf{R}$  such that  $(\varepsilon_1, \dots, \varepsilon_n) \notin \varphi(A)$ . Put

$$f(x) = g(x) - \varepsilon_1 x_1 - \dots - \varepsilon_n x_n.$$

A point  $x \in \Omega$  is a critical point of  $f$  if and only if  $\partial g / \partial x_j = \varepsilon_j$ , ( $j=1, \dots, n$ ).

At such points  $\varphi(x) = (\varepsilon_1, \dots, \varepsilon_n) \in \varphi(A)$  and hence  $\det(\partial^2 g / \partial x_i \partial x_j) \neq 0$ . Hence all critical points of  $f$  are non-degenerate.

Since  $\varepsilon_1, \dots, \varepsilon_n$  can be chosen arbitrarily small, the lemma is proved in the case  $X = \Omega$ .

The general case now follows by a category argument. From the special case we conclude that we can cover  $X$  by denumerably many relatively

compact open subsets  $U_j$  of  $X$ , such that  $\mathcal{U}_j$  is dense in the space of real  $C^\infty$ -functions, where  $\mathcal{U}_j$  denotes the set of real  $C^\infty$ -functions, whose critical points in  $\bar{U}_j$  are all non-degenerate. It is also easy to see that every  $\mathcal{U}_j$  is open in the space of real  $C^\infty$ -functions. Since this space is a real Fréchet space, we can therefore use Baire's theorem to conclude that the set of all real  $C^\infty$ -functions, whose critical points in  $X$  are all non-degenerate, i.e.  $\cap \mathcal{U}_j$ , is dense. This proves the lemma of Morse.

*Proof of Theorem 7.2.* Let  $X$  be a Stein manifold of dimension  $d$ , and let  $K$  be a compact subset of  $X$  such that

$$K = \{x \in X; |f(x)| \leq \|f\|_K, \quad \forall f \text{ holomorphic on } X\}.$$

(Since  $X$  is a Stein manifold, every compact subset of  $X$  is contained in some  $K$  of this kind.) Choose an open set  $U$  such that  $K \subset U \subset \subset X$ . For every  $a \in \partial U$  we can find a holomorphic function  $f$  on  $X$  such that  $|f(x)| \geq 1$  in a neighbourhood of  $a$  and  $\|f\|_K < 1$ . Since  $\partial U$  is compact, we can therefore choose holomorphic functions  $f_1, \dots, f_k$  on  $X$  such that

$$\max |f_j(a)| \geq 1, \quad (\forall a \in \partial U),$$

and

$$\|f_j\|_K < 1, \quad (\forall j).$$

By replacing each  $f_j$  by a sufficiently high power, we can also arrange that the function

$$p(x) = \sum |f_j(x)|^2$$

satisfies  $p(x) < 1$  on  $K$  and  $p(x) \geq 1$  on  $\partial U$ . We can also assume that the rank of  $(f_1, \dots, f_k)$  is maximal at all points of  $U$ .

Now  $p \in C^\infty(X)$ ,  $p \geq 0$ , and  $U_\beta = \{x \in U; p(x) \leq \beta\}$  is compact and contains  $K$  if  $\beta < 1$  is chosen so that  $p(x) < \beta$  in  $K$ . By calculating the Levi form and using the maximality of the rank of  $(f_1, \dots, f_k)$ , we see that  $p$  is strongly plurisubharmonic.

Because of Morse's lemma we can also assume that all critical points of  $p$  in  $U_\beta$  are non-degenerate. We shall prove that they are all of index  $\leq d$ .

We expand  $p$  at a critical point  $a \in U_\beta$  in a local coordinate system:

$$\begin{aligned} p(x) &= p(a) + 2\operatorname{Re} \sum \frac{\partial^2 p(a)}{\partial z_i \partial z_j} (z_i - a_i)(z_j - a_j) \\ &\quad + \sum \frac{\partial^2 p(a)}{\partial z_i \partial \bar{z}_j} (z_i - a_i)(\bar{z}_j - \bar{a}_j) + \dots \\ &= p(a) + \operatorname{Re} Q(z - a) + L(z - a) + \dots \end{aligned}$$

Here  $L(z-a)$  is the Levi form of  $p$  at the point  $a$ . Now, since  $p$  is strongly plurisubharmonic, we can choose the coordinates so that  $L(z-a) = |z-a|^2$ . Then we see that if  $\zeta$  is an eigenvector corresponding to an eigenvalue  $< 0$  of the symmetric matrix of the real quadratic form  $\operatorname{Re} Q(z) + L(z)$ , then  $i\zeta$  is an eigenvector corresponding to an eigenvalue  $> 0$ . Hence the number of negative eigenvalues is  $\leq d$ , since the real dimension of  $X$  is  $2d$ . Thus the index of the critical point  $a$  is  $\leq d$ .

Now using Lemma 7.3 (b), we see that

$$H_r(U_\beta, \mathbf{Z}) = 0, \quad (\forall r > d).$$

From this it follows that

$$H_r(X, \mathbf{Z}) = 0, \quad (\forall r > d),$$

because the singular cycles defining the homology groups  $H_r(X, \mathbf{Z})$  have compact supports, and any compact subset of  $X$  is contained in some compact set  $K$  with a corresponding  $U_\beta \supset K$ .

A refinement of the above argument leads to the stronger (homotopy) statement:

*Any Stein manifold of (complex) dimension  $d$  has the same homotopy type as a CW complex of (real) dimension  $\leq d$ . (See [6]).*

Moreover, the Lefschetz theorem has an analogue in homology and in homotopy [6]. The latter, for example, asserts that, if  $V, D$  are as in Th. 7.1, then the relative homotopy groups  $\pi_q(V, D) = 0$  for  $q < d$ .

Th. 7.2 has been generalised in various directions. It has a relative analogue (relative to a Runge domain). Further, Th. 7.2 remains true if  $X$  is any Stein space (with singularities) of complex dimension  $d$ , but the corresponding cohomology statement is proved only for some other coefficient groups [5, 7]. Note that in view of the use of Poincaré duality, this does not lead to a Lefschetz theorem for algebraic varieties with singularities.

## REFERENCES

- [1] ANDREOTTI, A. and FRANKEL, Theodore, The Lefschetz theorem on hyperplane sections. *Ann. of Math.* 69 (1959), 713-717.
- [2] GRAUERT, H., Ueber Modifikationen und exzeptionelle analytische Mengen. *Math. Ann.* 146 (1962), 331-368.
- [3] GRAUERT, H., and REMMERT, R., Bilder und Urbilder analytischer Garben. *Ann. of Math.* 68 (1958), 393-443.
- [4] HÖRMANDER, L., *An introduction to complex analysis in several variables*. Van Nostrand, 1966.