

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	15 (1969)
Heft:	1: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	SIMPLE PROOFS OF TWO THEOREMS ON MINIMAL SURFACES
Autor:	Chern, Shiing-shen
Kapitel:	3. Proof of Theorem 2
DOI:	https://doi.org/10.5169/seals-43204

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3. PROOF OF THEOREM 2

Let S^3 be the unit sphere in the euclidean 4-space E^4 . By an orthonormal frame in E^4 is meant an ordered set of vectors e_α , $0 \leq \alpha \leq 3$, satisfying

$$(9) \quad (e_\alpha, e_\beta) = \delta_{\alpha\beta}, \quad 0 \leq \alpha, \beta, \gamma \leq 3,$$

where the left-hand side is the scalar product of the vectors in question. The space of all orthonormal frames can be identified with the group $SO(4)$. We introduce in $SO(4)$ the Maurer-Cartan forms $\omega_{\alpha\beta}$ according to the equations

$$(10) \quad de_\alpha = \sum_\beta \omega_{\alpha\beta} e_\beta$$

or

$$(11) \quad \omega_{\alpha\beta} = (de_\alpha, e_\beta).$$

It follows from (9) that

$$(12) \quad \omega_{\alpha\beta} + \omega_{\beta\alpha} = 0.$$

Exterior differentiation of (10) gives the Maurer-Cartan structure equations of $SO(4)$, which are

$$(13) \quad d\omega_{\alpha\beta} = \sum_\gamma \omega_{\alpha\gamma} \wedge \omega_{\gamma\beta}.$$

There is a fibering

$$(14) \quad SO(4) \rightarrow S^3 = SO(4)/SO(3),$$

with the projection defined by sending the frame $e_0 e_1 e_2 e_3$ to the unit vector e_0 .

Suppose a smooth surface

$$(15) \quad M \rightarrow S^3$$

be described by the vector e_0 . We restrict to frames such that e_3 is the unit normal vector to M at e_0 . There are two choices for e_3 , any one of which is called an orientation of M . Suppose M be oriented. Then the frames are defined up to a rotation of the vectors e_1, e_2 in the tangent plane. In other words, our restricted family of frames is a circle bundle over M , for which the structure equations (13) are valid.

The condition that e_3 is a normal vector at e_0 implies

$$(16) \quad \omega_{03} = 0.$$

Taking its exterior derivative and using (13), we get

$$\omega_{01} \wedge \omega_{13} + \omega_{02} \wedge \omega_{23} = 0.$$

Since M is an immersed surface, we have $\omega_{01} \wedge \omega_{02} \neq 0$ and Cartan's lemma allows us to set

$$(17) \quad \omega_{13} = a\omega_{01} + b\omega_{02}, \quad \omega_{23} = b\omega_{01} + c\omega_{02}.$$

The condition for a minimal surface is the vanishing of the mean curvature:

$$(18) \quad a + c = 0.$$

Let

$$(19) \quad \alpha = \omega_{01} + i\omega_{02}, \quad \beta = \omega_{13} + i\omega_{23}.$$

The structure equations (13) give

$$d\alpha = -i\omega_{12} \wedge \alpha,$$

$$(20) \quad d\beta = -i\omega_{12} \wedge \beta.$$

Under a rotation of e_1 e_2 both α and β will be multiplied by the same complex number of absolute value 1. It follows that

$$(21) \quad \alpha \wedge \bar{\beta}, \quad \alpha \bar{\beta},$$

which are exterior and ordinary two-forms respectively, are globally defined on our oriented surface M .

Suppose from now on that M is a minimal surface. Condition (18) can be written

$$(22) \quad \beta = A\bar{\alpha}, \quad A = a + ib.$$

In this case the first form in (21) vanishes identically, while

$$(23) \quad \alpha \bar{\beta} = \bar{A} \alpha^2.$$

Taking the exterior derivative of (22) and using (20), we get

$$(24) \quad dA + 2iA\omega_{12} \equiv 0, \quad \text{mod } \bar{\alpha}.$$

The induced riemannian metric on M has an underlying complex structure which makes M into a Riemann surface. We wish to show that

the form in (23) is a quadratic differential in the sense of Riemann surfaces. For this purpose let z be a local complex coordinate on M , so that

$$(25) \quad \alpha = \lambda dz .$$

Then we have, locally,

$$\alpha\bar{\beta} = (\lambda^2 \bar{A}) dz^2 .$$

Exterior differentiation of (25) and use of (20) give

$$d\lambda + i\lambda\omega_{12} \equiv 0 , \quad \text{mod } dz .$$

Combining with (24), we get

$$\frac{\partial}{\partial \bar{z}} (\lambda^2 \bar{A}) = 0$$

i.e., the coefficient of dz^2 in $\alpha\bar{\beta}$ is holomorphic.

Since M is of genus zero, the quadratic differential must vanish. This implies $A = 0$ and that M is a great sphere.

The proof given above is not essentially different from those of Almgren and Calabi. The main idea of using the quadratic differential in surface theory goes back to H. Hopf. The formalism developed in this proof should also be useful in the study of other problems on surfaces in S^3 .

4. A FORMULA ON NON-PARAMETRIC MINIMAL HYPERSURFACES IN EUCLIDEAN SPACE

Instead of proving formula (2) we will establish a more general formula for a non-parametric minimal hypersurface in the euclidean $(n+1)$ -space E^{n+1} , which seems to have an independent interest.

Suppose $x: M \rightarrow E^{n+1}$ be an immersion of an n -dimensional manifold M in E^{n+1} . We consider orthonormal frames $x e_1 \dots e_{n+1}$ in E^{n+1} , such that $x \in M$ and e_{n+1} is the unit normal vector to M at x . We have then

$$(26) \quad \begin{aligned} dx &= \sum_i \omega_i e_i , \\ de_i &= \sum_k \omega_{ik} e_k + \omega_{i,n+1} e_{n+1} , \quad 1 \leq i, j, k, l \leq n , \\ de_{n+1} &= - \sum_i \omega_{i,n+1} e_i , \end{aligned}$$