

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	15 (1969)
Heft:	1: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	ONE-SIDED ANALOGUES OF KARAMATA's REGULAR VARIATION
Autor:	Feller, William
Kapitel:	4. Other conditions
DOI:	https://doi.org/10.5169/seals-43209

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

4. OTHER CONDITIONS

By theorem 1 both U and U_p vary dominatedly whenever $r > 0$ and $\bar{r} < \infty$. The next theorem gives even simpler criteria for dominated variation that remain applicable in the limiting situations $r = 0$ and $\bar{r} = \infty$.

THEOREM 3. *If there exists a number $\xi > 1$ such that*

$$(4.1) \quad \liminf \frac{U(\xi t)}{U(t)} > 1$$

then U_p varies dominatedly. Similarly the relation

$$(4.2) \quad \liminf \frac{U_p(t/\xi)}{U_p(t)} > 1$$

implies the dominated variation of U .

PROOF. Clearly

$$(4.3) \quad \frac{t^p U_p(t)}{U(t)} \geq \frac{t^p [U_p(t) - U_p(t\xi)]}{U(t)} \geq \xi^{-p} \frac{U(t\xi) - U(t)}{U(t)}.$$

When the right side is bounded away from 0 this implies $r > 0$, and so U_p varies dominatedly by theorem 1.

We can go a step further. If, besides (4.1), it is known that U varies dominatedly with exponent $\gamma < p$, then $t^{-p} U(t)/U_p(t)$ is bounded away from 0, and hence the second inequality in (4.3) implies that

$$(4.4) \quad \liminf \frac{U_p(t) - U_p(t\xi)}{U_p(t)} > 0.$$

This is equivalent to (4.2).

Similarly

$$(4.5) \quad \begin{aligned} \frac{U_p(t) - U_p(t\xi)}{U_p(t\xi)} &\leq t^{-p} \frac{U(t\xi) - U(t)}{U_p(t\xi)} = \\ &= \frac{\xi^p}{R_U(t\xi)} \frac{U(t\xi) - U(t)}{U(t\xi)}. \end{aligned}$$

The second fraction on the right does not exceed 1, and so (4.2) ensures that $R_U(t\xi)$ remains bounded, and hence that U is of dominated variation.

Again, if it is known that R_U is bounded away from 0 then (4.5) shows that (4.2) implies (4.1).

We have thus proved the

COROLLARY. *If U is of dominated variation with exponent $\gamma < p$ then (4.1) implies (4.2). Similarly, if U_p is of dominated variation with exponent $-q$ where $q < p$, then (4.2) entails (4.1). (In each case both functions are of dominated variation.)*

5. RATIO LIMIT THEOREMS

Let U and V be non-decreasing unbounded functions, and suppose that L is slowly varying (= regularly varying with exponent 0).

DEFINITION. *We shall say that U and V are L -equivalent and write*

$$(5.1) \quad V \leftrightarrow UL$$

if the ratio UL/V tends to 1 at all points of continuity.

More precisely, it is required that for each $\varepsilon > 0$ and fixed $\lambda > 1$

$$(5.2) \quad (1 - \varepsilon) L(t) U(t/\lambda) \leq V(t) \leq (1 + \varepsilon) L(t) U(t\lambda)$$

for all t sufficiently large.

THEOREM 4. *Let U be of dominated variation. In order that there exist a slowly varying function L such that (5.1) holds it is necessary and sufficient that*

$$(5.3) \quad R_U(t) - R_V(t) \rightarrow 0 \quad \text{boundedly.}$$

Needless to say, R_V and \mathcal{J}_V are defined by analogy with R_U in (1.5) and \mathcal{J}_U in (3.2).

PROOF. (a) *Necessity.* Assume (5.1) and suppose that U satisfies the basic inequality (2.2). Obviously the slow variation of L implies that for t sufficiently large and all $x > 1$

$$(5.4) \quad \frac{V(tx)}{V(t)} < C' x^{\gamma'}$$

for any pair of constants $C' > C$ and $\gamma' > \gamma$. Thus V is of dominated variation, and since $p > \gamma$ the function V_p exists.