

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 15 (1969)  
**Heft:** 1: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ASYMPTOTIC BEHAVIOUR OF A CLASS OF DETERMINANTS  
**Autor:** Kac, Mark  
**DOI:** <https://doi.org/10.5169/seals-43216>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# ASYMPTOTIC BEHAVIOUR OF A CLASS OF DETERMINANTS

by Mark KAC

*To the memory of J. Karamata*

1° Let  $f(\theta)$ ,  $0 < \theta < 1$ , be a function which in addition to certain regularity conditions, which will be specified later on, is bounded from below by a number larger than 2. In other words,

$$(1.1) \quad f(\theta) > m > 2, \quad 0 < \theta < 1.$$

Consider the determinant

$$(1.2) \quad D_n(f) = \begin{vmatrix} f\left(\frac{1}{n}\right), & -1, & 0, & \dots & \dots & 0 \\ -1, & f\left(\frac{2}{n}\right), & -1, & \dots & \dots & 0 \\ 0, & -1, & f\left(\frac{3}{n}\right), & -1, & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0, & 0, & \dots & -1, & f\left(\frac{n}{n}\right) & \end{vmatrix}$$

i.e. the determinant of the matrix

$$(1.3) \quad a_{ij} = f\left(\frac{i}{n}\right) \delta_{ij} - \delta_{i,j-1} - \delta_{i,j+1}$$

where  $\delta$ , as usual, denotes the Kronecker delta.

If  $f(\theta)$  is Riemann integrable it can be shown that

$$(1.4) \quad \lim_{n \rightarrow \infty} D_n^{\frac{1}{n}}(f) = \exp \left\{ \frac{1}{2\pi} \int_{-\pi}^{\pi} d\omega \int_0^1 \log (f(\theta) - 2 \cos \omega) d\theta \right\} = G(f).$$

This is a special case of a more general result of Kac, Murdock and Szegő [1953].

Assuming that  $f(\theta)$  is twice differentiable, with a bounded second derivative, one can go a good deal farther.

In fact, one can determine

$$(1.5) \quad \lim_{n \rightarrow \infty} \frac{D_n(f)}{G^n(f)}$$

with  $G(f)$  as defined in (1.4).

Although the result is again a special case of a more general result obtained by Mejbo and Schmidt [1962] the method which we shall use is quite different and of some interest in itself. The method has also the advantage of explaining the role of smoothness conditions imposed on  $f$ .

2° We begin with the elementary formula

$$(2.1) \quad D_n^{-\frac{1}{2}}(f) = \left( \frac{1}{(\sqrt{\pi})^n} \right)^n \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \exp \left\{ - \sum_1^n f\left(\frac{k}{n}\right) x_k^2 + 2 \sum_1^{n-1} x_k x_{k+1} \right\} dx_1 dx_2 \dots dx_n$$

and note that setting

$$(2.2) \quad K\left(x, y; \frac{l}{n}\right) = \frac{1}{\sqrt{\pi}} \exp \left\{ - \frac{1}{2} f\left(\frac{l}{n}\right) x^2 + 2xy - \frac{1}{2} f\left(\frac{l+1}{n}\right) y^2 \right\}$$

we can rewrite (2.1) in the form

$$(2.3) \quad D_n^{-\frac{1}{2}}(f) = \frac{1}{(\sqrt{\pi})^n} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \exp \left\{ - \frac{1}{2} f\left(\frac{1}{n}\right) x_1^2 \right\} K\left(x_1, x_2; \frac{1}{n}\right) K\left(x_2, x_3; \frac{2}{n}\right) \dots \\ \dots K\left(x_{n-1}, x_n; \frac{n}{n}\right) \exp \left\{ - \frac{1}{2} f\left(\frac{n}{n}\right) x_n^2 \right\} dx_1 dx_2 \dots dx_n.$$

The kernels  $K(x, y; \frac{l}{n})$  are not symmetric and we must replace them by symmetric ones without introducing too large an error.

This is done by noting that

$$(2.4) \quad f\left(\frac{l}{n}\right) = \frac{1}{2} f\left(\frac{l-1}{n}\right) + \frac{1}{4n} f'\left(\frac{l-1}{n}\right) + \frac{1}{2} f\left(\frac{l}{n}\right) + \frac{1}{4n} f'\left(\frac{l}{n}\right) + O\left(\frac{1}{n^2}\right)$$

where the error  $O(1/n^2)$  is actually less than  $M/n^2$  where  $M$  is a bound for  $f''(\theta)$  in  $(0, 1)$ .

It follows that

$$(2.5) \quad \left| \sum_{l=1}^n f\left(\frac{l}{n}\right) x_l^2 - \left\{ \left( \frac{1}{2}f(0) + \frac{1}{4n}f'(0) \right) x_1^2 + \right. \right. \\ \left. \left. + \sum_{l=1}^{n-1} \left[ \left( \frac{1}{2}f\left(\frac{l}{n}\right) + \frac{1}{4n}f'\left(\frac{l}{n}\right) \right) x_l^2 + \left( \frac{1}{2}f\left(\frac{l}{n}\right) + \frac{1}{4n}f'\left(\frac{l}{n}\right) x_{l+1}^2 \right] \right. \right. \\ \left. \left. + \left( \frac{1}{2}f(1) + \frac{1}{4n}f'(1) \right) x_n^2 \right\} \right| < \frac{M}{n^2} \sum_1^n x_l^2$$

and this suggests that we introduce the symmetric kernels

$$(2.6) \quad \tilde{K}_\alpha\left(x, y; \frac{l}{n}\right) = \frac{1}{\sqrt{\pi}} \exp \left\{ - \left( \frac{1}{2}f\left(\frac{l}{n}\right) + \frac{1}{4n}f'\left(\frac{l}{n}\right) + \frac{\alpha}{2n^2} \right) x^2 + 2xy \right. \\ \left. - \left( \frac{1}{2}f\left(\frac{l}{n}\right) + \frac{1}{4n}f'\left(\frac{l}{n}\right) + \frac{\alpha}{2n^2} \right) y^2 \right\}.$$

It should now be clear from (2.3) and (2.5) that  $D_n^{-\frac{1}{2}}(f)$  is contained between the value of the integral

$$(2.7) \quad \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \exp \left\{ - \left( \frac{1}{2}f(0) + \frac{1}{4n}f'(0) \right) x_1^2 \right\} \prod_{l=1}^{n-1} K_\alpha\left(x_l, x_{l+1}; \frac{l}{n}\right) \\ \exp \left\{ - \left( \frac{1}{2}f(1) + \frac{1}{4n}f'(1) \right) x_n^2 \right\} dx_1 \dots dx_n$$

for  $\alpha = -M$  and its value for  $\alpha = M$ .

3° It will turn out that the asymptotic behaviour of (2.7), to the required accuracy, does not depend on  $\alpha$  so that we may as well set  $\alpha = 0$ .

The kernels  $K_0(x, y; l/n)$  do not commute. However they “almost commute” in the following sense:

Let  $\lambda_1(l/n), \lambda_2(l/n), \dots$  be the eigenvalues of  $K_0(x, y; l/n)$  in the decreasing order and let

$$\varphi_1(x; l/n), \quad \varphi_2(x; l/n), \dots$$

be the corresponding normalized eigenfunctions.

Then

$$(3.1) \quad \int_{-\infty}^{\infty} \varphi_i\left(x; \frac{l}{n}\right) \varphi_j\left(x; \frac{l+1}{n}\right) dx = \delta_{ij} + O\left(\frac{1}{n^2}\right).$$

This follows by a perturbation argument but can also be obtained directly since the eigenfunctions and eigenvalues of our kernels can be determined explicitly.

I shall now ask the reader to believe me that whenever we have (3.1) and whenever  $\lambda_1(l/n) - \lambda_2(l/n)$  is bounded away from 0 (uniform non-degeneracy) the integral (2.7) divided by

$$(3.2) \quad \prod_{l=1}^{n-1} \lambda_1\left(\frac{l}{n}\right)$$

approaches

(3.3)

$$\pi^{-\frac{1}{2}} \int_{-\infty}^{\infty} \varphi_1(x; 0) \exp\left\{-\frac{1}{2}f(0)x^2\right\} dx \int_{-\infty}^{\infty} \varphi_1(x; 1) \exp\left\{-\frac{1}{2}f(1)x^2\right\} dx .$$

In other words, the asymptotic behaviour of the integral (2.7) is the same *as if* the kernels  $K_0(x, y; l/n)$  commuted. The details of the proof are tedious but intuitively the assertions should be nearly obvious.

It remains now to calculate  $\lambda_1(l/n)$  and  $\varphi_1(x; l/n)$ .

The kernels  $K_0$  are of the form

$$(3.4) \quad \frac{1}{\sqrt{\pi}} \exp\left\{-\frac{a}{2}x^2 + 2xy - \frac{a}{2}y^2\right\}$$

with  $a > 2$  and it is easy to convince oneself that the principal eigenfunction is of the form

$$(3.5) \quad \frac{\sqrt[4]{b}}{\sqrt[4]{\pi}} \exp\left\{-\frac{b}{2}y^2\right\}$$

A simple calculation then gives

$$(3.6) \quad b = \sqrt{a^2 - 4}$$

and for the maximum eigenvalue  $\lambda_1(a)$  the formula

$$(3.7) \quad \lambda_1(a) = \left(\frac{a}{2} + \sqrt{\left(\frac{a}{2}\right)^2 - 1}\right)^{-\frac{1}{2}} .$$

Setting

$$(3.8) \quad a_l = f\left(\frac{l}{n}\right) + \frac{1}{2n}f'\left(\frac{l}{n}\right)$$

we finally obtain that

$$(3.9) \quad \lim_{n \rightarrow \infty} \frac{D_n(f)}{\prod_{l=1}^{n-1} \left( \frac{a_l}{2} + \sqrt{\left(\frac{a_l}{2}\right)^2 - 1} \right)} = \frac{f(0) + \sqrt{f^2(0) - 4}}{2 \sqrt[4]{f^2(0) - 4}} \cdot \frac{f(1) + \sqrt{f^2(1) - 4}}{2 \sqrt[4]{f^2(1) - 4}}.$$

It remains to determine the asymptotic behavior of the product

$$(3.10) \quad \prod_{l=1}^{n-1} \left( \frac{a_l}{2} + \sqrt{\left(\frac{a_l}{2}\right)^2 - 1} \right) = \exp \sum_{l=1}^{n-1} \log \left( \frac{a_l}{2} + \sqrt{\left(\frac{a_l}{2}\right)^2 - 1} \right).$$

Neglecting terms of order  $1/n^2$  and higher we find easily that

$$\log \left( \frac{a_l}{2} + \sqrt{\left(\frac{a_l}{2}\right)^2 - 1} \right) = \log \frac{f\left(\frac{l}{n}\right) + \sqrt{f^2\left(\frac{l}{n}\right) - 4}}{2} + \frac{1}{2n} \frac{f'\left(\frac{l}{n}\right)}{\sqrt{f^2\left(\frac{l}{n}\right) - 4}}$$

and hence

$$(3.11) \quad \begin{aligned} & \sum_{l=1}^{n-1} \log \left( \frac{a_l}{2} + \sqrt{\left(\frac{a_l}{2}\right)^2 - 1} \right) = \\ & \sum_{l=1}^{n-1} \log \frac{f\left(\frac{l}{n}\right) + \sqrt{f^2\left(\frac{l}{n}\right) - 4}}{2} + \frac{1}{2} \int_0^1 \frac{f'(\theta)}{\sqrt{f^2(\theta) - 4}} d\theta + o\left(\frac{1}{n}\right). \end{aligned}$$

Using the well known formula

$$(3.12) \quad \sum_{l=1}^{n-1} g\left(\frac{l}{n}\right) = n \int_0^1 g(\theta) d\theta - \frac{g(0) + g(1)}{2} + o\left(\frac{1}{n}\right)$$

for

$$(3.13) \quad g(\theta) = \log \frac{f(\theta) + \sqrt{f^2(\theta) - 4}}{2}$$

and noting that

$$\int_0^1 \frac{f'(\theta)}{\sqrt{f^2(\theta) - 4}} d\theta = \log \frac{f(1) + \sqrt{f^2(1) - 4}}{2} - \log \frac{f(0) + \sqrt{f^2(0) - 4}}{2}$$

we obtain

$$(3.14) \quad \sum_{l=1}^{n-1} \log \left( \frac{a_l}{2} + \sqrt{\left( \frac{a_l}{2} \right)^2 - 1} \right) = \\ n \int_0^1 \log \frac{f(\theta) + \sqrt{f^2(\theta) - 4}}{2} d\theta - \log \frac{f(0) + \sqrt{f^2(0) - 4}}{2} + O\left(\frac{1}{n}\right).$$

Combining (3.14) with (3.9) and noting that

$$\log G(f) = \frac{1}{2\pi} \int_0^1 d\theta \int_{-\pi}^{\pi} \log (f(\theta) - 2 \cos \omega) d\omega = \\ \int_0^1 \log \frac{f(\theta) + \sqrt{f^2(\theta) - 4}}{2} d\theta$$

we are led to the final result that

$$(3.15) \quad \lim_{n \rightarrow \infty} \frac{D_n(f)}{G^n(f)} = \frac{f(1) + \sqrt{f^2(1) - 4}}{\sqrt[4]{f^2(0) - 4} \sqrt[4]{f^2(1) - 4}}.$$

4° If we set

$$(4.1) \quad \log (f(\theta) - 2 \cos \omega) = \sum_{-\infty}^{\infty} h_v(\theta) e^{iv\omega}$$

the right hand side of (3.15) can be shown to be equal to

$$(4.2) \quad \exp \left\{ \frac{1}{2} (h_0(1) - h_0(0)) \right\} \exp \left\{ \frac{1}{2} \sum_1^{\infty} v h_v^2(0) + \frac{1}{2} \sum_1^{\infty} v h_v^2(1) \right\}.$$

This is in apparent disagreement with the formula of Mejbo and Schmidt since in their formula  $h_0(1) - h_0(0)$  is replaced by  $h_0(0) - h_0(1)$ . We would get complete agreement if instead of our determinant (1.2) we would consider a modified determinant with  $f(0), f(\frac{1}{n}), f(\frac{2}{n}), \dots, f(\frac{n-1}{n})$  on the main diagonal. So accurate is formula (3.15) that it is sensitive to so slight a modification!

REFERENCES

KAC, M., W. L. MURDOCK and G. SZEGÖ, On the eigenvalues of certain Hermitian forms.  
*J. Rat. Mech. Anal.*, 2 (1953), pp. 767-800.

MEJLBO, Lars and Palle SCHMIDT, On the eigenvalues of generalized Toeplitz matrices.  
*Math. Scand.*, 10 (1952), pp. 5-16.

(Reçu le 1<sup>er</sup> mai 1968)

Mark Kac  
Rockefeller University  
New-York, N.Y. 10021

**vide-leer-empty**