

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 15 (1969)  
**Heft:** 1: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE DEGREE OF CONVERGENCE OF FEJÉR-LEBESGUE SUMS  
**Autor:** Aljani, S. / Bojanic, R. / Tomi, M.  
**DOI:** <https://doi.org/10.5169/seals-43201>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 03.03.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# ON THE DEGREE OF CONVERGENCE OF FEJÉR-LEBESGUE SUMS

S. ALJANČIĆ, R. BOJANIC and M. TOMIĆ

*To the memory of J. Karamata*

1. Let  $f$  be a  $2\pi$ -periodic and  $L$ -integrable function and  $(\sigma_n[f])$  the sequence of Fejér-Lebesgue sums of the Fourier series of  $f$ , defined by

$$\sigma_n[f](x) = \frac{1}{2\pi(n+1)} \int_{-\pi}^{\pi} f(t) \left( \frac{\sin \frac{1}{2}(n+1)(x-t)}{\sin \frac{1}{2}(x-t)} \right)^2 dt .$$

The classical result of Fejér [1] states that if  $f$  is a  $2\pi$ -periodic, bounded and  $R$ -integrable function, continuous at the point  $x$ , then

$$(1.1) \quad \lim_{n \rightarrow \infty} \sigma_n[f](x) = f(x)$$

and the convergence is uniform on any closed interval where  $f$  is continuous. In particular, if  $f$  is a  $2\pi$ -periodic and continuous function, then

$$(1.2) \quad \|\sigma_n[f] - f\| = \sup_{-\infty < x < \infty} |\sigma_n[f](x) - f(x)| \rightarrow 0 \quad (n \rightarrow \infty) .$$

Fejér's result has been generalized by Lebesgue [2] who has proved that (1.1) holds whenever  $f$  is a  $2\pi$ -periodic and  $L$ -integrable function and

$$(1.3) \quad \int_0^h |f(x+u) - f(x)| du = o(h) \quad (h \rightarrow 0) .$$

At the same time, Lebesgue has shown that this condition is satisfied almost everywhere. In particular, (1.3) holds at each point  $x$  where  $f$  is continuous.

For  $2\pi$ -periodic and continuous functions more precise versions of Fejér's result (1.2) usually give estimate of the rate of convergence of the sequence  $(\sigma_n[f])$  to  $f$  either in terms of the modulus of continuity  $\omega_f$  which is defined by

$$\omega_f(h) = \sup \{ |f(x') - f(x'')| : |x' - x''| \leq h \}$$

or in terms of the best approximation  $E_n^T(f)$  which is defined by

$$E_n^T(f) = \inf \{ \|f - t\| : t \in T_n \}$$

where  $T_n$  is the set of all trigonometric polynomials of degree  $\leq n$ . The following results are known.

If  $f$  is a  $2\pi$ -periodic and continuous function such that  $\omega_f(h) h^{-\eta}$  is decreasing for some  $\eta \in (0, 1)$ , then

$$(1.4) \quad \|\sigma_n[f] - f\| \leq A\omega_f\left(\frac{1}{n}\right)$$

(see [3], Vol. 1, Ch. 3, Th. 3.16). For arbitrary  $2\pi$ -periodic and continuous  $f$  we have the inequality

$$(1.5) \quad \|\sigma_n[f] - f\| \leq B\omega_f\left(\frac{\log n}{n}\right) \quad (n \geq 2)$$

which in a somewhat weaker form was obtained by D. Jackson (see [4], p. 64), and the inequality

$$(1.6) \quad \|\sigma_n[f] - f\| \leq \frac{C}{n+1} \sum_{k=0}^n \omega_f\left(\frac{1}{k+1}\right)$$

which was obtained recently by S. B. Stečkin [5]. Stečkin's inequality follows from a still more precise inequality

$$\|\sigma_n[f] - f\| \leq \frac{12}{n+1} \sum_{k=0}^n E_{k+1}^T(f)$$

and Jackson's theorem which states that  $E_n^T(f) \leq 12\omega_f(1/n)$ . Using the well known properties of the modulus of continuity it is easy to see that Stečkin's inequality (1.6) is more precise than (1.5).

The aim of this paper is to obtain the analog of Stečkin's inequality (1.6) for arbitrary  $2\pi$ -periodic and  $L$ -integrable functions and to show that this result cannot be improved for certain classes of functions. As a measure of deviation of  $\sigma_n[f](x)$  from  $f(x)$  we shall take the function  $w[f, x]$  defined by

$$w[f, x](h) = \sup \left\{ (2t)^{-1} \int_{-t}^t |f(x+u) - f(x)| du : 0 < t \leq h \right\}.$$

Clearly,  $w[f, x]$  is a non-decreasing, real valued function and, by (1.3),  $w[f, x](h) \rightarrow 0$  ( $h \rightarrow 0$ ) almost everywhere. If  $f$  is a continuous function we have for  $0 < t \leq h$

$$\int_{-t}^t |f(x+u) - f(x)| du \leq \int_{-t}^t \omega_f(|u|) du \leq 2t\omega_f(h)$$

and consequently

$$(1.7) \quad w[f, x](h) \leq \omega_f(h).$$

Our first result can be stated as follows.

**THEOREM 1.** *Let  $f$  be a  $2\pi$ -periodic and  $L$ -integrable function. Then we have for all  $n \geq 0$  the inequality*

$$(1.8) \quad |\sigma_n[f](x) - f(x)| \leq \frac{3}{n+1} \sum_{k=0}^n w[f, x]\left(\frac{\pi}{k+1}\right).$$

For continuous  $2\pi$ -periodic functions in view of (1.7) we obtain immediately from (1.8) Stečkin's inequality

$$\|\sigma_n[f] - f\| \leq \frac{3}{n+1} \sum_{k=0}^n \omega_f\left(\frac{\pi}{k+1}\right).$$

From Theorem 1 we can obtain similar results valid for all functions in certain classes of  $2\pi$ -periodic and  $L$ -integrable functions. We shall consider here two classes of these functions generated by a non-negative and increasing real valued function  $\Omega$  defined on  $[0, \pi]$  with  $\Omega(0) = 0$ . The first class  $L(\Omega)$  consists of all  $2\pi$ -periodic and  $L$ -integrable functions  $f$  such that

$$A_f = \sup_{0 < h \leq \pi} \frac{w[f, x](h)}{\Omega(h)} < \infty.$$

The second, somewhat larger class  $L^*(\Omega)$  consists of all  $2\pi$ -periodic and  $L$ -integrable functions  $f$  such that

$$B_f = \limsup_{h \rightarrow 0+} \frac{w[f, x](h)}{\Omega(h)} < \infty.$$

From (1.8) follows immediately that for any  $f \in L(\Omega)$  and  $n = 0, 1, 2, \dots$  we have

$$|\sigma_n[f](x) - f(x)| \leq \frac{3A_f}{n+1} \sum_{k=0}^n \Omega\left(\frac{\pi}{k+1}\right)$$

and so

$$(1.9) \quad \sup_{n \geq 0} \left( \frac{|\sigma_n[f](x) - f(x)|}{\frac{1}{n+1} \sum_{k=0}^n \Omega\left(\frac{\pi}{k+1}\right)} \right) \leq 3 \sup_{0 < h \leq \pi} \frac{w[f, x](h)}{\Omega(h)}.$$

On the other hand, if  $\sum \Omega(\pi/k+1)$  is a divergent series we have for every  $f \in L^*(\Omega)$  the inequality

$$(1.10) \quad \limsup_{n \rightarrow \infty} \frac{|\sigma_n[f](x) - f(x)|}{\frac{1}{n+1} \sum_{k=0}^n \Omega\left(\frac{\pi}{k+1}\right)} \leq 3 \limsup_{h \rightarrow 0+} \frac{w[f, x](h)}{\Omega(h)}.$$

In order to show how (1.10) can be obtained from (1.8), observe that for  $k \geq N'_f$  we have  $w[f, x](\pi/k+1) \leq (B_f + \varepsilon) \Omega(\pi/k+1)$ . Consequently, from (1.8) follows that for  $n \geq N'_f$

$$|\sigma_n[f](x) - f(x)| \leq \frac{3}{n+1} \sum_{k=0}^{N'_f} w[f, x]\left(\frac{\pi}{k+1}\right) + \frac{3(B_f + \varepsilon)}{n+1} \sum_{k=N'_f+1}^n \Omega\left(\frac{\pi}{k+1}\right).$$

Since  $\sum \Omega(\pi/k+1)$  is a divergent series, we can find  $N''_f$  such that for  $n > N''_f$

$$\sum_{k=0}^{N'_f} w[f, x]\left(\frac{\pi}{k+1}\right) \leq \varepsilon \sum_{k=0}^n \Omega\left(\frac{\pi}{k+1}\right).$$

Hence, for  $n > \max(N'_f, N''_f)$  we have

$$|\sigma_n[f](x) - f(x)| \leq \frac{3(B_f + 2\varepsilon)}{n+1} \sum_{k=0}^n \Omega\left(\frac{\pi}{k+1}\right)$$

and (1.10) follows.

In order to show that the inequality (1.8) cannot be essentially improved we shall consider the class  $L_M(\Omega) \subseteq L(\Omega)$  of  $2\pi$ -periodic and  $L$ -integrable functions  $f$  such that

$$A_f = \sup_{0 < h \leq \pi} \frac{w[f, x](h)}{\Omega(h)} \leq M.$$

We have then the following result.

**THEOREM 2.** *There exists  $m$  ( $0 < m < M$ ) such that for all  $n \geq 2$  we have the inequalities*

$$(1.11) \quad \frac{m}{n+1} \sum_{k=2}^n \Omega\left(\frac{\pi}{k+1}\right) \leq \sup_{f \in L_M(\Omega)} |\sigma_n[f](x) - f(x)| \leq \leq \frac{3M}{n+1} \sum_{k=0}^n \Omega\left(\frac{\pi}{k+1}\right).$$

2. *Proof of Theorem 1.* The following proof of Theorem 1 is based on a small modification of the classical proof of the Fejér-Lebesgue's theorem (see [3], Vol. 1, Ch. 3, Th. 3.9).

Let  $0 \leq t \leq \pi$  and

$$F_x(t) = \int_0^t |f(x+u) + f(x-u) - 2f(x)| du.$$

We have then

$$(2.1) \quad F_x(t) \leq \int_{-t}^t |f(x+u) - f(x)| du \leq 2tw[f, x](t).$$

Since

$$\begin{aligned} \sigma_n[f](x) - f(x) &= \\ &= \frac{1}{2\pi(n+1)} \int_0^\pi (f(x+t) + f(x-t) - 2f(x)) \left( \frac{\sin \frac{1}{2}(n+1)t}{\sin \frac{1}{2}t} \right)^2 dt \end{aligned}$$

we have

$$\begin{aligned} |\sigma_n[f](x) - f(x)| &\leq \\ &\leq \frac{1}{2\pi(n+1)} \left( \int_0^{\pi/n+1} + \int_{\pi/n+1}^\pi \right) |f(x+t) + f(x-t) - 2f(x)| \left( \frac{\sin \frac{1}{2}(n+1)t}{\sin \frac{1}{2}t} \right)^2 dt \\ &\leq P_n + Q_n. \end{aligned}$$

Since  $|\sin nt| \leq n |\sin t|$  we have

$$(2.2) \quad \begin{aligned} P_n &\leq \frac{1}{2\pi} (n+1) \int_0^{\pi/n+1} |f(x+t) + f(x-t) - 2f(x)| dt = \\ &= \frac{1}{2\pi} (n+1) F_x \left( \frac{\pi}{n+1} \right). \end{aligned}$$

Next, since  $\sin \frac{1}{2}t \geq t/\pi$  for  $t \in [0, \pi]$ , we have

$$Q_n \leq \frac{\pi}{2(n+1)} \int_{\pi/n+1}^\pi |f(x+t) + f(x-t) - 2f(x)| \frac{dt}{t^2}.$$

Using partial integration we find that

$$(2.3) \quad Q_n \leq \frac{1}{2\pi(n+1)} F_x(\pi) - \frac{1}{2\pi}(n+1) F_x\left(\frac{\pi}{n+1}\right) + \\ + \frac{\pi}{n+1} \int_{\pi/n+1}^{\pi} F_x(t) \frac{dt}{t^3}.$$

Adding (2.2) and (2.3) and using (2.1) we find that

$$(2.4) \quad |\sigma_n[f](x) - f(x)| \leq \frac{1}{n+1} w[f, x](\pi) + \\ + \frac{2\pi}{n+1} \int_{\pi/n+1}^{\pi} w[f, x](t) \frac{dt}{t^2}.$$

Since  $w[f, x]$  is non-negative and non-decreasing we have

$$w[f, x](\pi) \leq \sum_{k=0}^n w[f, x](\pi/k + 1)$$

and

$$\int_{\pi/n+1}^{\pi} w[f, x](t) \frac{dt}{t^2} = \frac{1}{\pi} \int_1^{n+1} w[f, x]\left(\frac{\pi}{t}\right) dt \leq \frac{1}{\pi} \sum_{k=0}^n w[f, x]\left(\frac{\pi}{k+1}\right)$$

and (1.8) follows from the last two inequalities and (2.4).

*Proof of Theorem 2.* The right hand side inequality (1.11) follows immediately from (1.9). To prove the left hand side inequality (1.11) let  $g_x(t) = M \Omega(|t-x|)$  if  $|t-x| \leq \pi$  and  $g_x(t+2\pi) = g_x(t)$ . Then  $g_x$  is clearly a  $2\pi$ -periodic and  $L$ -integrable function. Since for  $0 < t \leq h \leq \pi$  we have

$$\int_{-t}^t |g_x(x+u) - g_x(x)| du = M \int_{-t}^t \Omega(|u|) du \leq 2t M \Omega(h),$$

it follows that  $w[g_x, x](h) \leq M \Omega(h)$  and so  $g_x \in L_M(\Omega)$ .

Next,

$$(2.5) \quad \sup_{f \in L_M(\Omega)} |\sigma_n[f](x) - f(x)| \geq |\sigma_n[g_x](x) - g_x(x)| = \sigma_n[g_x](x).$$

We have first

$$\sigma_n[g_x](x) = \frac{1}{2\pi(n+1)} \int_0^{\pi} (g_x(x+t) + g_x(x-t)) \left( \frac{\sin \frac{1}{2}(n+1)t}{\sin \frac{1}{2}t} \right)^2 dt$$

$$\begin{aligned}
 &= \frac{M}{\pi(n+1)} \int_0^{\pi} \Omega(t) \left( \frac{\sin \frac{1}{2}(n+1)t}{\sin \frac{1}{2}t} \right)^2 dt \\
 &\geq \frac{4M}{\pi(n+1)} \int_{\frac{\pi}{n+1}}^{\pi} \Omega(t) \left( \frac{\sin \frac{1}{2}(n+1)t}{\sin \frac{1}{2}t} \right)^2 dt.
 \end{aligned}$$

Since  $|\sin x| \leq |x|$  it follows that

$$(2.6) \quad \sigma_n[g_x](x) \geq \frac{4M}{\pi(n+1)} \int_{\frac{\pi}{n+1}}^{\pi} \Omega(t) (\sin \frac{1}{2}(n+1)t)^2 \frac{dt}{t^2} = A_n.$$

Since  $\Omega$  is non-decreasing, we have

$$\begin{aligned}
 A_n &= \frac{4M}{\pi(n+1)} \sum_{k=1}^n \int_{\frac{k\pi}{n+1}}^{\frac{(k+1)\pi}{n+1}} \Omega(t) (\sin \frac{1}{2}(n+1)t)^2 \frac{dt}{t^2} \\
 &\geq \frac{4M}{\pi(n+1)} \sum_{k=1}^n \Omega\left(\frac{k\pi}{n+1}\right) \left(\frac{n+1}{k+1}\right)^2 \frac{1}{\pi^2} \int_{\frac{k\pi}{n+1}}^{\frac{(k+1)\pi}{n+1}} (\sin \frac{1}{2}(n+1)t)^2 dt.
 \end{aligned}$$

Since

$$\int_{\frac{k\pi}{n+1}}^{\frac{(k+1)\pi}{n+1}} (\sin \frac{1}{2}(n+1)t)^2 dt = \frac{\pi}{2(n+1)},$$

we find that for  $n \geq 2$

$$A_n \geq 2M\pi^{-2} \sum_{k=1}^n \frac{1}{(k+1)^2} \Omega\left(\frac{k\pi}{n+1}\right) \geq 2M\pi^{-2} \sum_{k=2}^n \frac{1}{(k+1)^2} \Omega\left(\frac{k\pi}{n+1}\right).$$

Next, for  $2 \leq k \leq n$

$$\int_{\frac{(k-1)\pi}{n+1}}^{\frac{k\pi}{n+1}} \Omega(t) \frac{dt}{t^2} \leq \frac{n+1}{\pi} \cdot \frac{1}{k(k-1)} \Omega\left(\frac{k\pi}{n+1}\right)$$

and so

$$\frac{1}{(k+1)^2} \Omega\left(\frac{k\pi}{n+1}\right) \geq \frac{\pi}{n+1} \frac{k(k-1)}{(k+1)^2} \int_{\frac{(k-1)\pi}{n+1}}^{\frac{k\pi}{n+1}} \Omega(t) \frac{dt}{t^2} \geq \frac{\pi}{5(n+1)} \int_{\frac{(k-1)\pi}{n+1}}^{\frac{k\pi}{n+1}} \Omega(t) \frac{dt}{t^2}.$$

Hence, for  $n \geq 2$ ,

$$A_n \geq \frac{2M}{5\pi(n+1)} \sum_{k=2}^n \int_{\frac{(k-1)\pi}{n+1}}^{\frac{k\pi}{n+1}} \Omega(t) \frac{dt}{t^2} \geq \frac{2M}{5\pi(n+1)} \int_{\frac{\pi}{n+1}}^{\frac{\pi}{2}} \Omega(t) \frac{dt}{t^2}.$$

Since  $\Omega$  is non-decreasing, we have

$$\int_{\frac{\pi}{n+1}}^{\frac{\pi}{2}} \Omega(t) \frac{dt}{t^2} = \frac{1}{\pi} \int_2^{n+1} \Omega\left(\frac{\pi}{t}\right) dt \geq \frac{1}{\pi} \sum_{k=2}^n \Omega\left(\frac{\pi}{k+1}\right).$$

Hence

$$(2.7) \quad A_n \geq \frac{2M}{5\pi^2(n+1)} \sum_{k=2}^n \Omega\left(\frac{\pi}{k+1}\right)$$

and the left hand side inequality (1.11) follows from (2.5), (2.6) and (2.7).

#### REFERENCES

- [1] FEJÉR, L., Untersuchungen über Fouriersche Reihen. *Math. Annalen*, 58 (1904), 501-569.
- [2] LEBESGUE, H., Recherches sur la convergence des séries de Fourier. *Math. Annalen*, 61 (1905), 251-280.
- [3] ZYGMUND, A., *Trigonometric Series*. Cambridge University Press, 1959.
- [4] JACKSON, D., *The Theory of Approximation*. American Math. Soc. Coll. Publ., Vol. XI, New York, 1930.
- [5] STEČKIN, S. B., On the approximation of periodic functions by Fejér sums (Russian). *Trudy Mat. Inst. V. A. Steklova*, LXII (1961), 48-60.

(Reçu le 2 novembre 1968.)

Proleterskih Brigada 62  
Belgrade, Yugoslavia

Department of Mathematics  
Ohio State University  
Columbus, Ohio 43210

Istarska 22  
Belgrade, Yugoslavia