

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 15 (1969)  
**Heft:** 1: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** SOME CONVERSE THEOREMS ON THE ABSCISSAE OF SUMMABILITY OF GENERAL DIRICHLET SERIES  
**Autor:** Rajagopal, C. T.  
**Kapitel:** §2. A BASIC THEOREM  
**DOI:** <https://doi.org/10.5169/seals-43225>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 09.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## § 2. A BASIC THEOREM

The theorem which follows supplies a basis for all the other theorems of this paper, whether by itself or not.

**THEOREM I. (A)** *For the Dirichlet series*

$$\sum_1^{\infty} \frac{a_n}{l_n^s}, \quad s = \sigma + i\tau,$$

*suppose that  $\sigma_r < \rho$  for some  $r > 0$ . Suppose also that there is a  $\gamma$  and an associated  $\theta(x)$  such that*

$$\sigma_r < \gamma < \rho, \quad \theta(x) \equiv x^{1-(\rho-\gamma)/r}, \quad (2.1)$$

*with*

EITHER (a)

$$\left. \begin{aligned} \overline{\lim}_{n \rightarrow \infty} \max_{l_n \leq l_m < \varepsilon \theta(l_n)} \frac{a_n + a_{n+1} + \dots + a_m}{l_n^\rho} &= o_R(1), \varepsilon \rightarrow 0, \\ \text{OR (b)} \quad \overline{\lim}_{n \rightarrow \infty} \max_{l_n \leq l_m < \varepsilon \theta(l_n)} \frac{|a_{n+1} + a_{n+2} + \dots + a_m|}{l_n^\rho} &= o(1), \varepsilon \rightarrow 0. \end{aligned} \right\} \quad (2.2)$$

*Then*

$$\sigma_k \leq \frac{(r-k)\rho + k\sigma_r}{r} \quad (0 \leq k < r). \quad (2.3)$$

(B) *If  $\sigma_r \geq \rho$ , instead of  $\sigma_r < \rho$  as in (A), and  $\rho$  is such that*

EITHER (a)

$$\left. \begin{aligned} \overline{\lim}_{n \rightarrow \infty} \max_{l_n \leq l_m < \varepsilon l_n} \frac{a_n + a_{n+1} + \dots + a_m}{l_n^\rho} &= o_R(1), \varepsilon \rightarrow 0, \\ \text{OR (b)} \quad \overline{\lim}_{n \rightarrow \infty} \max_{l_n \leq l_m < \varepsilon l_n} \frac{|a_{n+1} + a_{n+2} + \dots + a_m|}{l_n^\rho} &= o(1), \varepsilon \rightarrow 0, \end{aligned} \right\} \quad (2.4)$$

*then*

$$\sigma_k = \sigma_r \quad (0 \leq k < r). \quad (2.5)$$

*Proof.* (A) The proof is given below only for the case in which  $\{a_n\}$  satisfies the hypothesis in alternative (2.2) (a), the remaining case of alternative (2.2) (b) being exactly similar.

By the definition of  $\sigma_r$  and that of  $\gamma$  in (2.1),  $\sum a_n l_n^{-\gamma}$  is summable  $(R, l_n, r)$  to sum  $S$  (say), and so

$$b_1 + b_2 + \dots = (a_1 l_1^{-\gamma} - S) + a_2 l_2^{-\gamma} + \dots \text{ is summable } (R, l_n, r) \text{ to } 0,$$

i.e.,

$$B^r(x) = o(x^r), \quad x \rightarrow \infty, \quad (2.6)$$

while it is easy to prove that

$$\overline{\lim}_{n \rightarrow \infty} \max_{l_n \leq l_m < l_n + \varepsilon \theta(l_n)} \frac{b_n + b_{n+1} + \dots + b_m}{l_n^{\rho-\gamma}} = o_R(1), \quad \varepsilon \rightarrow 0, \quad (2.7)$$

distinguishing between the case  $\gamma \geq 0$  and  $\gamma < 0$ . If  $\gamma \geq 0$ , then

$$\begin{aligned} b_n + b_{n+1} + \dots + b_m &= a_n l_n^{-\gamma} + a_{n+1} l_{n+1}^{-\gamma} + \dots + a_m l_m^{-\gamma} \quad (n > 1) \\ &\leq l_n^{-\gamma} \max_{n \leq v \leq m} (a_n + a_{n+1} + \dots + a_v), \end{aligned}$$

from which and (2.2) (a) we have (2.7) as an immediate consequence. On the other hand, if  $\gamma < 0$ , then

$$b_n + b_{n+1} + \dots + b_m \leq l_m^{-\gamma} \max_{n \leq v \leq m} (a_v + a_{v+1} + \dots + a_m), \quad (2.8)$$

where

$$l_n \leq l_v \leq l_m < l_n + \varepsilon \theta(l_n) < (1 + \varepsilon) l_n,$$

$$\frac{\theta(l_n)}{\theta(l_v)} = \left(\frac{l_n}{l_v}\right)^{(r-\rho+\gamma)/r} < (1 + \varepsilon)^{|r-\rho+\gamma|/r} = K \text{ (say).}$$

Hence (2.8) gives us

$$\begin{aligned} &\max_{l_n \leq l_m < l_n + \varepsilon \theta(l_n)} \frac{b_n + b_{n+1} + \dots + b_m}{l_n^{\rho-\gamma}} \\ &\leq \frac{l_m^{-\gamma}}{l_n^{\rho-\gamma}} \max_{l_v \leq l_m < l_n + \varepsilon \theta(l_n)} (a_v + a_{v+1} + \dots + a_m) \\ &\leq \left(\frac{l_m}{l_v}\right)^{-\gamma} \left(\frac{l_v}{l_n}\right)^{\rho-\gamma} \max_{l_v \leq l_m < l_v + \varepsilon K \theta(l_v)} \frac{a_v + a_{v+1} + \dots + a_m}{l_v^\rho}. \end{aligned} \quad (2.9)$$

In (2.9),

$$\left(\frac{l_m}{l_v}\right)^{-\gamma} \leqslant 1, \left(\frac{l_v}{l_n}\right)^{\rho-\gamma} < (1+\varepsilon)^{\rho-\gamma}$$

since  $\gamma < 0$ ,  $\rho - \gamma > 0$ . Hence (2.9) in conjunction with hypothesis (2.2) (a) leads to (2.7) again.

After this we appeal to Theorem A with (1.1) and (1.2) (a) replaced by (2.6) and (2.7) respectively, to obtain

$$B(x) \equiv B(l_n) = o(l_n^{\rho-\gamma}) = o(x^{\rho-\gamma}), \quad l_{n+1} > x \geqslant l_n \rightarrow \infty.$$

From the last step and (2.6), we get, by using Lemma 3,

$$\begin{aligned} B^k(x) &= o\left\{x^{\left(1-\frac{k}{r}\right)(\rho-\gamma)+\frac{k}{r}}\right\} \text{ for } 0 \leqslant k < r \\ &= o(x^{k+\beta}) \text{ where } \beta = \left(1-\frac{k}{r}\right)(\rho-\gamma) > 0. \end{aligned}$$

Hence  $\sum a_n l_n^{-\gamma-\beta}$ , being summable  $(R, l_n, r)$ , is also summable  $(R, l_n, k)$  by Lemma 2, i.e.,  $\sum a_n l_n^{-s}$  is summable  $(R, l_n, k)$  for

$$\sigma \geqslant \gamma + \beta = \gamma + \left(1-\frac{k}{r}\right)(\rho-\gamma) = \left(1-\frac{k}{r}\right)\rho + \frac{k}{r}\gamma. \quad (2.10)$$

Since  $\gamma > \sigma_r$  may be taken as near to  $\sigma_r$  as we please<sup>1)</sup>, (2.10) immediately gives us the conclusion (2.3).

In arriving at (2.3) we have tacitly assumed that  $\sigma_r > -\infty$ . When  $\sigma_r = -\infty$ , we still reach (2.3) in the sense that  $\sigma_k = -\infty$  for  $0 \leqslant k < r$ , as we may see by taking  $\gamma = -G$  ( $G$  positive and arbitrarily large) in the preceding argument.

(B) As in (A), we confine ourselves to the hypothesis (2.4) (a), the treatment of (2.4) (b) being precisely similar. Defining as in (A)

$$b_1 + b_2 + \dots \equiv (a_1 l_1^{-\gamma} - S) + a_2 l_2^{-\gamma} + \dots \quad (\gamma > \sigma_r),$$

we see that (2.6) holds again, while (2.4) (a) implies

$$\lim_{n \rightarrow \infty} \max_{l_n \leq l_m < l_n + \varepsilon l_n} \frac{(b_n + b_{n+1} + \dots + b_m)}{l_n^{\rho-\gamma}} = o_R(1), \quad \varepsilon \rightarrow 0,$$

<sup>1)</sup> The truth of hypothesis (2.2) for some  $\gamma$ ,  $\sigma_r < \gamma < \rho$ , implies its truth for any  $\gamma'$ ,  $\sigma_r < \gamma' < \gamma$ , so that  $\gamma$  may be replaced by  $\gamma'$ .

exactly as (2.2) (a) implies (2.7). Since now  $\gamma > \sigma_r \geq \rho$ , the above condition in its turn implies

$$\overline{\lim}_{n \rightarrow \infty} \max_{l_n \leq l_m < l_n + \varepsilon l_n} (b_n + b_{n+1} + \dots + b_m) = o_R(1), \quad \varepsilon \rightarrow 0.$$

By Theorem A with hypothesis (1.2) (a) and  $a = b = 0$ , it follows that  $\sum a_n l_n^{-s}$  is convergent for any  $\sigma$  such that  $\sigma \geq \gamma > \sigma_r$  and therefore  $\sigma_0 \leq \sigma_r$ . But, in any case,  $\sigma_0 \geq \sigma_k \geq \sigma_r$  for  $0 \leq k < r$  and so we have the conclusion (2.5).

In the preceding argument we have supposed that  $\sigma_r < \infty$  since  $\sigma_r = \infty$  implies trivially  $\sigma_k = \infty$ .

### § 3. APPLICATIONS TO THEOREMS OF THE SCHNEE-LANDAU TYPE

Theorem II given next is the simplest of the theorems of the type mentioned above and it is a direct combination of Theorems I, B. Theorems V, VI are generalizations, respectively of Ananda-Rau's and Ganapathy Iyer's extensions of the Schnee-Landau theorem ([2], Theorem 9; [7], Theorem 10), as given by Chandrasekharan and Minakshisundaram ([6], pp. 88-9, Corollaries 3.73, 3.74). Theorems III, IV are apparently new counterparts of Theorems V, VI, the newness consisting in the replacement of the two-sided Tauberian conditions of the latter pair of theorems by analogous one-sided conditions suitably supplemented.

**THEOREM II.** *Suppose that (i) the Dirichlet series,*

$$\sum_1^\infty \frac{a_n}{l_n^s}, \quad s = \sigma + i\tau,$$

*is summable ( $R, l_n, q$ ) for some  $q \geq 0$  when  $\sigma > \rho$ , (ii) the sum-function  $f(s)$  thus defined is regular for  $\sigma > \eta$  when  $\eta < \rho$ , and satisfies the condition*

$$f(s) = O(|\tau|^r), \quad r > 0, \quad \text{uniformly for } \sigma \geq \eta + \varepsilon > \eta,$$

(iii) *the coefficients  $a_n$  of the Dirichlet series satisfy ONE of the two alternatives (a), (b) of (2.2), but with  $\theta(x) \equiv x^{1-(\rho-\eta)/r}$ . Then the Dirichlet series is summable ( $R, l_n, k$ ),  $0 \leq k < r$ , for*

$$\sigma \geq \frac{(r-k)\rho + k\eta}{r}.$$

*Proof.* By Theorem B, the Dirichlet series is summable ( $R, l_n, r'$ ),  $r' > r$ , for  $\sigma > \eta$  and hence  $\sigma_r \leq \eta < \rho$ . Therefore it is evident from the proof of