

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 16 (1970)  
**Heft:** 1: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** FUNCTIONAL EQUATIONS CONNECTED WITH ROTATIONS AND THEIR GEOMETRIC APPLICATIONS  
**Autor:** Schneider, Rolf  
**Kapitel:** 1. Introduction  
**DOI:** <https://doi.org/10.5169/seals-43867>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 14.03.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# FUNCTIONAL EQUATIONS CONNECTED WITH ROTATIONS AND THEIR GEOMETRIC APPLICATIONS

by Rolf SCHNEIDER

## 1. INTRODUCTION

In the geometry of convex bodies there is a certain type of questions which lead to some linear functional equations on the Euclidean sphere. Among these equations are, e.g., certain integral equations as well as relations between the values of a function at certain sets of finitely many arguments. What we mean by "linear functional equations on the sphere" is perhaps best described by giving some typical results:

Let  $E^d$  ( $d \geq 3$ ) denote  $d$ -dimensional Euclidean space with scalar product  $\langle \cdot, \cdot \rangle$ , and let

$$\Omega_d := \{x \in E^d \mid \langle x, x \rangle = 1\}$$

be its unit sphere.

**THEOREM 1.1.** *Let  $f$  be a real continuous function on  $\Omega_d$  satisfying*

$$(1.1) \quad f(u_1) + \dots + f(u_d) = 0$$

*for any  $d$  pairwise orthogonal vectors  $u_1, \dots, u_d \in \Omega_d$ . Then  $f$  is a spherical harmonic of degree 2.*

**THEOREM 1.2.** *Let  $f$  be a real continuous function on  $\Omega_d$  satisfying*

$$(1.2) \quad f(u_1) + \dots + f(u_{d+1}) = 0$$

*whenever  $u_1, \dots, u_{d+1}$  are the vertices of a regular simplex inscribed in  $\Omega_d$ . Then  $f$  is a sum of spherical harmonics of degrees 0, 1, 2, 5 if  $d = 3$ , and of degrees 0, 1, 2 if  $d \geq 4$ .*

For  $v \in \Omega_d$  let

$$s_v := \{u \in \Omega_d \mid \langle u, v \rangle = 0\}$$

be the great sphere with pole  $v$ ; and let  $\lambda_v$  denote the  $(d-2)$ -dimensional Lebesgue measure on  $s_v$ .

THEOREM 1.3. *If  $f$  is a real, even, continuous function on  $\Omega_d$  satisfying*

$$(1.3) \quad \int_{S_v} f d\lambda_v = 0$$

*for each  $v \in \Omega_d$ , then  $f = 0$ .*

In the following, by a measure on  $\Omega_d$  we understand a real valued, countably additive set function, defined on the  $\sigma$ -field of Borel subsets of  $\Omega_d$ . A measure  $\varphi$  on  $\Omega_d$  is called *even* (respectively *odd*) if  $\varphi(B) = \varphi(B^*)$  ( $\varphi(B) = -\varphi(B^*)$ ) for any two antipodal Borel subsets  $B, B^*$  of  $\Omega_d$ .

THEOREM 1.4. *If  $\varphi$  is an even measure on  $\Omega_d$  satisfying*

$$(1.4) \quad \int_{\Omega_d} |\langle u, v \rangle| d\varphi(u) = 0$$

*for each  $v \in \Omega_d$ , then  $\varphi = 0$ .*

For  $v \in \Omega_d$  let

$$S_v := \{u \in \Omega_d \mid \langle u, v \rangle > 0\}$$

be the open hemisphere with center  $v$ .

THEOREM 1.5. *If  $\varphi$  is an odd measure on  $\Omega_d$  satisfying*

$$(1.5) \quad \varphi(S_v) = 0$$

*for each  $v \in \Omega_d$ , then  $\varphi = 0$ .*

To functional relations of the types (1.1)–(1.5) one is lead by some uniqueness and characterization problems in the theory of convex bodies; and the theorems quoted above (all of which are essentially known—see section 3) have interesting geometric interpretations. The main purpose of this note is to treat the above functional equations from a unifying point of view and to exhibit them as special cases of one general equation.

In fact, each of these equations can be written in the form (see section 3)

$$(1.6) \quad \int_{\Omega_d} f(\delta u) d\varphi(u) = 0 \quad \text{for each } \delta \in SO(d).$$

Here  $SO(d)$  denotes the  $d$ -dimensional rotation group acting on  $\Omega_d$ ,  $f$  is a function and  $\varphi$  a measure on  $\Omega_d$ . Equation (1.6) may be read in two ways: Either  $\varphi$  is given, then (1.6) is a functional equation for  $f$ , or  $f$  is given and  $\varphi$  is to be determined.

We shall now state a theorem which gives necessary and sufficient conditions for a pair  $f, \varphi$  in order that (1.6) be true. From these conditions the uniqueness theorems 1.1–1.5, and some others, can be immediately deduced via some elementary computations.