

<b>Zeitschrift:</b>	L'Enseignement Mathématique
<b>Herausgeber:</b>	Commission Internationale de l'Enseignement Mathématique
<b>Band:</b>	16 (1970)
<b>Heft:</b>	1: L'ENSEIGNEMENT MATHÉMATIQUE
<b>Artikel:</b>	FUNCTIONAL EQUATIONS CONNECTED WITH ROTATIONS AND THEIR GEOMETRIC APPLICATIONS
<b>Autor:</b>	Schneider, Rolf
<b>Kapitel:</b>	2. The main theorem
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-43867">https://doi.org/10.5169/seals-43867</a>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 02.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## 2. THE MAIN THEOREM

Let us first give some definitions. In the following, all functions are complex valued and continuous. Let  $\omega$  denote Lebesgue measure on  $\Omega_d$ , normalized so that  $\omega(\Omega_d) = 1$ .

For functions  $f, g$  and measures  $\varphi$  on  $\Omega_d$  we write  $(f, g) := \int f \bar{g} d\omega$  and  $(f, \varphi) := \int f d\varphi$ , where the integrals are extended over  $\Omega_d$ . For  $n = 0, 1, 2, \dots$  let  $\mathfrak{H}_n$  denote the complex vector space of spherical harmonics of degree  $n$  on  $\Omega_d$ ; let  $N_{d,n}$  be its dimension. If  $f$  is a function on  $\Omega_d$ , we say that  $\mathfrak{H}_n$  occurs in  $f$  if and only if the orthogonal projection of  $f$  onto  $\mathfrak{H}_n$  does not vanish, i.e. if  $(f, Y_n) \neq 0$  for some spherical harmonic  $Y_n$  of degree  $n$  (or, equivalently, if  $\int f(u) C_n^v(\langle u, v \rangle) d\omega(u)$  does not vanish identically, where  $C_n^v$  is the Gegenbauer polynomial of degree  $n$  and order  $v = \frac{1}{2}(d-2)$ ). Analogously, we say that  $\mathfrak{H}_n$  occurs in the measure  $\varphi$  if and only if  $(Y_n, \varphi) \neq 0$  for some  $Y_n \in \mathfrak{H}_n$ . If  $f$  is a function on  $\Omega_d$  and  $\delta \in SO(d)$  is a rotation, the left translate  $\delta f$  of  $f$  by  $\delta$  is defined by  $(\delta f)(u) = f(\delta^{-1}u)$  for  $u \in \Omega_d$ .

**THEOREM 2.1.** *Let  $f$  be a continuous function and  $\varphi$  a measure on  $\Omega_d$ . In order that  $(\delta f, \varphi) = 0$  for each  $\delta \in SO(d)$ , it is necessary and sufficient that none of the spaces  $\mathfrak{H}_n$ ,  $n \in \{0, 1, 2, \dots\}$ , occurs in both,  $f$  and  $\varphi$ .*

We remark that this theorem, together with its proof to be given below, carries over to the following more general situation:  $SO(d)$  and  $\Omega_d$  may be replaced, respectively, by a compact connected topological group  $G$  and by the homogeneous manifold  $G/K$ , where  $K (= SO(d-1)$  in our case) is a closed subgroup of  $G$ . The rôle of the spherical harmonics is then played by their natural generalizations. We do not write down this generalization explicitly since we do not know any application of it.

*Proof of Theorem 2.1.* Let  $\{Y_{ni}; i=1, \dots, N_{d,n}\}$  be an orthonormal basis of  $\mathfrak{H}_n$  ( $n = 0, 1, 2, \dots$ ). Let us first suppose that  $f$  is a finite sum of spherical harmonics,

$$(2.1) \quad f = \sum_{n=0}^k \sum_{j=1}^{N_{n,d}} (f, Y_{nj}) Y_{nj}.$$

Since  $\mathfrak{H}_n$  is invariant under the action of  $SO(d)$  by left translation, we have a relation

$$(2.2) \quad Y_{nj}(\delta^{-1} u) = \sum_{i=1}^{N_{d,n}} t_{ij}^n(\delta) Y_{ni}(u)$$

for each  $\delta \in SO(d)$ , by which continuous functions  $t_{ij}^n$  on  $SO(d)$  are defined. It is well known that, for each  $n \in \{0, 1, 2, \dots\}$ , the mapping  $\delta \rightarrow (t_{ij}^n(\delta))$  is a unitary, irreducible matrix valued representation of the group  $SO(d)$ . From (2.1) and (2.2) we get

$$(2.3) \quad (\delta f, \varphi) = \sum_{n=0}^k \sum_{i,j=0}^{N_{d,n}} t_{ij}^n(\delta) (f, Y_{nj})(Y_{ni}, \varphi).$$

If  $f$  is an arbitrary continuous function on  $\Omega_d$ , then  $f$  can be uniformly approximated by a sequence  $f_1, f_2, \dots$ , where each  $f_k$  is a finite sum of spherical harmonics of those degrees  $n$  only, for which  $\mathfrak{H}_n$  occurs in  $f$  (see, e.g., Weyl [22], p. 499).

Let us now suppose that  $\mathfrak{H}_n$  does not occur in both,  $f$  and  $\varphi$  ( $n = 0, 1, 2, \dots$ ). Approximate  $f$  as explained above. Then if  $(Y_{ni}, \varphi) \neq 0$  for some  $n$  and some  $i \in \{1, \dots, N_{d,n}\}$ , the space  $\mathfrak{H}_n$  does not occur in  $f$ . Therefore we have  $(f_k, Y_{nj}) = 0$  ( $k = 1, 2, \dots$ ) for each  $j \in \{1, \dots, N_{d,n}\}$ , since  $f_k$  is a finite sum of spherical harmonics of degrees other than  $n$ . This shows that  $(f_k, Y_{nj})(Y_{ni}, \varphi) = 0$  for each possible choice of  $k, n, i, j$ , and hence  $(\delta f_k, \varphi) = 0$  by (2.3). For  $k \rightarrow \infty$  we get  $(\delta f, \varphi) = 0$ , which proves one half of the theorem.

In order to prove the other direction of Theorem 2.1, we multiply equation (2.3) by  $\overline{t_{km}^n(\delta)}$  and integrate over  $SO(d)$  with respect to the normalized Haar measure  $\mu$ . Using one of the well known orthogonality relations for the matrix elements of unitary, irreducible representations of a compact group, namely

$$N_{d,n} \int_{SO(d)} t_{ij}^n(\delta) \overline{t_{km}^n(\delta)} d\mu(\delta) = \delta_{ik} \delta_{jm},$$

we arrive at

$$N_{d,n} \int_{SO(d)} (\delta f, \varphi) \overline{t_{km}^n(\delta)} d\mu(\delta) = (f, Y_{nm})(Y_{nk}, \varphi),$$

provided  $f$  is a finite sum of spherical harmonics. By approximation, this holds for arbitrary continuous  $f$ . If now (1.6) is assumed, we get  $(f, Y_{nm})(Y_{nk}, \varphi) = 0$  for  $n = 0, 1, 2, \dots$  and  $k, m \in \{1, \dots, N_{d,n}\}$ , which shows that  $\mathfrak{H}_n$  does not occur in both,  $f$  and  $\varphi$ . Theorem 2.1 is proved.