Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 17 (1971)

Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: APPROXIMATION TO ALGEBRAIC NUMBERS

Autor: Schmidt, Wolfgang M.

Kapitel: 11. Generalizations and open problems

DOI: https://doi.org/10.5169/seals-44578

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 24.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

discuss what happens in the case (B). For simplicity we shall assume that K is totally real.

There is an element φ of the Galois group such that the sets $\{\alpha^{(1)}, ..., \alpha^{(\mu)}\}$ and $\{\varphi(\alpha^{(1)}), ..., \varphi(\alpha^{(\mu)})\}$ are neither identical nor disjoint. We may assume without loss of generality that

$$\left\{ \left. \varphi \left(\alpha ^{(1)} \right),...,\varphi \left(\alpha ^{(\mu)} \right) \right\} \right. \\ = \left. \left\{ \left. \alpha ^{(1)},...,\alpha ^{(l)},\alpha ^{(\mu +1)},...,\alpha ^{(2\,\mu -l)} \right\} \right. \\ .$$

Here $1 \le l \le \mu - 1$. The forms $L^{(1)}$, ..., $L^{(\mu)}$ have rank ρ , and hence also $L^{(1)}$, ..., $L^{(l)}$, $L^{(\mu+1)}$, ..., $L^{(2\mu-l)}$ have rank ρ . Denote the rank of $L^{(1)}$, ..., $L^{(l)}$ by r_1 and the rank of $L^{(1)}$, ..., $L^{(\mu)}$, ..., $L^{(2\mu-l)}$ by r_2 . It is easily seen that $r_2 \le 2\rho - r_1$, i.e. that

$$r_1 + r_2 \leq 2\rho$$
.

Since μ was chosen as small as possible with (10.10), and since $l \leq \mu - 1$, we have $l < r_1 q$. The number $2\mu - l$ of elements of $L^{(1)}, ..., L^{(\mu)}, ..., L^{(2\mu-l)}$ satisfies $2\mu - l \leq r_2 q$ by (10.9). Thus

$$2\mu = l + (2\mu - l) < r_1 q + r_2 q \le 2\rho q,$$

which contradicts (10.10). Hence (B) is impossible if K is totally real.

We have in fact used the hypothesis that K is totally real, for in general $L^{(1)}$, ..., $L^{(l)}$ need not be a Symmetric System, and $l < r_1 q$ need not hold. The situation is therefore somewhat more complicated if K is not totally real.

11. Generalizations and open problems

11.1. The theorems of §7 and §10 can almost certainly be generalized to include p-adic valuations. I understand that work on this question is being done now. (p-adic versions of the results of §2 were discussed in §4.5). Next, suppose that K is an algebraic number field and that $\alpha_1, ..., \alpha_l$ are algebraic numbers such that $1, \alpha_1, ..., \alpha_l$ are linearly independent over K. It is likely that for every $\delta > 0$ there are only finitely many l-tuples of elements $\beta_1, ..., \beta_l$ of K with

$$(11.1) |\alpha_i - \beta_i| < \mathcal{H}(\beta)^{-1 - (1/l) - \delta} (i = 1, ..., l),$$

where $\mathcal{H}(\beta)$ is a suitably defined height of $\beta = (\beta_1, ..., \beta_l)$. A possible definition for $\mathcal{H}(\beta)$ is

$$\mathcal{H}(\beta) = \prod_{v} \max (1, ||\beta_1||_v, ..., ||\beta_l||_v),$$

where v runs through the valuations of K and where $|| ||_v$ is defined as in §4.5. In view of (4.9), $\mathcal{H}(\beta)$ is almost the same as $H_K(\beta)$ if l=1, and hence a theorem on (11.1) would generalize Le Veque's Theorem 4A. One could try to obtain a still more general theorem which would contain both the p-adic case and the case of a number field K. Such a result was put forward as a conjecture by Lang (1962, Ch. 6).

11.2. We have already said in §2.2 that it would be desirable to replace the factor q^{δ} in Roth's Theorem by something smaller, say by a power of log q. The same is true of the generalizations of Roth's Theorem to simultaneous approximation.

The theorems of §2, 7 and 10 are non-effective. For approximation to a single algebraic number α there are the effective results of Baker (see §5), but for simultaneous approximation there are only the relatively special effective theorems of Baker (1967a), Feldman (1970a, 1970b) and Osgood (1970).

11.3. The following questions also appear to be very difficult. Suppose $(\alpha_1, ..., \alpha_l)$ is a point of transcendence degree d < l. The theorems of §7 deal with the case when the point is algebraic, i.e. when d = 0. What can one say for other values of d? Perron (1932) and Schmidt (1962) obtained results, of about the same level of sophistication as Liouville's Theorem, which can be used to show that certain given points have transcendence degree l.

A better question perhaps is how close rational points can come to a given algebraic variety. We may reformulate this question in a homogeneous setting. Let V be a homogeneous variety defined over the rationals (i.e. one defined by homogeneous polynomial equations with rational coefficients) in E^n with $n \ge 2$. For every $\mathbf{x} \ne \mathbf{0}$ we put

$$\psi(V, \mathbf{x}) = \Delta(V, \mathbf{x}) |\mathbf{x}|^{-1}$$

where $\Delta(V, \mathbf{x})$ is the distance from \mathbf{x} to V. It is clear that $\psi(V, \lambda \mathbf{x}) = \psi(V, \mathbf{x})$; the function $\psi(V, \mathbf{x})$ may be interpreted as the "angle" between V and the vector \mathbf{x} . We are interested in inequalities of the type

$$(11.2) \psi(V, \mathbf{x}) < c |\mathbf{x}|^{-\omega}$$

where x runs through the integer points. We saw in §6 that Theorems 6A and 6C had such an interpretation. The best value of ω for which (11.2) has

infinitely many integer solutions can always be found if V is linear, i.e. is a subspace. For the non-linear case we have neither a good generalization of Dirichlet's Theorem nor anything like Roth's Theorem.


Suppose now that V is a hypersurface containing no integer point $\mathbf{x} \neq \mathbf{0}$ and defined by the equation $F(\mathbf{x}) = 0$ where F is a form of degree d with rational integer coefficients. For every integer point $\mathbf{x} \neq \mathbf{0}$ we have

 $|F(\mathbf{x})| \ge 1$, and since $|\frac{\partial}{\partial x_i} F(\mathbf{x})| \le c_1 |\mathbf{x}|^{d-1}$ (i=1,...,n), the distance from \mathbf{x} to V is $\ge c_2 |\mathbf{x}|^{1-d}$, which in turn implies that

$$\psi(V, \mathbf{x}) \geq c_3 |\mathbf{x}|^{-d}$$
,

where the constants depend only on V. This inequality may be interpreted as a generalization of Liouville's Theorem. Any improvement of this inequality, even though perhaps it may apply only to special classes of non-linear hypersurfaces, would be of great interest and would shed light on certain diophantine equations different from the equations with norm forms discussed in §10.

REFERENCES



— (1967a). Simultaneous approximations to certain algebraic numbers. Proc. Camb.

– (1967b). Linear forms in the logarithms of algebraic numbers (II). Mathematika 14,

204-216.

102-107.

Phil. Soc. 63, 693-702.