

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 19 (1973)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ASYMPTOTIC PROPERTIES OF LINEAR OPERATORS  
**Autor:** Bojanic, R. / Vuilleumier, M.  
**Kapitel:** 2. Boundedness and convergence preserving regular operators.  
**DOI:** <https://doi.org/10.5169/seals-46293>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

$V_\Psi$  is additive on non-negative functions in  $\mathcal{F}_1$  and homogeneous with respect to multiplication by positive real numbers:

$$\begin{aligned} f_1 \geq 0 \text{ and } f_2 \geq 0 &\Rightarrow V_\Psi(f_1 + f_2, x) = V_\Psi(f_1, x) + V_\Psi(f_2, x), \\ \alpha > 0 \text{ and } f \geq 0 &\Rightarrow V_\Psi(\alpha f, x) = \alpha V_\Psi(f, x). \end{aligned}$$

The operator  $V_\Psi$  is then extended to a positive linear operator  $\Phi_\Psi: \mathcal{F}_1 \rightarrow \mathcal{F}$ , which coincides with  $V_\Psi$  on non-negative functions, in the usual way: if  $f = f^+ - f^-$ , then  $\Phi_\Psi(f, x) = V_\Psi(f^+, x) - V_\Psi(f^-, x)$ . Finally, if  $f \in \mathcal{F}_1$  we have  $|\Psi(f, x)| \leq V_\Psi(|f|, x) = \Phi_\Psi(|f|, x)$ . Hence, by Theorem I, the operator  $\Psi$  is regular.

Conversely, if  $\Psi$  is regular, by Theorem I, we have a positive linear operator  $\Phi: \mathcal{F}_1 \rightarrow \mathcal{F}$  such that

$$|\Psi(g, x)| \leq \Phi(|g|, x) \leq \Phi(f, x)$$

for every  $g \in \mathcal{F}_1$ ,  $|g| \leq f$ , and every  $x \in R^+$ . Hence, the condition (1.4) of Theorem II is satisfied.

If we consider the operator  $G$  defined by (1.1), then

$$V_G(f, x) = \int_0^\infty |\psi(x, t)| f(t) dt.$$

From the statement of the theorems of Hahn and Raff mentioned earlier, we can expect that the operator  $V_\Psi$  will play an important role in the extension of these results to general regular operators. In fact, as in the theory of positive linear operators, some asymptotic property of the regular operator  $\Psi$  will hold for a large class of functions if and only if the operator  $V_\Psi$  has certain properties on a much smaller class of functions.

## 2. BOUNDEDNESS AND CONVERGENCE PRESERVING REGULAR OPERATORS.

In this section and the following one we shall extend to regular operators some of the well-known results about the asymptotic behavior of the special transform  $G$  defined by (1.1).

Let us consider the linear space  $\mathcal{M}$  of real valued measurable functions on  $R^+$  and let  $\mathcal{M}_0$  be the subspace of  $\mathcal{M}$  consisting of all measurable functions on  $R^+$  which are bounded on every finite interval of  $R^+$ .

The basic result which characterizes regular operators from  $\mathcal{M}_0$  into  $\mathcal{F}_0$  that preserve boundedness can be stated as follows:

THEOREM 1. Let  $\Psi: \mathcal{M}_0 \rightarrow \mathcal{F}_0$  be a regular operator. In order that, as  $x \rightarrow \infty$ ,

$$(2.1) \quad f \in \mathcal{M}_0 \text{ and } f(x) = O(1) \Rightarrow \Psi(f, x) = O(1)$$

it is necessary and sufficient that

$$(2.2) \quad V_\Psi(1, x) = O(1)$$

where  $V_\Psi$  is defined by (1.5).

This result is clearly a natural extension of the results for the special operator  $G$  mentioned in section 1.2 under  $A$ . The corresponding result for regular operators which transform functions in  $\mathcal{S}_0$ , converging to zero as  $x \rightarrow \infty$  into bounded functions is given by the following theorem.

THEOREM 2. Let  $\Psi: \mathcal{M}_0 \rightarrow \mathcal{F}_0$  be a regular operator. In order that, as  $x \rightarrow \infty$ ,

$$(2.3) \quad f \in \mathcal{M}_0 \text{ and } f(x) \rightarrow 0 \Rightarrow \Psi(f, x) = O(1)$$

it is necessary and sufficient that

$$(2.4) \quad W_\Psi(1, x) = O(1)$$

where  $W_\Psi$  is defined for every  $f \in \mathcal{M}_0, f \geq 0$ , by

$$(2.5) \quad W_\Psi(f, x) = \sup \{ |\Psi(g, x)| : g \in \mathcal{M}_0, |g| \leq f, g = o(f) \}^1.$$

Condition (2.4) in Theorem 2 is less restrictive than the corresponding condition (2.2) in Theorem 1, since  $W_\Psi(f, x) \leq V_\Psi(f, x)$  for every  $f \in \mathcal{M}_0, f \geq 0$  and for every  $x \in R^+$ .

It is now easy to obtain a generalization of the results for the special operator  $G$  mentioned in section 1.2 under  $B$ , i.e. to establish necessary and sufficient conditions for a regular operator to be convergence preserving:

THEOREM 3. Let  $\Psi: \mathcal{M}_0 \rightarrow \mathcal{F}_0$  be a regular operator. In order that, as  $x \rightarrow \infty$ ,

$$(2.6) \quad f \in \mathcal{M}_0 \text{ and } f(x) \rightarrow c \Rightarrow \Psi(f, x) \rightarrow c$$

it is necessary and sufficient that

$$(2.7) \quad \Psi(1, x) \rightarrow 1,$$

---

<sup>1)</sup>  $g = o(f)$  means  $g(t) = o(f(t)) (t \rightarrow \infty)$ .

$$(2.8) \quad \Psi(\chi_E, x) \rightarrow 0$$

for every bounded measurable subset  $E$  of  $R^+$ , and

$$(2.9) \quad W_\Psi(1, x) = O(1).$$

### 3. TRANSFORMATIONS OF $O$ -REGULAR AND SLOWLY VARYING FUNCTIONS BY REGULAR OPERATORS.

3.1. The class of positive functions which are eventually bounded away from zero and infinity has been extended to the class of  $O$ -regular functions defined as follows:

A positive, measurable function  $l$  on  $R^+$  is  $O$ -regular if

$$(3.1) \quad \frac{l(\lambda x)}{l(x)} = O(1) \quad (x \rightarrow \infty)$$

for every  $\lambda > 0$ .

For example, any function  $l$  such that  $ax^\alpha \leq l(x) \leq Ax^\alpha$ , where  $\alpha \in R$ , clearly satisfies condition (3.1).

The class of  $O$ -regular functions and related classes of functions have been studied extensively by V. G. Avakumović [8, 9, 10, 11], J. Karamata [14], N. K. Bari, S. B. Stečkin [15], M. A. Krasnoselskiĭ, T. B. Rutickiĭ [16], W. Matuszewska [17] and others.

The closely related class of slowly varying ( $SV$ ) functions, introduced by J. Karamata ([12], [13]), generalizes the class of functions converging to a positive limit. A positive, measurable function  $L$  defined on  $R^+$  is a slowly varying function if

$$(3.2) \quad \lim_{x \rightarrow \infty} \frac{L(\lambda x)}{L(x)} = 1$$

for every  $\lambda > 0$ .

Clearly, every measurable function on  $R^+$  which converges to a positive limit as  $x \rightarrow \infty$  is a  $SV$  function. Also, functions like

$$\varphi(x) = \begin{cases} 1, & 0 \leq x < e, \\ \log x, & x \geq e, \end{cases}, \quad h(x) = \left(2 + \frac{\sin x}{x}\right) \varphi(x),$$

and their iterations are  $SV$  functions. More generally, any measurable function  $g$  on  $R^+$  such that  $\varphi(x) \leq g(x) \leq \varphi(x) + \sqrt{\varphi(x)}$  is a  $SV$  function.

The most important properties of  $O$ -regular and  $SV$  functions can be stated as follows: