

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 19 (1973)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AN INTEGRAL INEQUALITY IN ANALYTIC FUNCTION THEORY
Autor: Haruki, Hiroshi
DOI: <https://doi.org/10.5169/seals-46294>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

AN INTEGRAL INEQUALITY IN ANALYTIC FUNCTION THEORY

by Hiroshi HARUKI

The following theorem was proved in [1]:

THEOREM A. *Suppose that $f = f(z)$ is an entire function of a complex variable z . Then the only solutions of the functional inequality*

$$(1) \quad |f((x+y)/2)| \leq (|f(x)| + |f(y)|)/2,$$

where x, y are complex variables, are $f(z) = (Az + B)^n$ and $f(z) = \exp(Az + B)$ where A, B are arbitrary complex constants and n is an arbitrary positive integer.

Now, we shall prove that (1) implies the following integral inequality:

$$(2) \quad |1/(y-x) \int_C f(z) dz| \leq (|f(x)| + |f(y)|)/2,$$

where $f = f(z)$ is an entire function of z , x, y are complex variables ($x \neq y$) and C is an arbitrary contour joining two points x and y .

To this end we shall apply the following lemma, the easy proof of which is omitted:

LEMMA 1. If $k = k(t)$ is a real-valued continuous function of a real variable t and if k is convex on $[a, b]$, then

$$1/(b-a) \int_a^b k(t) dt \leq (k(a) + k(b))/2.$$

Now, we put $g(t) = |f(x+(y-x)t)|$, where $t \in [0, 1]$ and x, y are arbitrary distinct complex constants. $g(t)$ is a real-valued continuous function on $[0, 1]$. By (1) $g(t)$ is convex on $[0, 1]$. Hence, by Lemma 1 we have

$$\int_0^1 g(t) dt \leq (g(0) + g(1))/2.$$

Therefore we have

$$\int_0^1 |f(x + (y-x)t)| dt \leq (|f(x)| + |f(y)|)/2$$

and

$$(3) \quad |1/(y-x) \int_L f(z) dz| \leq (|f(x)| + |f(y)|)/2,$$

where L is the line segment joining the two points x and y . By Cauchy's Integral Theorem (3) implies (2).

The purpose of this note is to solve (2), i.e., to prove the following

THEOREM. *Suppose that $f = f(z)$ is an entire function of z . Then the only solutions of (2) are $f(z) = (Az+B)^n$ and $f(z) = \exp(Az+B)$ where A, B are arbitrary complex constants and n is an arbitrary positive integer.*

To this end we shall apply the following two lemmas:

LEMMA 2. (See [1].) Suppose that $f = f(z), g = g(z)$ are entire functions of z . If $|f(z)| \leq |g(z)|$ holds in $|z| < +\infty$, then $f(z) = Cg(z)$ where C is a complex constant with $|C| \leq 1$.

Proof. The proof is clear from Riemann's Theorem concerning a removable singularity and Liouville's Theorem.

LEMMA 3. Suppose that $H = H(z)$ is an entire function of z . If $A(t) = |H(t \exp(i\varphi))|^2$ where t, φ are real and φ is arbitrarily fixed, then we have

$$(i) \quad A''(0) = 2\operatorname{Re}(\exp(2i\varphi) H''(0) \overline{H(0)}) + 2|H'(0)|^2,$$

$$(ii) \quad A^{(4)}(0) = 2\operatorname{Re}(\exp(4i\varphi) H^{(4)}(0) \overline{H(0)} \\ + 4\exp(2i\varphi) H^{(3)}(0) \overline{H'(0)}) + 6|H''(0)|^2.$$

Proof. Since the proof is easy, we omit it.

We may now prove our theorem.

Let $F = F(z)$ be an entire function such that $F'(z) = f(z)$. By (2) we have for all complex x, y

$$2|F(y) - F(x)| \leq |y-x|(|f(x)| + |f(y)|).$$

By a corollary of Schwarz's Inequality ($(a+b)^2 \leq 2(a^2+b^2)$, a, b real) we have

$$2|F(x) - F(y)|^2 \leq |x-y|^2(|f(x)|^2 + |f(y)|^2).$$

Replacing x, y by $x+y, x-y$, respectively, we get

$$|F(x+y) - F(x-y)|^2 \leq 2|y|^2 (|f(x+y)|^2 + |f(x-y)|^2).$$

Putting $y = t \exp(i\varphi)$ where t, φ are real we have

$$\begin{aligned} & |F(x+t \exp(i\varphi)) - F(x-t \exp(i\varphi))|^2 \\ & \leq 2t^2 (|f(x+t \exp(i\varphi))|^2 + |f(x-t \exp(i\varphi))|^2). \end{aligned}$$

Keeping x, φ arbitrarily fixed and putting

$$\begin{aligned} p(t) = & 2t^2 (|f(x+t \exp(i\varphi))|^2 + |f(x-t \exp(i\varphi))|^2) \\ & - |F(x+t \exp(i\varphi)) - F(x-t \exp(i\varphi))|^2, \end{aligned}$$

$p(t)$ is a real-valued function of t and is of course four times differentiable on $|t| < +\infty$. Further $p(t)$ is an even function of t . Hence we have

$$(4) \quad p'(0) = 0, \quad p^{(3)}(0) = 0.$$

By Lemma 3 we have

$$\begin{aligned} (5) \quad p''(0) &= 8(|f(x)|^2 - |F'(x)|^2) = 0, \\ p^{(4)}(0) &= 96 \operatorname{Re}(\exp(2i\varphi) f''(x) \overline{f(x)}) + 96 |f'(x)|^2 \\ &\quad - 32 \operatorname{Re}(\exp(2i\varphi) F^{(3)}(x) \overline{F'(x)}) \\ &= 64 \operatorname{Re}(\exp(2i\varphi) f''(x) \overline{f(x)}) + 96 |f'(x)|^2. \end{aligned}$$

$p(t)$ has a minimum at $t = 0$ ($p(t) \geq 0$ on $|t| < +\infty, p(0) = 0$). Hence, by (4), (5) we have $p^{(4)}(0) \geq 0$, or

$$2 \operatorname{Re}(\exp(2i\varphi) f''(x) \overline{f(x)}) + 3 |f'(x)|^2 \geq 0.$$

x, φ were arbitrarily fixed. An appropriate choice of φ_0 gives

$$\operatorname{Re}(\exp(2i\varphi_0) f''(x) \overline{f(x)}) = -|f''(x) \overline{f(x)}| = -|f''(x) f(x)|.$$

Hence we have in $|x| < +\infty$

$$2|f''(x) f(x)| \leq 3|f'(x)|^2,$$

and by Lemma 2

$$(6) \quad f''(x) f(x) = K f'(x)^2,$$

where K is a complex constant with $|K| \leq 3/2$.

Solving (6) and taking into account the fact that f is an entire function, we have

$$(7) \quad f(z) = (Az + B)^n \quad \text{or} \quad f(z) = \exp(Az + B),$$

where A, B are complex constants and n is an arbitrary positive integer. By Theorem A, (7) satisfies (1). Since (1) implies (2) as already proved, (7) satisfies (2). Q.E.D.

REFERENCE

- [1] HIROSHI HARUKI. On the functional inequality $|f((x+y)/2)| \leq (|f(x)| + |f(y)|)/2$, *Journal of the Mathematical Society of Japan*, 16 (1964), pp. 39-41.

(Reçu le 5 mars 1973)

Hiroshi Haruki
Faculty of Mathematics
University of Waterloo
Waterloo, Ontario
Canada