

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 20 (1974)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SOME CLASSICAL THEOREMS ON DIVISION RINGS
Autor: Taylor, D. E.
Kapitel: 1. SOME LINEAR ALGEBRA
DOI: <https://doi.org/10.5169/seals-46912>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

SOME CLASSICAL THEOREMS ON DIVISION RINGS

by D. E. TAYLOR

The theorem of Wedderburn [15] that every finite division ring is a field, and the theorem of Frobenius [6] characterizing the quaternions as a non-commutative real division algebra can both be obtained as immediate and easy consequences of theorems on central simple algebras—particularly the Skolem-Noether theorem (van der Waerden [14, p. 199]). The purpose of this note is to use elementary linear algebra to prove a version of the Skolem-Noether theorem sufficient to yield the results of Wedderburn and Frobenius.

1. SOME LINEAR ALGEBRA

All the results of this section are quite elementary and can be found in most texts on linear algebra (for example: Hoffman and Kunze [9]).

Let V be a vector space over a field F and let T be a linear transformation of V . Suppose that $f(X)$ is a polynomial with coefficients in F such that $f(T) = 0$. If $f(X) = f_1(X)f_2(X)$ where $f_1(X)$ and $f_2(X)$ are coprime, then there are polynomials $g_1(X)$ and $g_2(X)$ such that $1 = f_1(X)g_1(X) + f_2(X)g_2(X)$. Then for each v in V the vector $v_1 = f_2(T)g_2(T)v$ belongs to the kernel, V_1 , of $f_1(T)$, the vector $v_2 = f_1(T)g_1(T)v$ belongs to the kernel, V_2 , of $f_2(T)$ and $v = v_1 + v_2$. Thus V is the (direct) sum of V_1 and V_2 . Moreover, the restriction T_i of T to V_i satisfied the equation $f_i(T_i) = 0$ for $i = 1, 2$.

It follows by induction on the degree that if $f(X)$ can be factorized over F into distinct linear factors, then V is the direct sum of the eigenspaces of T . Note that V is not assumed to be finite dimensional.

Recall that the minimal polynomial of T is the monic polynomial $m(X)$ of least degree such that $m(T) = 0$. It is immediate that each eigenvalue λ of T satisfies the equation $m(\lambda) = 0$ and conversely, the above considerations show that each root of $m(X)$ is an eigenvalue of T .