

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 20 (1974)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** SOME CLASSICAL THEOREMS ON DIVISION RINGS  
**Autor:** Taylor, D. E.  
**Kapitel:** 4. Frobenius' theorem  
**DOI:** <https://doi.org/10.5169/seals-46912>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 03.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

number of conjugates of a subgroup is the index of its normalizer, so  $H$  has at most  $|G : H|$  conjugates in  $G$  and hence the union of the conjugates contains at most  $|G : H| (|H| - 1) + 1 = |G| - |G : H| + 1$  elements. This number is less than  $|G|$  except when  $G = H$ . Hence  $D = F$  is a field.

#### 4. FROBENIUS' THEOREM

Let  $\mathbf{R}$  denote the field of real numbers,  $\mathbf{C}$  the field of complex numbers and  $\mathbf{H}$  the division ring of quaternions. The following proof makes use of the fundamental theorem that every polynomial with coefficients in  $\mathbf{C}$  has a root in  $\mathbf{C}$ .

**THEOREM.** *Let  $D$  be a division ring which contains the real numbers  $\mathbf{R}$  in its centre and suppose that every element of  $D$  satisfies a polynomial with coefficients in  $\mathbf{R}$ . Then  $D$  is isomorphic to one of  $\mathbf{R}$ ,  $\mathbf{C}$  or  $\mathbf{H}$ .*

*Proof.* Suppose that  $D$  is not isomorphic to  $\mathbf{R}$  or  $\mathbf{C}$ . It follows that the maximal subfield  $F$  of  $D$  is isomorphic to  $\mathbf{C}$ , the centre  $K$  of  $D$  is isomorphic to  $\mathbf{R}$  and  $F = K(i)$  where  $i^2 = -1$ . Let  $j$  be an eigenvector of  $T_i$  corresponding to the eigenvalue  $-i$ . Then  $ji = -ij$  and  $j^2$  commutes with  $j$  and  $F$ . From (2.2) and (2.3) the elements 1 and  $j$  form an  $F$ -basis for  $D$  and therefore  $j^2 = \alpha$  belongs to  $K$ . If  $\alpha = \beta^2$  for some  $\beta \in K$  then  $(j - \beta)(j + \beta) = 0$  and  $j$  belongs to  $K$ , which is not the case; hence  $\alpha = -\beta^2$  for some  $\beta \in K$ . Replacing  $j$  by  $j\beta^{-1}$  we obtain a  $K$ -basis 1,  $i$ ,  $j$ ,  $ij$  for  $D$  such that  $i^2 = j^2 = -1$  and  $ij = -ji$ . That is,  $D$  is isomorphic to  $\mathbf{H}$ .

An almost identical argument shows that if the dimension of  $D$  over its centre  $K$  is 4 and the characteristic is not 2, then  $D$  has a  $K$ -basis 1,  $i$ ,  $j$ ,  $ij$  where  $i^2 = \alpha$ ,  $j^2 = \beta$  and  $ij = -ji$  for some  $\alpha, \beta \in K$ .

#### 5. OTHER PROOFS OF WEDDERBURN'S THEOREM

The original proofs of the theorem of §3 were given first by Wedderburn [15] in 1905 and then by Dickson [5] in the same year; they depend on certain divisibility properties of the integers. The neatest proof along these lines is that of Witt [16]. Elementary proofs which avoid the use of such number theory have been given by Artin [1] and Herstein [7]. And