

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 20 (1974)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SOME CLASSICAL THEOREMS ON DIVISION RINGS
Autor: Taylor, D. E.
Kapitel: 5. Other proofs of Wedderburn's theorem
DOI: <https://doi.org/10.5169/seals-46912>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

number of conjugates of a subgroup is the index of its normalizer, so H has at most $|G : H|$ conjugates in G and hence the union of the conjugates contains at most $|G : H|(|H| - 1) + 1 = |G| - |G : H| + 1$ elements. This number is less than $|G|$ except when $G = H$. Hence $D = F$ is a field.

4. FROBENIUS' THEOREM

Let \mathbf{R} denote the field of real numbers, \mathbf{C} the field of complex numbers and \mathbf{H} the division ring of quaternions. The following proof makes use of the fundamental theorem that every polynomial with coefficients in \mathbf{C} has a root in \mathbf{C} .

THEOREM. *Let D be a division ring which contains the real numbers \mathbf{R} in its centre and suppose that every element of D satisfies a polynomial with coefficients in \mathbf{R} . Then D is isomorphic to one of \mathbf{R} , \mathbf{C} or \mathbf{H} .*

Proof. Suppose that D is not isomorphic to \mathbf{R} or \mathbf{C} . It follows that the maximal subfield F of D is isomorphic to \mathbf{C} , the centre K of D is isomorphic to \mathbf{R} and $F = K(i)$ where $i^2 = -1$. Let j be an eigenvector of T_i corresponding to the eigenvalue $-i$. Then $ji = -ij$ and j^2 commutes with j and F . From (2.2) and (2.3) the elements 1 and j form an F -basis for D and therefore $j^2 = \alpha$ belongs to K . If $\alpha = \beta^2$ for some $\beta \in K$ then $(j-\beta)(j+\beta) = 0$ and j belongs to K , which is not the case; hence $\alpha = -\beta^2$ for some $\beta \in K$. Replacing j by $j\beta^{-1}$ we obtain a K -basis 1, i , j , ij for D such that $i^2 = j^2 = -1$ and $ij = -ji$. That is, D is isomorphic to \mathbf{H} .

An almost identical argument shows that if the dimension of D over its centre K is 4 and the characteristic is not 2, then D has a K -basis 1, i , j , ij where $i^2 = \alpha$, $j^2 = \beta$ and $ij = -ji$ for some $\alpha, \beta \in K$.

5. OTHER PROOFS OF WEDDERBURN'S THEOREM

The original proofs of the theorem of §3 were given first by Wedderburn [15] in 1905 and then by Dickson [5] in the same year; they depend on certain divisibility properties of the integers. The neatest proof along these lines is that of Witt [16]. Elementary proofs which avoid the use of such number theory have been given by Artin [1] and Herstein [7]. And

proofs which deduce the theorem using finite group theory have been given by Zassenhaus [17], Brandis [3] and Scott [11, p. 426].

Perhaps the most interesting proofs are those which present the result as a consequence of a more general theory. There are two such proofs in the book of van der Waerden [14]: the first (on p. 203) uses the theory of central simple algebras, the second (sketched on p. 215) relates the theorem to cohomology and the Brauer group (see also, Serre [12, p. 170]). The theorem is also a consequence of the work of Tsen [13] and Chevalley [4]. Further comments on the history of the theorem can be found in an article by Artin [2] and in the book by Herstein [8] where many interesting generalisations are also given. One such generalization is a theorem of Jacobson: a division ring in which $x^{n(x)} = x$ for all x is commutative. Laffey [10] has recently given an elementary proof of this using Wedderburn's theorem and linear algebra similar to that used here. See also [18].

REFERENCES

- [1] ARTIN, E. Über einen Satz von Herrn J. H. MacLagan-Wedderburn. *Abh. Math. Sem. Univ. Hamburg*, 5 (1928), pp. 245-250.
- [2] —— The influence of J. H. M. Wedderburn on the development of modern algebra. *Bull. Amer. Math. Soc.*, 56 (1950), pp. 65-72.
- [3] BRANDIS, A. Ein gruppentheoretischer Beweis für die Kommutativität endlicher Divisionringe, *Abh. Math. Sem. Univ. Hamburg*, 26 (1964), pp. 234-236.
- [4] CHEVALLEY, C. Démonstration d'une hypothèse de M. Artin. *Abh. Math. Sem. Univ. Hamburg*, 11 (1936), pp. 73-75.
- [5] DICKSON, L. E. On finite algebras. *Nachr. Ges. Wiss. Göttingen*, (1905), p. 379.
- [6] FROBENIUS, G. Über lineare Substitutionen und bilineare Formen. *Crelle*, 84 (1878), p. 59.
- [7] HERSTEIN, I. N. Wedderburn's theorem and a theorem of Jacobson. *Amer. Math. Monthly*, 68 (1961), pp. 249-251.
- [8] —— *Noncommutative rings*. Carus Monograph 15, The Mathematical Society of America, 1968.
- [9] HOFFMAN K. and R. KUNZE. *Linear algebra*. Prentice-Hall, Englewood Cliffs, New Jersey, 1971.
- [10] LAFFEY, T. J. Infinite rings with all proper subrings finite. *Amer. Math. Monthly*, 81 (1974), pp. 270-272.
- [11] SCOTT, W. R. *Group theory*. Prentice-Hall, Englewood Cliffs, New Jersey, 1965.
- [12] SERRE, J.-P. *Corps locaux*. Hermann, Paris, 1968.
- [13] TSEN, C. C. Divisionalgebren über Funktionenkörpern. *Nachr. Ges. Wiss. Göttingen* (1933), p. 335.
- [14] van der WAERDEN, B. L. *Modern Algebra, vol. II*. Fredereick Ungar, New York, 1950.
- [15] MACLAGAN-WEDDERBURN, J. H. A theorem on finite algebras. *Trans. Amer. Math. Soc.*, 6 (1905), pp. 349-352.