

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 20 (1974)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: NON-STANDARD ANALYSIS: AN EXPOSITION
Autor: Levitz, Hilbert
Kapitel: 9. Derivatives
DOI: <https://doi.org/10.5169/seals-46892>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$f(c) \approx f(a_i^\alpha) \text{ and } f(c) \approx f(a_{i-1}^\alpha).$$

Taking this together with the fact (seen already) that

$$f(a_i^\alpha) \geq 0 \text{ and } f(a_i^\alpha) < 0$$

we have (in summary) that $f(c)$ is a standard number infinitely close to a negative number and a non-negative number. Thus $f(c) = 0$.

(Q.E.D.)

9. DERIVATIVES

Let $f(x)$ be a standard function defined on a standard open interval (a, b) and having the point x_0 as an interior point. Using the non-standard characterization of limit, the condition that $f(x)$ be differentiable at x_0 is that there exist a standard number L such that

$$\frac{f(x_0 + dx) - f(x_0)}{dx} \approx L$$

for all non-zero infinitesimals dx . L , of course, will be the derivative. If $f(x)$ is differentiable, then writing $dy = f(x_0 + dx) - f(x_0)$ we have

(using the notation for “standard part” introduced in Section 2) $\circ\left(\frac{dy}{dx}\right)$

$= f'(x_0)$. This says that the quotient of the infinitesimal increments need not in general be the derivative, but it must be infinitely close to it.

Example 9.1. Suppose we wish to calculate the derivative of $f(x) = x^2$. Let dx be an arbitrary non-zero infinitesimal, then

$$\frac{dy}{dx} = \frac{(x + dx)^2 - x^2}{dx}$$

After squaring and cancelling we get, $\frac{dy}{dx} = 2x + dx \approx 2x$ therefore

$$\circ\left(\frac{dy}{dx}\right) = 2x.$$

That is, the function x^2 is differentiable with derivative $2x$.

Example 9.2. Let's see how to prove the Chain Rule! Suppose $f(x)$ and $g(x)$ are differentiable at the appropriate places and we wish to show

that the function $h(x) = f(g(x))$ is differentiable with derivative $h'(x) = f'(g(x))g'(x)$. For any non-zero infinitesimal dx , write $dg = g(x+dx) - g(x)$ and $dh = h(x+dx) - h(x)$ then

$$dh = f(g(x+dx)) - f(g(x)) = f(g(x) + dg) - f(g(x)).$$

We want to show that for any non-zero infinitesimal dx ,

$$(1) \quad \frac{dh}{dx} \approx f'(g(x))g'(x).$$

Let non-zero infinitesimal dx be given. By continuity of $g(x)$, dg is also infinitesimal.

Case 1. $dg = 0$. Then $dh = 0$, so $\circ(\frac{dg}{dx}) = g'(x) = 0$ and $\frac{dh}{dx} = 0$. Thus

both sides of (1) are zero, so (1) holds.

Case 2. $dg \neq 0$. Then $\frac{dh}{dx} = \frac{dh}{dg} \cdot \frac{dg}{dx}$ that is

$$(2) \quad \frac{dh}{dx} = \frac{f(g(x)+dg) - f(g(x))}{dg} \cdot \frac{g(x+dx) - g(x)}{dx}.$$

The two factors of the right side of (2) are infinitely close to $f'(g(x))$ and $g'(x)$ respectively. Now using the rules given in Section 2 for manipulating the symbol \approx we get

$$\frac{dh}{dx} \approx f'(g(x)) \cdot g'(x)$$

as desired.

10. INTEGRATION

Let $f(x)$ be a standard function integrable on the standard interval $[a, b]$. For each standard n let

$$a = a_0^n < a_1^n < \cdots < a_n^n = b$$

be a partition of the interval into n subintervals of equal length. The Riemann sums

$$S_n = \sum_{i=1}^n f(a_i^n) (a_i^n - a_{i-1}^n)$$