

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 21 (1975)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: NOTES ON THE CONGRUENCE $y^2 \equiv x^5 - a \pmod{p}$
Autor: Rajwade, A. R.
Kapitel: 4. Expressions allied to $\Delta_a(\pi)$
DOI: <https://doi.org/10.5169/seals-47329>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Since the choice of g is arbitrary, we change g to another primitive root g^r with $(r, p-1) = 1$, $r = i \pmod{5}$, $i = 1, 2, 3, 4$. This does not alter Δ_a (as Δ_a is independent of g) but replaces π by any desired π_i so that $\Delta_a(\pi) = \Delta_a$ (any other π). Note that such an r exists, for all we want is, for $i = 1, 2, 3, 4$, a λ such that $(i+5\lambda, p-1) = 1$. Now $i+5\lambda$ takes infinitely many prime values as λ takes positive integer values since $(i, 5) = 1$; so λ may be chosen so that $i+5\lambda$ is a prime avoiding the primes occurring in $p-1$.

4. EXPRESSIONS ALLIED TO $\Delta_a(\pi)$

We fix our π now with $(g/\pi)_5 = \zeta$ and normalize it too. It is clear that there are only 3 expressions allied to $\Delta_a(\pi)$ viz $(-a/p)_Z (4a/\pi)_5 \cdot \pi \cdot \pi^\sigma +$ conjugates, $(-a/p)_Z (4a/\pi)_5 \cdot \pi^\sigma \cdot \pi^{\sigma^2} +$ conjugates and $(-a/p)_Z (4a/\pi)_5 \cdot \pi^{\sigma^2} \cdot \pi^{\sigma^3} +$ conjugates. This is so because changing the first term of $\Delta_a(\pi)$ fixes the changes in the other terms (otherwise we will not even get a rational integer!). Let us look at the first of these (the others would be similar), which equals $\text{Tr} [(-a/p)_Z (4a/\pi)_5 \cdot \pi \pi^\sigma]$. We have the following theorem:

THEOREM 3. $\text{Tr} [(-a/p)_Z (4a/\pi)_5 \cdot \pi \pi^\sigma] = \Delta_{au} - 1(\pi)$, where $(u/p)_Z = 1$ and $(u/\pi)_5 = (4a/\pi)_5$.

Proof. We have

$$\begin{aligned} \Delta_a(\pi) &= \text{Tr} [(-a/p)_Z (4a/\pi)_5 \cdot \pi \cdot \pi^{\sigma^3}] \\ &= \text{Tr} [(-a/p)_Z (4a/\pi^\sigma)_5 \cdot \pi^\sigma \cdot \pi^{\sigma^3}] \text{ by 3 on letting } \pi \rightarrow \pi^\sigma, \\ &= \text{Tr} [(-a/p)_Z (16a^2/\pi)_5 \cdot \pi^\sigma \cdot \pi] \text{ since } (4a/\pi^\sigma)_5 = (g^v/\pi_2)_5 \\ &= (g^v/\pi_1)_5^2 = (4a/\pi)_5^2 = (16a^2/\pi)_5, \\ &= \text{Tr} [(-au/p)_Z (4(au)/\pi)_5 \cdot \pi \pi^\sigma], \text{ where } (u/p)_Z = 1 \text{ and } (u/p)_5 \\ &= (4a/\pi)_5. \end{aligned}$$

Now writing a for au we get the theorem.

It follows that the expressions allied to $\Delta_a(\pi)$ also represent the number of solutions of the congruence (1) for a suitable value of a .

5. THE SET $\{\Delta_a \mid a = 1, 2, 3, \dots, p-1\}$

Dickson's paper on cyclotomy [1] includes the following Theorem (theorem 8 of [1]). Let $p \equiv 1 \pmod{5}$ be a rational prime. Then the Diophantine equations