

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 21 (1975)
Heft: 1: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON COMMON ZEROS OF EXPONENTIAL POLYNOMIALS
Autor: van der Poorten, A. J. / Tijdeman, R.
Kapitel: 1. Introduction
DOI: <https://doi.org/10.5169/seals-47330>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ON COMMON ZEROS OF EXPONENTIAL POLYNOMIALS

by A. J. van der POORTEN and R. TIJDEMAN

1. INTRODUCTION

At the 1974 Bolyai Janos Society Colloquium on Number Theory, H. L. Montgomery mentioned the following problem, which he attributed to H. S. Shapiro [13]:

Denote by E the collection of all exponential polynomials

$$(1) \quad E = \{a_1 e^{\alpha_1 z} + a_2 e^{\alpha_2 z} + \dots + a_n e^{\alpha_n z} : a_1, \dots, a_n \in \mathbf{C}, \alpha_1, \dots, \alpha_n \in \mathbf{C}, n \in \mathbf{N}\}$$

Suppose $f, g \in E$ have infinitely many zeros in common. Then is it the case that there exists an h in E , such that h has infinitely many zeros, and h is a common factor of f and g in the ring E ?

As we see below, it is equivalent to ask whether there exists an h in E , such that h has infinitely many zeros, and such that all the zeros of h are common zeros of f and g . Henceforth in this note we refer to Shapiro's problem simply as "the problem".

The problem is mentioned by H. S. Shapiro [13] in the context of his study of mean-periodic functions satisfying a certain functional equation. There, [13], p. 18 the problem appears in the form of a conjecture:

If two exponential polynomials have infinitely many zeros in common they are both multiples of some third (entire transcendental) exponential polynomial.

In this note we survey those ideas that appear relevant to settling this conjecture. Many of the ideas we mention here independently in response to Montgomery's question, are already alluded to in [13]. In particular we should remark that the conjecture arises as a generalisation of the Skolem-Mahler-Lech theorem which we describe in Section 4.

In the sequel we refer to the quantities $\alpha_1, \dots, \alpha_n$ in (1) as the *frequencies* of the exponential polynomial, and the quantities a_1, \dots, a_n as the *coefficients*. Unless otherwise indicated we shall always suppose given frequencies to be distinct and given coefficients to be non-zero. Similarly we shall suppose an exponential polynomial to have at least two distinct terms, hence to have a zero, and indeed hence to have infinitely many zeros. We mention some results on zeros of exponential polynomials in section 2.

As Professor Turán pointed out to us, one source of information on the problem is the papers of J. F. Ritt. Ritt provides a factorisation theory for exponential polynomials [10], and shows *inter alia* that if a quotient of exponential polynomials is an entire function then it is an exponential polynomial [11]. We describe these results in section 3.

In the case where all the frequencies of one of the exponential polynomials are rational we confirm that there is indeed a common factor. Even this special case seems to require a non-trivial argument; we employ the theorem of Skolem-Mahler-Lech on recurrence sequences with infinitely many vanishing terms (Lech [4], Mahler [6]). Conversely we observe in section 4 that an affirmative answer to the problem implies a generalised form of the Skolem-Mahler-Lech theorem. For a similar application of this theorem to zeros of exponential polynomials see Jager [3].

It follows from the results mentioned in sections 3 and 4 that one can define the greatest common divisor $h \in E$ of two exponential polynomials $f, g \in E$. An affirmative answer to the problem then implies that the set of zeros of the gcd h is all but at most finitely many of the common zeros of f and g . We make these and other remarks in section 5. We conclude this section with an example due to Montgomery which shows that “approximate methods” in the obvious manner are doomed to failure.

2. ZEROS OF EXPONENTIAL POLYNOMIALS

Given an exponential polynomial,

$$f(z) = \sum a_j e^{\alpha_j z} = a_1 e^{\alpha_1 z} + \dots + a_n e^{\alpha_n z}$$

denote by C_f the convex polygon in the complex plane defined by the complex conjugates of the frequencies; that is, the convex hull of the points $\bar{\alpha}_1, \bar{\alpha}_2, \dots, \bar{\alpha}_n$. Then *the zeros of f lie in half-strips in the directions of the exterior normals to C_f* . More quantitatively, *suppose an edge of the polygon C_f has length l . Then the number of zeros of $f(z)$ in the half-strip perpendicular to that edge and of absolute value less than R is*

$$(2) \quad \frac{lR}{2\pi} + O(1) \quad ; \text{ see Pólya [8], D. G. Dickson [2].}$$

It can also be shown that near every line in and parallel to the sides of a strip of zeros lie infinitely many zeros of the exponential polynomial, see Moreno [7], van der Poorten [9]. From a different point of view, one can