

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 21 (1975)  
**Heft:** 1: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON COMMON ZEROS OF EXPONENTIAL POLYNOMIALS  
**Autor:** van der Poorten, A. J. / Tijdeman, R.  
**Kapitel:** 2. Zeros of Exponential Polynomials  
**DOI:** <https://doi.org/10.5169/seals-47330>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

As Professor Turán pointed out to us, one source of information on the problem is the papers of J. F. Ritt. Ritt provides a factorisation theory for exponential polynomials [10], and shows *inter alia* that if a quotient of exponential polynomials is an entire function then it is an exponential polynomial [11]. We describe these results in section 3.

In the case where all the frequencies of one of the exponential polynomials are rational we confirm that there is indeed a common factor. Even this special case seems to require a non-trivial argument; we employ the theorem of Skolem-Mahler-Lech on recurrence sequences with infinitely many vanishing terms (Lech [4], Mahler [6]). Conversely we observe in section 4 that an affirmative answer to the problem implies a generalised form of the Skolem-Mahler-Lech theorem. For a similar application of this theorem to zeros of exponential polynomials see Jager [3].

It follows from the results mentioned in sections 3 and 4 that one can define the greatest common divisor  $h \in E$  of two exponential polynomials  $f, g \in E$ . An affirmative answer to the problem then implies that the set of zeros of the gcd  $h$  is all but at most finitely many of the common zeros of  $f$  and  $g$ . We make these and other remarks in section 5. We conclude this section with an example due to Montgomery which shows that “approximate methods” in the obvious manner are doomed to failure.

## 2. ZEROS OF EXPONENTIAL POLYNOMIALS

Given an exponential polynomial,

$$f(z) = \sum a_j e^{\alpha_j z} = a_1 e^{\alpha_1 z} + \dots + a_n e^{\alpha_n z}$$

denote by  $C_f$  the convex polygon in the complex plane defined by the complex conjugates of the frequencies; that is, the convex hull of the points  $\bar{\alpha}_1, \bar{\alpha}_2, \dots, \bar{\alpha}_n$ . Then *the zeros of  $f$  lie in half-strips in the directions of the exterior normals to  $C_f$ .* More quantitatively, suppose an edge of the polygon  $C_f$  has length  $l$ . Then the number of zeros of  $f(z)$  in the half-strip perpendicular to that edge and of absolute value less than  $R$  is

$$(2) \quad \frac{lR}{2\pi} + O(1) \quad ; \text{ see Pólya [8], D. G. Dickson [2].}$$

It can also be shown that near every line in and parallel to the sides of a strip of zeros lie infinitely many zeros of the exponential polynomial, see Moreno [7], van der Poorten [9]. From a different point of view, one can

obtain upper bounds for the number of zeros of  $f(z)$  in any circle of radius  $R$  in the complex plane; if  $\Delta = \max_k |\alpha_k|$ , then such a bound is

$$3(n-1) + 4R\Delta \quad ; \text{ see Tijdeman [17].}$$

For further details the reader is referred to the cited articles and the appropriate references mentioned therein.

### 3. FACTORISATION OF EXPONENTIAL POLYNOMIALS

We describe the results of J. F. Ritt [10], [11].

Define an ordering on the set of complex numbers by:  $\alpha < \beta$  if  $\Re \alpha \leq \Re \beta$ , and if  $\Re \alpha = \Re \beta$  then  $\Im \alpha < \Im \beta$ . We will suppose in the sequel, that, unless indicated otherwise, any exponential polynomial

$$\sum a_j e^{\alpha_j z} = a_1 e^{\alpha_1 z} + \dots + a_n e^{\alpha_n z}$$

is so normalised that  $a_1 = 1$  and  $0 = \alpha_1 < \alpha_2 < \dots < \alpha_n$ . Of course the normalisation is effected by multiplying by some unit  $be^{\beta z}$ ,  $b \neq 0$  thus not affecting the zeros of the exponential polynomial. Many of the remarks below are invalid if the normalisation is not assumed.

Ritt [10] firstly shows that *if the exponential polynomial  $\sum b_j e^{\beta_j z}$  divides the exponential polynomial  $\sum a_j e^{\alpha_j z}$  (in the ring  $E$  of exponential polynomials) then the frequencies  $\beta_1, \dots, \beta_m$  are linear combinations with rational coefficients of the frequencies  $\alpha_1, \dots, \alpha_n$ .* Now, following Ritt, call an exponential polynomial  $\sum a_j e^{\alpha_j z}$  simple if its frequencies are commensurable, that is, there is a minimal (in the sense of the ordering on  $\mathbb{C}$ ) number  $\alpha$  such that each  $\alpha_j$  is a non-negative integer multiple of  $\alpha$ . So such a simple exponential polynomial  $f(z)$  is a polynomial in  $e^{\alpha z}$  and factorises into a finite product of functions of the shape  $1 + ae^{\alpha z}$ ,  $a \in \mathbb{C}$ . Of course,  $1 + ae^{\alpha z}$  has factors of the shape  $1 + a'e^{(\alpha/m)z}$  for each  $m = 1, 2, 3, \dots$  but it follows from Ritt's lemma mentioned above that every factor of  $f(z)$  is a product of such factors. Similarly, call an exponential polynomial irreducible if it has no non-trivial (that is, other than units and associates) factors in the ring  $E$ . Then Ritt's principal result is that *an exponential polynomial can be factorised uniquely as a finite product of simple exponential polynomials such that their sets of frequencies are pairwise incommensurable, and a finite product of irreducible exponential polynomials.*

We outline the structure of the proof. Firstly one shows that there exist complex numbers  $\mu_1, \mu_2, \dots, \mu_p$  linearly independent over the field of