Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 22 (1976)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CLASSICAL THEOREMS ON QUADRATIC RESIDUES
Autor: Berndt, Bruce C.
Kapitel: 5. SUMS OVER INTERVALS OF LENGTH k/5.
DOI: https://doi.org/10.5169/seals-48188

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 14.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

ences for $h(-3p)$ are then consequences of (4.6). The number of summands in $S_{31}(\chi_{4p})$ is $8m + [4j/3]$. If $p \equiv 7 \pmod{12}$, the number of non-zero summands is $4m$; if $p \equiv 11 \pmod{12}$, the number of non-zero summands is $4m + 2$. In either case, the number of non-zero summands is even, and so it follows from (4.6) that $h(-12p) \equiv 0 \pmod 4$ when $p \equiv 3 \pmod 4$. Lastly, the number of summands in $S_{31}(\chi_{8p})$ is $16m + [8j/3]$. If $j = 1$, there are $8m$ non-zero summands; if $j = 5$, there are $8m + 6$ non-zero summands. In either case, $S_{31}(\chi_{8p})$ is even, and we deduce from (4.6) that $h(-24p) \equiv 0 \pmod 4$.

COROLLARY 4.5. Let p and q be distinct primes with $p, q > 3$ and $p \equiv q \pmod 4$. Then $h(-3pq) \equiv 0 \pmod 4$.

Proof. Let $p = 6m + j$ and $q = 6m' + j'$, where $j, j' = 1$ or 5 and m and m' are non-negative integers. The number of summands in $S_{31}(\chi_{pq})$ is $[pq/3]$, and we observe that $[pq/3] \equiv [jj'/3] \pmod 2$. Of these summands, $[q/3] = 2m' + [j'/3]$ are multiples of p, and $[p/3] = 2m + [j/3]$ are multiples of q. Thus,

$$S_{31}(\chi_{pq}) \equiv [jj'/3] - [j'/3] - [j/3] \pmod 2 .$$

By examining all of the possibilities for the pair j, j', we find that $S_{31}(\chi_{pq})$ is always even. The result now follows from (4.6).

It is clear that the same type of argument yields congruences from $h(-12pq)$ and $h(-24pq)$.

The class number formulae (4.6) and (4.7) appear to be due originally to Lerch [44, pp. 402, 408]. Holden [36] has also given a proof of (4.7).

5. SUMS OVER INTERVALS OF LENGTH $k/5$.

THEOREM 5.1. Let χ be odd and let $\chi_{5k}(n) = \left(\frac{n}{5}\right)\chi(n)$. Then

$$(5.1) \qquad S_{51} = \frac{1}{4\pi i} G(\chi)\{(5 - \bar{\chi}(5))\, L(1, \bar{\chi}) - 5^{1/2} L(1, \bar{\chi}_{5k})\}$$

and

$$(5.2) \qquad S_{52} = \frac{1}{2\pi i} 5^{1/2} G(\chi)\, L(1, \bar{\chi}_{5k}) .$$

Proof. Let

$$f(x) = \begin{cases} 1, & 0 < x < 2\pi/5\,, \\ 1/2, & x = 2\pi/5\,, \\ 0, & 2\pi/5 < x \leqslant \pi\,, \end{cases}$$

be an odd function of period 2π. Calculating the Fourier series of f, we find that, for all x,

$$(5.3) \qquad f(x) = \frac{2}{\pi}\sum_{n=1}^{\infty}\frac{\sin(nx)}{n} - \frac{2}{5\pi}\sum_{n=1}^{\infty}\frac{\sin(5nx)}{n}$$

$$+ \frac{2}{\pi}\cos(\pi/5)\sum_{\substack{n=1\\ n\equiv 2,3 \pmod 5}}^{\infty}\frac{\sin(nx)}{n} - \frac{2}{\pi}\cos(2\pi/5)\sum_{\substack{n=1\\ n\equiv 1,4 \pmod 5}}^{\infty}\frac{\sin(nx)}{n}$$

$$= \frac{2}{\pi}\sum_{n=1}^{\infty}\frac{\sin(nx)}{n} - \frac{2}{5\pi}\sum_{n=1}^{\infty}\frac{\sin(5nx)}{n}$$

$$+ \frac{1}{\pi}\cos(\pi/5)\sum_{n=1}^{\infty}\left\{1 - \left(\frac{n}{5}\right)\right\}\frac{\sin(nx)}{n}$$

$$- \frac{1}{\pi}\cos(2\pi/5)\sum_{n=1}^{\infty}\left\{1 + \left(\frac{n}{5}\right)\right\}\frac{\sin(nx)}{n}$$

$$+ \frac{1}{5\pi}\{\cos(2\pi/5) - \cos(\pi/5)\}\sum_{n=1}^{\infty}\frac{\sin(5nx)}{n}$$

$$= \frac{1}{2\pi}\sum_{n=1}^{\infty}\left\{5 - 5^{1/2}\left(\frac{n}{5}\right)\right\}\frac{\sin(nx)}{n} - \frac{1}{2\pi}\sum_{n=1}^{\infty}\frac{\sin(5n\ x)}{n}\,,$$

since $\cos(\pi/5) = (5^{1/2}+1)/4$ and $\cos(2\pi/5) = (5^{1/2}-1)/4$.

Now, multiply both sides of (2.1) by $\left\{5 - 5^{1/2}\left(\frac{n}{5}\right)\right\}/(2\pi n)$ and sum on n, $1 \leqq n < \infty$. Next, replace n by $5n$ in (2.1) and then multiply both sides of (2.1) by $-1/(2\pi n)$ and sum on n, $1 \leqq n < \infty$. Adding the resulting two equations and using (5.3), we get

$$2i\,S_{51} = i\sum_{j=1}^{k-1}\chi(j)f(2\pi j/k)$$

$$= \frac{G(\chi)}{2\pi}\left\{\sum_{n=1}^{\infty}\left\{5 - 5^{1/2}\left(\frac{n}{5}\right)\right\}\frac{\bar{\chi}(n)}{n} - \sum_{n=1}^{\infty}\frac{\bar{\chi}(5n)}{n}\right\}$$

$$= \frac{G(\chi)}{2\pi}\{5L(1,\bar{\chi}) - 5^{1/2}L(1,\bar{\chi}_{5k}) - \bar{\chi}(5)L(1,\bar{\chi})\}\,,$$

from which (5.1) follows immediately.

The proof of (5.2) is similar. In this case, we let

$$f(x) = \begin{cases} 0, & 0 \leqslant x < 2\pi/5,\ 4\pi/5 < x \leqslant \pi, \\ 1/2, & x = 2\pi/5,\ 4\pi/5, \\ 1, & 2\pi/5 < x < 4\pi/5, \end{cases}$$

be an odd function with period 2π. The Fourier series of f is given by

$$f(x) = \frac{5^{1/2}}{\pi} \sum_{n=1}^{\infty} \left(\frac{n}{5}\right) \frac{\sin(nx)}{n} \qquad (-\infty < x < \infty).$$

We then proceed in the same fashion as above.

COROLLARY 5.2. If χ is real and odd, then $S_{52} > 0$.

COROLLARY 5.3. If $d < 0$ and $5 \nmid d$, then

$$S_{51}(\chi_{-d}) = \frac{1}{4}\left\{5 - \left(\frac{d}{5}\right)\right\} h(d) - \frac{1}{4} h(5d) \tag{5.4}$$

and

$$S_{52}(\chi_{-d}) = \frac{1}{2} h(5d). \tag{5.5}$$

Formula (5.5) is due to Lerch [44, p. 407]. By combining (5.4) and (5.5), we can derive a formula for $h(d)$ which is also due to Lerch [44, p. 404].

COROLLARY 5.4. If $p \neq 5$, we have the following consequences:

(5.6) $h(-5p) \equiv 0 \pmod 8$, if $p \equiv 19 \pmod{20}$,

(5.7) $h(-5p) \equiv 4 \pmod 8$, if $p \equiv 11 \pmod{20}$,

(5.8) $h(-5p) \equiv 2h(-p) \pmod 8$, if $p \equiv 7 \pmod{20}$,

(5.9) $h(-5p) \equiv 4 + 2h(-p) \pmod 8$, if $p \equiv 3 \pmod{20}$,

(5.10) $h(-20p) \equiv 0 \pmod 8$, if $p \equiv 1,9 \pmod{20}$ or if $p \equiv 13,37 \pmod{40}$,

(5.11) $h(-20p) \equiv 4 \pmod 8$, if $p \equiv 17,33 \pmod{40}$,

(5.12) $h(-40p) \equiv 4 \pmod 8$, if $p \equiv 2,3 \pmod 5$,

and

(5.13) $h(-40p) \equiv 2h(-8p) \pmod 8$, if $p \equiv 1,4 \pmod 5$.

Proof. If $p \equiv j \pmod{10}$, $1 \leq j \leq 9$, then $S_{51}(\chi_p) \equiv [j/5] \pmod 2$. With the use of (5.4) and the above, and recalling that $h(-p)$ is odd, we deduce (5.6)-(5.9).

If $p \equiv j \pmod 5$, $1 \leq j \leq 4$, the number of non-zero summands in $S_{51}(\chi_{4p})$ is even if $j = 1$ or 4 and is odd if $j = 2$ or 3. Using also Corollary 3.10, we readily deduce (5.10) and (5.11) from (5.4).

If $p \equiv j \pmod 5$, $1 \leq j \leq 4$, the number of non-zero summands in $S_{51}(\chi_{8p})$ is even if $j = 1$ or 4 and is odd if $j = 2$ or 3. Using also the fact that $h(-8p)$ is even, we may deduce (5.12) and (5.13) from (5.4).

COROLLARY 5.5. Let p and q be primes with $p, q \neq 5$ and with $p \equiv q + 2 \pmod 4$. Then

$$h(-5pq) \equiv 0 \pmod 8, \text{ if } p \equiv 1,9 \pmod{20} \text{ and } q \equiv 11,19 \pmod{20},$$

$$h(-5pq) \equiv 4 \pmod 8, \text{ if } p \equiv 13,17 \pmod{20} \text{ and } q \equiv 3,7 \pmod{20},$$

and

$$h(-5pq) \equiv 2h(-pq), \text{ if } p \equiv 1,9 \pmod{20} \text{ and } q \equiv 3,7 \pmod{20},$$
$$\text{or if } p \equiv 13,17 \pmod{20} \text{ and } q = 11,19 \pmod{20}.$$

Of course, the same congruences for $h(-5pq)$ hold if the congruences for p and q are interchanged.

Proof. Let $p \equiv j \pmod{10}$ and $q \equiv j' \pmod{10}$, where $1 \leq j, j' \leq 9$. Observe that $S_{51}(\chi_{pq})$ contains $[pq/5]$ terms of which $[q/5]$ are multiples of p and $[p/5]$ are multiples of q. From (5.4), we then find that

$$4([jj'/5] - [j/5] - [j'/5])$$
$$\equiv \left\{5 - \left(\frac{5}{p}\right)\left(\frac{5}{q}\right)\right\} h(-pq) - h(-5pq) \pmod 8.$$

Since $h(-pq)$ is even, each of the desired congruences readily follows.

In the case that χ is even, we can state a theorem analogous to Theorem 5.1. However, the L-functions in the representations of S_{51} and S_{52} involve quartic characters. For example,

$$(5.14) \qquad S_{51} = \frac{G(\chi)}{\pi}\left\{\sin(2\pi/5) \sum_{\substack{n=1 \\ n \equiv 1,4 \pmod 5}}^{\infty} \frac{(-1)^{n+1}\bar{\chi}(n)}{n} + \sin(\pi/5) \sum_{\substack{n \equiv 1 \\ n \equiv 2,3 \pmod 5}}^{\infty} \frac{(-1)^n \bar{\chi}(n)}{n}\right\};$$

the series on the right side of (5.14) may be written in terms of L-functions of quartic characters. Thus, we are unable to derive any positivity results for character sums.

6. Sums over intervals of length $k/6$.

THEOREM 6.1. Let χ be even and let $\chi_{3k}(n) = \left(\frac{n}{3}\right)\chi(n)$. Then

$$S_{61} = \frac{3^{1/2}\,G(\chi)}{2\pi}\{1 + \bar{\chi}(2)\}\,L(1, \bar{\chi}_{3k}), \tag{6.1}$$

$$S_{62} = -\frac{3^{1/2}\,G(\chi)}{2\pi}\,\bar{\chi}(2)\,L(1, \bar{\chi}_{3k}), \tag{6.2}$$

and

$$S_{63} = -\frac{3^{1/2}\,G(\chi)}{2\pi}\,L(1, \bar{\chi}_{3k}). \tag{6.3}$$

Let χ be odd. Then

$$S_{61} = \frac{G(\chi)}{2\pi i}\{1 + \bar{\chi}(2) + \bar{\chi}(3) - \bar{\chi}(6)\}\,L(1, \bar{\chi}), \tag{6.4}$$

$$S_{62} = \frac{G(\chi)}{2\pi i}\{2 - \bar{\chi}(2) - 2\bar{\chi}(3) + \bar{\chi}(6)\}\,L(1, \bar{\chi}), \tag{6.5}$$

and

$$S_{63} = \frac{G(\chi)}{2\pi i}\{1 - 2\bar{\chi}(2) + \bar{\chi}(3)\}\,L(1, \bar{\chi}). \tag{6.6}$$

We shall not give a proof of Theorem 6.1, because all of the formulas may be deduced from Theorems 3.2 and 4.1 and elementary considerations.

COROLLARY 6.2. If $d > 0$, we have

$S_{61} > 0$, if d is even, or if $\chi(2) = 1$;

$S_{61} = 0$, if $\chi(2) = -1$;

$S_{62} > 0$, if $\chi(2) = -1$;

$S_{62} = 0$, if d is even;

$S_{62} < 0$, if $\chi(2) = 1$;

$S_{63} < 0$, for all d;