

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	22 (1976)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	CLASSICAL THEOREMS ON QUADRATIC RESIDUES
Autor:	Berndt, Bruce C.
Kapitel:	12. Sums over several intervals of equal length
DOI:	https://doi.org/10.5169/seals-48188

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Proof. The desired congruences follow from Corollaries 7.5, 9.3, and 11.6.

Lerch [44, pp. 409, 410] has derived some class number formulas in terms of the sums $S_{24,i}$, $1 \leq i \leq 12$. Karpinski [42] and Rédei [57] have also established class number relations of this sort.

12. SUMS OVER SEVERAL INTERVALS OF EQUAL LENGTH

In this section, it will be convenient to use the following character analogues of the Poisson summation formula [6, Theorem 2.3], [7, equations (4.1), (4.2)]. Let f be continuous and of bounded variation on $[c, d]$. Let χ be a primitive character of modulus k . If χ is even, then

$$(12.1) \quad \sum'_{c \leq n \leq d} \chi(n) f(n) = \frac{2G(\chi)}{k} \sum_{n=1}^{\infty} \bar{\chi}(n) \int_c^d f(x) \cos(2\pi nx/k) dx;$$

if χ is odd, then

$$(12.2) \quad \sum'_{c \leq n \leq d} \chi(n) f(n) = -\frac{2iG(\chi)}{k} \sum_{n=1}^{\infty} \bar{\chi}(n) \int_c^d f(x) \sin(2\pi nx/k) dx.$$

The primes ' on the summation signs on the left sides of (12.1) and (12.2) indicate that if c or d is an integer, then the associated summands must be halved.

Throughout the section, it is assumed that χ is a primitive character of modulus k . For each of the theorems below, deductions concerning the signs of the pertinent character sums are trivial. Likewise, the corresponding formulas for class numbers are immediate from (2.4). Thus, none of these obvious corollaries shall be explicitly stated.

THEOREM 12.1. Let χ be even, and let m be any positive integer. Then

$$(12.3) \quad S_{4m,1} + S_{4m,4} + S_{4m,5} + S_{4m,8} + S_{4m,9} + \dots + S_{4m,4m} \\ = \frac{2G(\chi)}{\pi} \bar{\chi}(m) L(1, \bar{\chi}_{4k}).$$

Proof. Apply (12.1) several times with $f(x) \equiv 1$ in each case and with $(c, d) = (0, k/4m)$, $(3k/4m, 5k/4m)$, $(7k/4m, 9k/4m)$, ..., $((4m-1)k/4m, k)$. We then get

$$\begin{aligned}
 S_{4m,1} &= \frac{G(\chi)}{\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \sin(2\pi n/4m), \\
 S_{4m,4} + S_{4m,5} &= \frac{G(\chi)}{\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \{ \sin(10\pi n/4m) - \sin(6\pi n/4m) \} \\
 &\quad \cdot \\
 &\quad \cdot \\
 S_{4m,4m} &= \frac{G(\chi)}{\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \{ -\sin(2\pi n(4m-1)/4m) \}.
 \end{aligned}$$

Adding the above equations, we find that

$$\begin{aligned}
 (12.4) \quad &S_{4m,1} + S_{4m,4} + S_{4m,5} + \dots + S_{4m,4m} \\
 &= \frac{G(\chi)}{\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \sum_{j=0}^{2m-1} (-1)^j \sin(2(2j+1)\pi n/4m).
 \end{aligned}$$

Now an elementary calculation shows that

$$\begin{aligned}
 (12.5) \quad &\sum_{j=0}^{2m-1} (-1)^j \sin(2(2j+1)\pi n/4m) \\
 &= \begin{cases} 2m(-1)^{\mu}, & \text{if } n = (2\mu+1)m, \\ 0, & \text{otherwise.} \end{cases}
 \end{aligned}$$

Putting (12.5) into (12.4), we conclude that

$$\begin{aligned}
 &S_{4m,1} + S_{4m,4} + S_{4m,5} + \dots + S_{4m,4m} \\
 &= \frac{2G(\chi)}{\pi} \sum_{\mu=0}^{\infty} \frac{(-1)^{\mu} \bar{\chi}((2\mu+1)m)}{2\mu+1} = \frac{2G(\chi)}{\pi} \bar{\chi}(m) L(1, \bar{\chi}_{4k}),
 \end{aligned}$$

which completes the proof.

Observe that if $m = 1$, Theorem 12.1 reduces to Theorem 3.7. If $m = 2, 3, 4$, and 6 , then Theorem 12.1 reduces to results that can be derived from Theorems 7.1, 9.1, 10.1, and 11.1, respectively.

THEOREM 12.2. Let χ be odd, and let m be a positive integer. If m is odd, then

$$(12.6) \quad \sum_{1 \leq j \leq m/2} \left(\frac{m+1}{2} - j \right) S_{m,j} = \frac{G(\chi)}{2\pi i} \{ m - \bar{\chi}(m) \} L(1, \bar{\chi});$$

if m is even, then

$$(12.7) \quad \sum_{1 \leq j \leq m/2} \left(\frac{m+2}{2} - j \right) S_{m,j} = \frac{G(\chi)}{2\pi i} \{ m+2 - \bar{\chi}(2) - \bar{\chi}(m) \} L(1, \bar{\chi}).$$

Proof. Apply (12.2) several times with $f(x) \equiv 1$ in each case and with $(c, d) = (0, k/m), (k/m, 2k/m), \dots, (([m/2]-1)k/m, [m/2]k/m)$. We then get

$$\begin{aligned} S_{m,1} &= \frac{iG(\chi)}{\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \left\{ \cos(2\pi n/m) - 1 \right\}, \\ S_{m,2} &= \frac{iG(\chi)}{\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \left\{ \cos(4\pi n/m) - \cos(2\pi n/m) \right\}, \\ &\quad \vdots \\ &\quad \vdots \\ S_{m,[m/2]} &= \frac{iG(\chi)}{\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \left\{ \cos(2[m/2]\pi n/m) \right. \\ &\quad \left. - \cos(2\{[m/2]-1\}\pi n/m) \right\}. \end{aligned}$$

Multiply the j -th equation above by $[m/2] + 1 - j$, $1 \leq j \leq [m/2]$, and add the resulting equations to obtain

$$(12.8) \quad \begin{aligned} &\sum_{1 \leq j \leq m/2} \{[m/2] + 1 - j\} S_{m,j} \\ &= \frac{iG(\chi)}{\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \left\{ -[m/2] + \sum_{j=1}^{[m/2]} \cos(2\pi nj/m) \right\}. \end{aligned}$$

First, suppose that m is odd. Then (12.8) becomes

$$\begin{aligned} \sum_{1 \leq j \leq m/2} \left(\frac{m+1}{2} - j \right) S_{m,j} &= \frac{iG(\chi)}{2\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \left\{ -m + \sum_{j=0}^{m-1} \cos(2\pi nj/m) \right\} \\ &= \frac{iG(\chi)}{2\pi} \{ -m + \bar{\chi}(m) \} L(1, \bar{\chi}), \end{aligned}$$

from which (12.6) follows.

Suppose next that m is even. Then (12.8) becomes

$$\begin{aligned} \sum_{1 \leq j \leq m/2} \left(\frac{m+2}{2} - j \right) S_{m,j} &= \frac{iG(\chi)}{2\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \left\{ -m - 1 + (-1)^n + \sum_{j=0}^{m-1} \cos(2\pi nj/m) \right\} \\ &= \frac{iG(\chi)}{2\pi} \{ -m - 1 + \bar{\chi}(2) - 1 + \bar{\chi}(m) \} L(1, \bar{\chi}), \end{aligned}$$

from which (12.7) follows.

We indicate some special cases of the previous theorem. If $m = 2$, (12.7) reduces to Theorem 3.2. If $m = 3$, (12.6) yields Theorem 4.1. If $m = 5, 6, 8, 10, 12$, and 24 in Theorem 12.2, we obtain results deducible from Theorems 5.1, 6.1, 7.1, 8.1, 9.1 and 11.1, respectively.

THEOREM 12.3. Let χ be even and let m be an arbitrary positive integer. Then

$$(12.9) \quad S_{8m,1} - S_{8m,4} - S_{8m,5} + S_{8m,8} + S_{8m,9} - \dots + \dots + S_{8m,8m} \\ = \frac{2^{3/2} G(\chi)}{\pi} \bar{\chi}(m) L(1, \bar{\chi}_{8k}).$$

Proof. Apply (12.1) several times with $f(x) \equiv 1$ in each instance and with $(c, d) = (0, k/8m), (3k/8m, 5k/8m), (7k/8m, 9k/8m), \dots, ((8m-1)k/8m, k)$. Accordingly, we find that

$$S_{8m,1} - S_{8m,4} - S_{8m,5} + S_{8m,8} + S_{8m,9} - \dots + \dots + S_{8m,8m} \\ = \frac{G(\chi)}{\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \sum_{j=0}^{8m-1} \chi_4(j) \chi_8(j) \sin(2\pi nj/8m) \\ = \frac{G(\chi)}{\pi} \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \sum_{v=0}^7 \chi_4(v) \chi_8(v) \sum_{\mu=0}^{m-1} \sin(2\pi n(8\mu+v)/8m) \\ = \frac{G(\chi)}{\pi} \bar{\chi}(m) \sum_{n=1}^{\infty} \frac{\bar{\chi}(n)}{n} \sum_{v=0}^7 \chi_4(v) \chi_8(v) \sin(2\pi nv/8).$$

The inner sum above is merely $-iG(n, \chi_4 \chi_8) = \chi_4(n) \chi_8(n) 2^{3/2}$, by (2.2). Hence, (12.9) immediately follows.

The special cases with $m = 1, 2$ and 3 of Theorem 12.3 may be deduced from Theorems 7.1, 10.1 and 11.1, respectively.

The proofs of the next four theorems are very similar to the preceding proofs and so will not be given.

THEOREM 12.4. Let χ be odd, and let m be an arbitrary positive integer. Then

$$S_{8m,2} + S_{8m,3} - S_{8m,6} - S_{8m,7} + \dots - S_{8m,8m-2} - S_{8m,8m-1} \\ = - \frac{i2^{3/2} G(\chi)}{\pi} \bar{\chi}(m) L(1, \bar{\chi}_{8k}).$$

The special cases of Theorem 12.4 with $m = 1, 2$ and 3 are consequences of Theorems 7.1, 10.1 and 11.1, respectively.

THEOREM 12.5. Let χ be even, and let m be an arbitrary positive integer.

Then

$$\sum_{j=0}^{m-1} S_{3m,3j+2} = -\frac{3^{1/2} G(\chi)}{\pi} \bar{\chi}(m) L(1, \bar{\chi}_{3k}).$$

The instances of Theorem 12.5 with $m = 1, 2, 4$ and 8 are consequences of Theorems 4.1, 6.1, 9.1 and 11.1, respectively.

THEOREM 12.6. Let χ be odd, and let m be an arbitrary positive integer.

Then

$$S_{5m,2} - S_{5m,4} + S_{5m,7} - S_{5m,9} + \cdots + S_{5m,5m-3} - S_{5m,5m-1} \\ = -\frac{i 5^{1/2} G(\chi)}{\pi} \bar{\chi}(m) L(1, \bar{\chi}_{5k}).$$

The special cases of Theorem 12.6 for $m = 1$ and $m = 2$ follow immediately from Theorems 5.1 and 8.1, respectively.

THEOREM 12.7. Let χ be odd, and let m be an arbitrary positive integer.

Then

$$S_{12m,2} + S_{12m,3} + S_{12m,4} + S_{12m,5} - S_{12m,8} - S_{12m,9} - S_{12m,10} - S_{12m,11} \\ + + + + - - - - \cdots - S_{12m,12m-4} - S_{12m,12m-3} - S_{12m,12m-2} - S_{12m,12m-1} \\ = -\frac{i (12)^{1/2} G(\chi)}{\pi} \bar{\chi}(m) L(1, \bar{\chi}_{12k}).$$

The special instances of $m = 1$ and $m = 2$ of Theorem 12.7 yield results that are easily deduced from Theorems 9.1 and 11.1, respectively.

The class number formula arising from Theorem 12.1 was first proved by Holden [39]. A less general form of Theorem 12.2 was also established by Holden [36] who in another paper [37] used his result to derive formulas for sums of the Legendre-Jacobi symbol over various residue classes. The special case $m = 1$ of the class number formula deducible from Theorem 12.7 is due to Lerch [44, p. 407]. Otherwise, the results of this section appear to be new.

13. SUMS OF QUADRATIC RESIDUES AND NONRESIDUES

We mentioned in the Introduction the two equivalent formulations of Dirichlet's theorem for primes that are congruent to 3 modulo 4. In this section, we state and prove as many theorems as we can that are of the same